

**American College of Radiology  
ACR Appropriateness Criteria®  
Postmenopausal Subacute or Chronic Pelvic Pain**

**Variant 1: Postmenopausal subacute or chronic pelvic pain, localized to the deep pelvis. Initial imaging.**

Procedure	Appropriateness Category	Relative Radiation Level
US pelvis transvaginal	Usually Appropriate	○
US duplex Doppler pelvis	Usually Appropriate	○
US pelvis transabdominal	Usually Appropriate	○
MRI pelvis without and with IV contrast	May Be Appropriate	○
CT abdomen and pelvis with IV contrast	May Be Appropriate	☼ ☼ ☼
CT pelvis with IV contrast	May Be Appropriate	☼ ☼ ☼
MRI pelvis without IV contrast	May Be Appropriate (Disagreement)	○
CT abdomen and pelvis without and with IV contrast	Usually Not Appropriate	☼ ☼ ☼ ☼
CT abdomen and pelvis without IV contrast	Usually Not Appropriate	☼ ☼ ☼
CT pelvis without and with IV contrast	Usually Not Appropriate	☼ ☼ ☼ ☼
CT pelvis without IV contrast	Usually Not Appropriate	☼ ☼ ☼
Radiography abdomen and pelvis	Usually Not Appropriate	☼ ☼ ☼

**Variant 2: Postmenopausal subacute or chronic pelvic pain, clinically suspected pathologies in perineum, vulva, or vagina. Initial imaging.**

Procedure	Appropriateness Category	Relative Radiation Level
US duplex Doppler pelvis	Usually Appropriate	○
US pelvis transabdominal	Usually Appropriate	○
US pelvis transvaginal	Usually Appropriate	○
MRI pelvis without and with IV contrast	May Be Appropriate	○
MRI pelvis without IV contrast	May Be Appropriate (Disagreement)	○
CT abdomen and pelvis with IV contrast	Usually Not Appropriate	☼ ☼ ☼
CT abdomen and pelvis without and with IV contrast	Usually Not Appropriate	☼ ☼ ☼ ☼
CT abdomen and pelvis without IV contrast	Usually Not Appropriate	☼ ☼ ☼
CT pelvis with IV contrast	Usually Not Appropriate	☼ ☼ ☼
CT pelvis without and with IV contrast	Usually Not Appropriate	☼ ☼ ☼ ☼
CT pelvis without IV contrast	Usually Not Appropriate	☼ ☼ ☼
Radiography abdomen and pelvis	Usually Not Appropriate	☼ ☼ ☼

## POSTMENOPAUSAL SUBACUTE OR CHRONIC PELVIC PAIN

Expert Panel on Women's Imaging: Katherine E. Maturen, MD, MS<sup>a</sup>; Esma A. Akin, MD<sup>b</sup>; Mark Dassel, MD<sup>c</sup>; Sandeep Prakash Deshmukh, MD<sup>d</sup>; Kika M. Dudiak, MD<sup>e</sup>; Tara L. Henrichsen, MD<sup>f</sup>; Lee A. Learman, MD, PhD<sup>g</sup>; Edward R. Oliver, MD, PhD<sup>h</sup>; Liina Poder, MD<sup>i</sup>; Elizabeth A. Sadowski, MD<sup>j</sup>; Hebert Alberto Vargas, MD<sup>k</sup>; Therese M. Weber, MD<sup>l</sup>; Tom Winter, MD<sup>m</sup>; Phyllis Glanc, MD.<sup>n</sup>

### **Summary of Literature Review**

#### **Introduction/Background**

Chronic pelvic pain, defined as cyclical or noncyclical pain involving the pelvis, lower abdomen, vulva, vagina, or perineum and lasting for at least 6 months, affects as many as a quarter of women worldwide and is the single most common presenting complaint at gynecologic office visits [1,2]. The morbidity, public health impact, and downstream costs are substantial but poorly quantified in part due to the large variety of etiologies and lack of definitions associated with chronic pelvic pain. For purposes of this document, the term “subacute” is added to distinguish our target entities from diagnoses that most commonly present with acute or even emergent symptoms. This guideline is limited to postmenopausal women, which further limits the range of potential pain etiologies.

Subacute or chronic pelvic pain is a broad clinical presentation common to a variety of gynecologic, urinary, gastrointestinal, and musculoskeletal disorders. There are specific ACR Appropriateness Criteria documents pertaining to many of these diagnoses, which are detailed in [Appendix 1](#). In particular, we emphasize the importance of both vaginal bleeding and suspected adnexal mass in postmenopausal women because of the prevalence of endometrial and ovarian neoplasia in this age group. These clinical features, if present, should take precedence over the general complaint of pelvic pain in directing the management algorithm. Patients with acute pain, suspected pelvic floor dysfunction, or urinary complaints may be managed in accordance with the respective algorithms for those conditions. Imaging evaluation for suspected endometriosis is not considered here as endometriosis is estrogen dependent and usually regresses after menopause [3]. If a postmenopausal woman is experiencing pain from endometriosis, it is likely secondary to scarring or reactivation that is due to postmenopausal hormonal therapy. In cases of persistent endometriosis-related symptoms after menopause, readers are referred to ACR Appropriateness Criteria guidance for the premenopausal age group (see [Appendix 1](#)).

Finally, like other types of chronic pain, pelvic pain is a complex process with incompletely mapped cognitive and neurologic contributors. As such, there is a growing body of literature regarding potential use of neurologic imaging in patients with chronic pelvic pain [4-7]. However, central nervous system functional imaging remains in the research domain for evaluation of chronic pelvic pain at this time, so we will not consider it formally among the discussed imaging procedures.

When all of these aspects of subacute and chronic pelvic pain in postmenopausal women are excluded from direct consideration, a handful of clinically significant conditions remain. We group these according to location of clinical symptoms: pain localized to the deep or internal pelvis, with potential etiologies and associated conditions, including pelvic venous disorders (commonly termed pelvic congestion syndrome), intraperitoneal adhesions, hydrosalpinx, chronic inflammatory disease, or cervical stenosis versus chronic pain localized to the perineum, vulva, or vagina that arises from suspected vaginal atrophy, vaginismus, vaginal or vulvar cysts, vulvodynia, or pelvic myofascial pain.

#### **Special Imaging Considerations**

When there is suspected local pathology in the vulva, perineum, or vaginal wall, translabial/transperineal ultrasound (US) or side-firing transvaginal probes may provide better visualization than end-firing transvaginal

<sup>a</sup>Panel Chair, University of Michigan, Ann Arbor, Michigan. <sup>b</sup>George Washington University Hospital, Washington, District of Columbia. <sup>c</sup>Cleveland Clinic, Cleveland, Ohio; American Congress of Obstetricians and Gynecologists. <sup>d</sup>Thomas Jefferson University Hospital, Philadelphia, Pennsylvania. <sup>e</sup>Mayo Clinic, Rochester, Minnesota. <sup>f</sup>Mayo Clinic, Rochester, Minnesota. <sup>g</sup>Florida Atlantic University, Boca Raton, Florida; American Congress of Obstetricians and Gynecologists. <sup>h</sup>Children's Hospital of Philadelphia and Perelman School of Medicine at the University of Pennsylvania, Philadelphia, Pennsylvania. <sup>i</sup>University of California San Francisco, San Francisco, California. <sup>j</sup>University of Wisconsin, Madison, Wisconsin. <sup>k</sup>Memorial Sloan Kettering Cancer Center, New York, New York. <sup>l</sup>University of Alabama at Birmingham, Birmingham, Alabama. <sup>m</sup>University of Utah, Salt Lake City, Utah. <sup>n</sup>Specialty Chair, University of Toronto and Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada.

The American College of Radiology seeks and encourages collaboration with other organizations on the development of the ACR Appropriateness Criteria through society representation on expert panels. Participation by representatives from collaborating societies on the expert panel does not necessarily imply individual or society endorsement of the final document.

Reprint requests to: [publications@acr.org](mailto:publications@acr.org)

US probes [8]. There is scant evidence for imaging recommendations at this level of specificity, and it is assumed that the performing sonographer and sonologist will make appropriate technical adjustments to optimize imaging in these relatively uncommon clinical scenarios.

## **Discussion of Procedures by Variant**

### **Variant 1: Postmenopausal subacute or chronic pelvic pain, localized to the deep pelvis. Initial imaging.**

#### **Radiography Abdomen and Pelvis**

To our knowledge, there is currently no evidence to support the use of radiography to evaluate postmenopausal subacute or chronic pelvic pain localized to the deep pelvis.

#### **US Pelvis Transvaginal**

Pelvic US using a combined transabdominal and transvaginal approach is the initial imaging study of choice to evaluate postmenopausal subacute or chronic pelvic pain localized to the deep pelvis [9-12]. US can provide anatomic information about uterine size and endometrial canal distension, fallopian tube dilation, ovaries, and adnexal masses. Regarding the sequencing of examinations, several authors have pointed out that if the etiology of pelvic pain remains obscure after CT, a subsequent US has the capacity to provide additional information about the adnexa in particular [13,14]. US is broadly used and clinically accepted worldwide. However, high-quality evidence, such as clinical trials supporting specific usefulness of US, is lacking.

Chronic pelvic inflammatory disease may be associated with pelvic fluid, hydrosalpinx or pyosalpinx, inflammatory adnexal masses, and peritoneal inclusions visible by US [15]. When pelvic adhesions are suspected, real-time dynamic US or cine clips may document abnormal adherence or lack of mobility of structures, particularly transvaginally. However, adhesive disease is a notoriously difficult diagnosis to confirm nonoperatively [16], and the evidence basis is anecdotal [15,17]. Furthermore, the causal linkage between adhesive disease and chronic pelvic pain remains unclear.

#### **US Pelvis Transabdominal**

As above, a combined transabdominal and transvaginal approach is most appropriate for pelvic imaging, combining the anatomic overview provided by the transabdominal approach with the greater spatial and contrast resolution of transvaginal imaging. These techniques should be performed together whenever possible. Please see the “US Pelvis Transvaginal” section for further details.

#### **US Duplex Doppler Pelvis**

Color and spectral Doppler are routinely employed in pelvic sonography to evaluate internal vascularity of pelvic observations and distinguish fluid and cysts from soft tissue. Although it is rated as a separate imaging procedure per ACR methodology, the expert panel considers Doppler imaging to be a standard component of pelvic sonography. Special considerations for women with chronic pelvic pain may include evaluation of uterine artery blood flow with low-resistance waveforms in women with chronic pelvic pain [12] and altered venous flow in the setting of pelvic congestion. When pelvic venous disorders are suspected clinically, color and spectral Doppler evaluation may be used to document engorged periuterine and periovarian veins ( $\geq 8$  mm), low-velocity flow, altered flow with Valsalva maneuver, retrograde (caudal) flow of the ovarian veins, and direct connection between engorged pelvic veins and myometrial arcuate veins [9-11]. Increased pelvic vascularity may also be present in the setting of uterine or tubo-ovarian neoplasia; these guidelines assume normal imaging appearance of the pelvic organs.

Many women with pelvic venous disorders have morphologic findings of polycystic ovarian syndrome (enlarged ovaries with exaggerated central stroma and multiple small peripherally located follicles), but the associated clinical features of hirsutism and amenorrhea are rare [11,18]. Multiple investigators have identified a component of estrogen overstimulation in pelvic venous disorders, and symptoms may subside after menopause in some women [15]. There is a lack of clear definition and high-quality evidence in the clinical domain of pelvic venous disorders.

#### **CT Abdomen and Pelvis**

When pelvic venous disorders are clinically suspected, contrast-enhanced CT of the abdomen and pelvis may demonstrate engorged periuterine and periovarian veins, venous anatomic variants, and occasional compression of the left renal vein resulting in asymmetric left-sided pelvic varicosities [10,19-22]. However, CT lacks the capacity of US or MR to provide dynamic flow information [18].

In chronic pelvic inflammatory disease, CT may demonstrate pelvic fluid, peritoneal thickening, hydrosalpinx or pyosalpinx, and even tubo-ovarian abscess [23]. CT has the capacity to demonstrate architectural distortion and tethering in adhesive disease, but CT sensitivity and specificity for this diagnosis have not been documented to our knowledge. When adhesive disease is severe, small-bowel obstruction may result, and CT of the abdomen and pelvis with intravenous (IV) contrast is the imaging examination of choice. See [Appendix 1](#) for the ACR Appropriateness Criteria® for “Suspected Small-Bowel Obstruction.”

### **CT Pelvis**

When pelvic venous disorders are clinically suspected, contrast-enhanced CT of the pelvis may demonstrate engorged periuterine and periovarian veins, although their drainage into the renal vein or cava will not be evaluated without CT coverage of the abdomen [10,19-22]. In chronic inflammatory disease, CT may demonstrate pelvic fluid, peritoneal thickening, hydrosalpinx or pyosalpinx, and tubo-ovarian abscess [23].

### **MRI Pelvis**

MRI is widely regarded as the problem-solving imaging examination of choice for chronic pelvic pain, particularly when US findings are nondiagnostic or inconclusive [11,15,24]. When MRI is clinically indicated, the use of a gadolinium-based IV contrast agent is preferred. Please see the ACR Manual on Contrast Media for additional information [25].

The diagnostic performance of MRI/MR angiography is comparable to conventional venography for identifying pelvic venous disorders [26,27]. The use of MRI for this indication is growing accordingly [28], necessitating standardized interpretation and reporting [29]. T2-weighted imaging has the capacity to demonstrate pelvic varices, but signal intensity varies with flow velocity. Vein conspicuity and flow directional assessment are superior using time-resolved postcontrast T1-weighted imaging, which can directly demonstrate ovarian vein reflux [10,30]. Noninvasive imaging with MRI has largely supplanted conventional venography for diagnostic purposes, but venography may still be performed in the context of intended intervention.

In chronic pelvic inflammatory disease, MRI with T2-weighted imaging may demonstrate edema, fluid collections, and distension of endometrial canal or fallopian tubes [15]. When infection is long standing, distinguishing between inflammatory and neoplastic masses is particularly difficult. Postcontrast T1-weighted imaging and diffusion-weighted imaging are particularly important in this setting [23]. Adhesive disease may be directly evident at MRI as low-signal bands between structures on nonfat saturated T2-weighted imaging or inferred in the presence of peritoneal inclusion cysts [11,15].

### **Variant 2: Postmenopausal subacute or chronic pelvic pain, clinically suspected pathologies in perineum, vulva, or vagina. Initial imaging.**

Physical examination is the foundation of clinical evaluation of suspected pathology in the perineum, vulva, or vagina. The evidence supporting the use of imaging procedures in this clinical context largely assumes that the physical examination is abnormal.

### **Radiography Abdomen and Pelvis**

To our knowledge, there is currently no evidence to support the use of radiography to evaluate postmenopausal subacute or chronic pelvic pain localized to the perineum, vulva, or vagina.

### **US Pelvis Transvaginal**

Physical examination is the basis of diagnosis for most conditions localized to the vulvar skin [31]. Perineal and vaginal cysts are subcutaneous but often palpable and are appropriately evaluated with either translabial or transvaginal US, or both [8]. As with pelvic pain localized to the deep pelvis, US is widely regarded as the initial imaging study of choice for pelvic pain localized to the perineum, vulva, or vagina, but there is little high-quality evidence specifically supporting its use.

### **US Pelvis Transabdominal**

As above, a combined transabdominal and transvaginal approach is most appropriate for pelvic imaging, combining the anatomic overview provided by the transabdominal approach with the greater spatial and contrast resolution of transvaginal imaging. These techniques should be performed together whenever possible. Please see the “US Pelvis Transvaginal” section for further details.

## US Duplex Doppler Pelvis

Color and spectral Doppler are routinely used in pelvic sonography to evaluate internal vascularity of pelvic observations and distinguish cysts from soft tissue. Although it is rated as a separate imaging procedure per ACR methodology, the expert panel considers Doppler imaging to be a standard component of pelvic sonography. Special considerations for women with chronic pelvic pain may include evaluation of uterine artery blood flow, with low-resistance waveforms having been described in women with chronic pelvic pain [12].

## CT Abdomen and Pelvis

To our knowledge, there is currently no evidence to support the use of CT for primary evaluation of postmenopausal subacute or chronic pelvic pain localized to the perineum, vulva, or vagina.

## CT Pelvis

To our knowledge, there is currently no evidence to support the use of CT for primary evaluation of postmenopausal subacute or chronic pelvic pain localized to the perineum, vulva, or vagina.

## MRI Pelvis

When MRI is clinically indicated, the use of a gadolinium-based IV contrast agent is preferred. Please see the ACR Manual on Contrast Media for additional information [25].

When a cyst or mass is identified by US in the perineum, vulva, or vagina, MRI provides additional anatomic detail and evaluation of any enhancing soft-tissue components that might favor infection or neoplasia [11,24,32,33]. MRI has an important role as a problem-solving examination for lesion characterization and surgical planning, but there is, to our knowledge, no direct evidence to support the use of MRI as the initial or primary imaging examination for evaluation of pelvic pain localized to the perineum, vulva, or vagina, particularly when the physical examination is normal. However, there is emerging evidence to support the first-line utility of MRI when endometriosis or fistulizing disease are suspected [34]; readers are again referred to specific ACR Appropriateness Criteria guidelines for these clinical scenarios (see [Appendix 1](#)).

MRI also enables accurate depiction of pelvic floor muscular anatomy, integrity, and function [35,36]. Pelvic floor dysfunction is discussed in detail in a separate ACR Appropriateness Criteria document (see [Appendix 1](#)), but specific note is made here of the usefulness of MRI for assessment of muscular hypertonicity in chronic pelvic pain syndromes [37].

## Summary of Recommendations

- **Variante 1:** US pelvis transvaginal, US duplex Doppler pelvis, and US pelvis transabdominal are usually appropriate for the initial imaging of postmenopausal subacute or chronic pelvic pain localized to the deep pelvis. These procedures are complementary and should be performed together.
- **Variante 2:** US duplex Doppler pelvis, US pelvis transvaginal, and US pelvis transabdominal are usually appropriate for the initial imaging of postmenopausal subacute or chronic pelvic pain with clinically suspected pathologies in the perineum, vulva, or vagina. These procedures are complementary and should be performed together.

## Summary of Evidence

Of the 38 references cited in the *ACR Appropriateness Criteria® Postmenopausal Subacute or Chronic Pelvic Pain* document, all of them are categorized as diagnostic references including 1 well-designed study, 4 good-quality studies, and 10 quality studies that may have design limitations. There are 23 references that may not be useful as primary evidence.

The 38 references cited in the *ACR Appropriateness Criteria® Postmenopausal Subacute or Chronic Pelvic Pain* document were published from 2005 to 2017.

Although there are references that report on studies with design limitations, 5 well-designed or good-quality studies provide good evidence.

## Appropriateness Category Names and Definitions

Appropriateness Category Name	Appropriateness Rating	Appropriateness Category Definition
Usually Appropriate	7, 8, or 9	The imaging procedure or treatment is indicated in the specified clinical scenarios at a favorable risk-benefit ratio for patients.
May Be Appropriate	4, 5, or 6	The imaging procedure or treatment may be indicated in the specified clinical scenarios as an alternative to imaging procedures or treatments with a more favorable risk-benefit ratio, or the risk-benefit ratio for patients is equivocal.
May Be Appropriate (Disagreement)	5	The individual ratings are too dispersed from the panel median. The different label provides transparency regarding the panel's recommendation. "May be appropriate" is the rating category and a rating of 5 is assigned.
Usually Not Appropriate	1, 2, or 3	The imaging procedure or treatment is unlikely to be indicated in the specified clinical scenarios, or the risk-benefit ratio for patients is likely to be unfavorable.

### Relative Radiation Level Information

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Patients in the pediatric age group are at inherently higher risk from exposure, both because of organ sensitivity and longer life expectancy (relevant to the long latency that appears to accompany radiation exposure). For these reasons, the RRL dose estimate ranges for pediatric examinations are lower as compared to those specified for adults (see Table below). Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® [Radiation Dose Assessment Introduction](#) document [38].

Relative Radiation Level Designations		
Relative Radiation Level*	Adult Effective Dose Estimate Range	Pediatric Effective Dose Estimate Range
○	0 mSv	0 mSv
⊗	<0.1 mSv	<0.03 mSv
⊗ ⊗	0.1-1 mSv	0.03-0.3 mSv
⊗ ⊗ ⊗	1-10 mSv	0.3-3 mSv
⊗ ⊗ ⊗ ⊗	10-30 mSv	3-10 mSv
⊗ ⊗ ⊗ ⊗ ⊗	30-100 mSv	10-30 mSv

\*RRL assignments for some of the examinations cannot be made, because the actual patient doses in these procedures vary as a function of a number of factors (eg, region of the body exposed to ionizing radiation, the imaging guidance that is used). The RRLs for these examinations are designated as "Varies".

### Supporting Documents

For additional information on the Appropriateness Criteria methodology and other supporting documents go to [www.acr.org/ac](http://www.acr.org/ac).

## References

1. Ahangari A. Prevalence of chronic pelvic pain among women: an updated review. *Pain Physician* 2014;17:E141-7.
2. Latthe P, Latthe M, Say L, Gulmezoglu M, Khan KS. WHO systematic review of prevalence of chronic pelvic pain: a neglected reproductive health morbidity. *BMC Public Health* 2006;6:177.
3. Gemmell LC, Webster KE, Kirtley S, Vincent K, Zondervan KT, Becker CM. The management of menopause in women with a history of endometriosis: a systematic review. *Hum Reprod Update* 2017:1-20.
4. Bagarinao E, Johnson KA, Martucci KT, et al. Preliminary structural MRI based brain classification of chronic pelvic pain: A MAPP network study. *Pain* 2014;155:2502-9.
5. Berman SM, Naliboff BD, Suyenobu B, et al. Reduced brainstem inhibition during anticipated pelvic visceral pain correlates with enhanced brain response to the visceral stimulus in women with irritable bowel syndrome. *J Neurosci* 2008;28:349-59.
6. Borg C, Georgiadis JR, Renken RJ, Spoelstra SK, Weijmar Schultz W, de Jong PJ. Brain processing of visual stimuli representing sexual penetration versus core and animal-reminder disgust in women with lifelong vaginismus. *PLoS One* 2014;9:e84882.
7. Gupta A, Rapkin AJ, Gill Z, et al. Disease-related differences in resting-state networks: a comparison between localized provoked vulvodynia, irritable bowel syndrome, and healthy control subjects. *Pain* 2015;156:809-19.
8. Shobeiri SA, Rostaminia G, White D, Quiroz LH, Nihira MA. Evaluation of vaginal cysts and masses by 3-dimensional endovaginal and endoanal sonography. *J Ultrasound Med* 2013;32:1499-507.
9. Cicchiello LA, Hamper UM, Scutt LM. Ultrasound evaluation of gynecologic causes of pelvic pain. *Obstet Gynecol Clin North Am* 2011;38:85-114, viii.
10. Ganeshan A, Upponi S, Hon LQ, Uthappa MC, Warakaulle DR, Uberoi R. Chronic pelvic pain due to pelvic congestion syndrome: the role of diagnostic and interventional radiology. *Cardiovasc Intervent Radiol* 2007;30:1105-11.
11. Kuligowska E, Deeds L, 3rd, Lu K, 3rd. Pelvic pain: overlooked and underdiagnosed gynecologic conditions. *Radiographics* 2005;25:3-20.
12. Somprasit C, Tanprasertkul C, Suwannarurk K, Pongroj paw D, Chanthasenanont A, Bhamarapratana K. Transvaginal color Doppler study of uterine artery: is there a role in chronic pelvic pain? *J Obstet Gynaecol Res* 2010;36:1174-8.
13. Patel MD, Dubinsky TJ. Reimaging the female pelvis with ultrasound after CT: general principles. *Ultrasound Q* 2007;23:177-87.
14. Yitta S, Mausner EV, Kim A, et al. Pelvic ultrasound immediately following MDCT in female patients with abdominal/pelvic pain: is it always necessary? *Emerg Radiol* 2011;18:371-80.
15. Juhan V. Chronic pelvic pain: An imaging approach. *Diagn Interv Imaging* 2015;96:997-1007.
16. Tabibian N, Swehli E, Boyd A, Umbreen A, Tabibian JH. Abdominal adhesions: A practical review of an often overlooked entity. *Ann Med Surg (Lond)* 2017;15:9-13.
17. Silva PD, Suarez SA. A Case of Disabling Urinary Frequency and Pelvic Pain Due to Postoperative Uterine Adhesions. *WMJ* 2016;115:43-5.
18. Ignacio EA, Dua R, Sarin S, et al. Pelvic congestion syndrome: diagnosis and treatment. *Semin Intervent Radiol* 2008;25:361-8.
19. Karaosmanoglu D, Karcaaltincaba M, Karcaaltincaba D, Akata D, Ozmen M. MDCT of the ovarian vein: normal anatomy and pathology. *AJR Am J Roentgenol* 2009;192:295-9.
20. Koc Z, Ulsan S, Oguzkurt L. Right ovarian vein drainage variant: is there a relationship with pelvic varices? *Eur J Radiol* 2006;59:465-71.
21. Koc Z, Ulsan S, Oguzkurt L. Association of left renal vein variations and pelvic varices in abdominal MDCT. *Eur Radiol* 2007;17:1267-74.
22. Wang R, Yan Y, Zhan S, et al. Diagnosis of ovarian vein syndrome (OVS) by computed tomography (CT) imaging: a retrospective study of 11 cases. *Medicine (Baltimore)* 2014;93:e53.
23. Thomassin-Naggara I, Darai E, Bazot M. Gynecological pelvic infection: what is the role of imaging? *Diagn Interv Imaging* 2012;93:491-9.
24. Valentini AL, Gui B, Basilico R, Di Molfetta IV, Micco M, Bonomo L. Magnetic resonance imaging in women with pelvic pain from gynaecological causes: a pictorial review. *Radiol Med* 2012;117:575-92.

25. American College of Radiology. *Manual on Contrast Media*. Available at: <https://www.acr.org/Clinical-Resources/Contrast-Manual>. Accessed September 30, 2018.
26. Ascitutto G, Mumme A, Marpe B, Koster O, Ascitutto KC, Geier B. MR venography in the detection of pelvic venous congestion. *Eur J Vasc Endovasc Surg* 2008;36:491-6.
27. Yang DM, Kim HC, Nam DH, Jahng GH, Huh CY, Lim JW. Time-resolved MR angiography for detecting and grading ovarian venous reflux: comparison with conventional venography. *Br J Radiol* 2012;85:e117-22.
28. Leiber LM, Thouveny F, Bouvier A, et al. MRI and venographic aspects of pelvic venous insufficiency. *Diagn Interv Imaging* 2014;95:1091-102.
29. Bharwani N, Tirlapur SA, Balogun M, et al. MRI reporting standard for chronic pelvic pain: consensus development. *Br J Radiol* 2016;89:20140615.
30. Dick EA, Burnett C, Anstee A, Hamady M, Black D, Gedroyc WM. Time-resolved imaging of contrast kinetics three-dimensional (3D) magnetic resonance venography in patients with pelvic congestion syndrome. *Br J Radiol* 2010;83:882-7.
31. van der Meijden WI, Boffa MJ, Ter Harmsel WA, et al. 2016 European guideline for the management of vulval conditions. *J Eur Acad Dermatol Venereol* 2017;31:925-41.
32. Hwang JH, Oh MJ, Lee NW, Hur JY, Lee KW, Lee JK. Multiple vaginal mullerian cysts: a case report and review of literature. *Arch Gynecol Obstet* 2009;280:137-9.
33. Surabhi VR, Menias CO, George V, Siegel CL, Prasad SR. Magnetic resonance imaging of female urethral and periurethral disorders. *Radiol Clin North Am* 2013;51:941-53.
34. Tirlapur SA, Daniels JP, Khan KS. Chronic pelvic pain: how does noninvasive imaging compare with diagnostic laparoscopy? *Curr Opin Obstet Gynecol* 2015;27:445-8.
35. Quinn M. Injuries to the levator ani in unexplained, chronic pelvic pain. *J Obstet Gynaecol* 2007;27:828-31.
36. Savoye-Collet C, Koning E, Dacher JN. Radiologic evaluation of pelvic floor disorders. *Gastroenterol Clin North Am* 2008;37:553-67, viii.
37. Ackerman AL, Lee UJ, Jellison FC, et al. MRI suggests increased tonicity of the levator ani in women with interstitial cystitis/bladder pain syndrome. *Int Urogynecol J* 2016;27:77-83.
38. American College of Radiology. ACR Appropriateness Criteria® Radiation Dose Assessment Introduction. Available at: <https://www.acr.org/-/media/ACR/Files/Appropriateness-Criteria/RadiationDoseAssessmentIntro.pdf>. Accessed September 30, 2018.

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.



## Appendix 1. Related ACR Appropriateness Criteria Topics

Subject area	AC topic
Vaginal bleeding in postmenopausal women	<a href="#">Abnormal Vaginal Bleeding</a>
Clinically suspected adnexal mass	<a href="#">Clinically Suspected Adnexal Mass</a>
Acute pelvic pain in postmenopausal women	Postmenopausal Acute Pelvic Pain – Topic under development
Pelvic floor dysfunction	<a href="#">Pelvic Floor Dysfunction</a>
Endometriosis	<a href="#">Infertility</a>
Pelvic inflammatory disease	<a href="#">Acute Pelvic Pain in the Reproductive Age Patient</a>
Leiomyomas	<a href="#">Abnormal Vaginal Bleeding</a> and <a href="#">Clinically Suspected Adnexal Mass</a>
Urinary tract infection	<a href="#">Recurrent Lower Urinary Tract Infections in Women</a>
Endometrial cancer	<a href="#">Pretreatment Evaluation and Follow-up of Endometrial Cancer</a>
Vaginal cancer	Staging and Follow-up of Vaginal Cancer – Topic under development
Vulvar cancer	Staging and Follow-up of Vulvar Cancer – Topic under development
Hematuria	<a href="#">Hematuria</a>
Diverticulitis	<a href="#">Left Lower Quadrant Pain-Suspected Diverticulitis</a>
Left lower quadrant pain	<a href="#">Left Lower Quadrant Pain-Suspected Diverticulitis</a>
Right lower quadrant pain	<a href="#">Right Lower Quadrant Pain-Suspected Appendicitis</a>
Suspected small bowel obstruction	<a href="#">Suspected Small-Bowel Obstruction</a>