

**American College of Radiology  
ACR Appropriateness Criteria®  
Workup of Noncerebral Systemic Arterial Embolic Source**

**Variant 1:**                    **Known upper extremity arterial occlusion. Suspected embolic etiology. Next imaging study to determine source.**

Procedure	Appropriateness Category	Relative Radiation Level
US echocardiography transesophageal	Usually Appropriate	○
US echocardiography transthoracic resting	Usually Appropriate	○
MRA chest without and with IV contrast	Usually Appropriate	○
MRI heart function and morphology without and with IV contrast	Usually Appropriate	○
MRI heart function and morphology without IV contrast	Usually Appropriate	○
CTA chest with IV contrast	Usually Appropriate	⊕⊕⊕
CT heart function and morphology with IV contrast	Usually Appropriate	⊕⊕⊕⊕
MRA chest without IV contrast	May Be Appropriate	○
US duplex Doppler abdomen	Usually Not Appropriate	○
Aortography chest	Usually Not Appropriate	⊕⊕⊕

**Variant 2:**                    **Known arterial occlusion in the mesenteric or renal arterial system or renal infarcts. Suspected embolic etiology. Next imaging study to determine source.**

Procedure	Appropriateness Category	Relative Radiation Level
US echocardiography transesophageal	Usually Appropriate	○
US echocardiography transthoracic resting	Usually Appropriate	○
MRA chest and abdomen without and with IV contrast	Usually Appropriate	○
MRI heart function and morphology without and with IV contrast	Usually Appropriate	○
MRI heart function and morphology without IV contrast	Usually Appropriate	○
CTA chest with IV contrast	Usually Appropriate	⊕⊕⊕
CT heart function and morphology with IV contrast	Usually Appropriate	⊕⊕⊕⊕
CTA chest and abdomen with IV contrast	Usually Appropriate	⊕⊕⊕⊕
US duplex Doppler abdomen	May Be Appropriate	○
MRA chest and abdomen without IV contrast	May Be Appropriate	○
MRA chest without and with IV contrast	May Be Appropriate	○
MRA chest without IV contrast	May Be Appropriate	○
Aortography chest and abdomen	Usually Not Appropriate	⊕⊕⊕⊕

**Variant 3:****Known lower extremity arterial occlusion. Suspected embolic etiology. Next imaging study to determine source.**

Procedure	Appropriateness Category	Relative Radiation Level
US echocardiography transesophageal	Usually Appropriate	○
US echocardiography transthoracic resting	Usually Appropriate	○
MRA chest abdomen pelvis without and with IV contrast	Usually Appropriate	○
MRA chest without and with IV contrast	Usually Appropriate	○
MRI heart function and morphology without and with IV contrast	Usually Appropriate	○
MRI heart function and morphology without IV contrast	Usually Appropriate	○
CTA chest with IV contrast	Usually Appropriate	⊕⊕⊕
CT heart function and morphology with IV contrast	Usually Appropriate	⊕⊕⊕⊕
CTA chest abdomen pelvis with IV contrast	Usually Appropriate	⊕⊕⊕⊕⊕
MRA chest abdomen pelvis without IV contrast	May Be Appropriate	○
MRA chest without IV contrast	May Be Appropriate	○
US duplex Doppler abdomen	Usually Not Appropriate	○
Aortography chest abdomen pelvis	Usually Not Appropriate	⊕⊕⊕⊕

**Variant 4:****Known multiorgan system arterial occlusions. Suspected embolic etiology. Next imaging study to determine source.**

Procedure	Appropriateness Category	Relative Radiation Level
US echocardiography transesophageal	Usually Appropriate	○
US echocardiography transthoracic resting	Usually Appropriate	○
MRA chest abdomen pelvis without and with IV contrast	Usually Appropriate	○
MRA chest abdomen pelvis without IV contrast	Usually Appropriate	○
MRA chest without and with IV contrast	Usually Appropriate	○
MRI heart function and morphology without and with IV contrast	Usually Appropriate	○
MRI heart function and morphology without IV contrast	Usually Appropriate	○
CTA chest with IV contrast	Usually Appropriate	⊕⊕⊕
CT heart function and morphology with IV contrast	Usually Appropriate	⊕⊕⊕⊕
CTA chest abdomen pelvis with IV contrast	Usually Appropriate	⊕⊕⊕⊕⊕
US duplex Doppler abdomen	May Be Appropriate	○
MRA chest without IV contrast	May Be Appropriate	○

## WORKUP OF NONCEREBRAL SYSTEMIC ARTERIAL EMBOLIC SOURCE

Expert Panels on Vascular Imaging and Cardiac Imaging: Vincent G. Parenti, MD<sup>a</sup>; Kanupriya Vijay, MD, MBBS<sup>b</sup>; Christopher D. Maroules, MD<sup>c</sup>; Bill S. Majdalany, MD<sup>d</sup>; Lynne M. Koweek, MD<sup>e</sup>; Minhajuddin S. Khaja, MD, MBA<sup>f</sup>; Brian B. Ghoshhajra, MD, MBA<sup>g</sup>; Prachi P. Agarwal, MD<sup>h</sup>; Benjamin N. Contrella, MD<sup>i</sup>; Nicole A. Keefe, MD<sup>j</sup>; Bruce M. Lo, MD, RDMS, MBA<sup>k</sup>; Sachin B. Malik, MD<sup>l</sup>; Devaki Shilpa Surasi, MD<sup>m</sup>; Kathleen Waite, MD<sup>n</sup>; Eric E. Williamson, MD<sup>o</sup>; Suhny Abbara, MD<sup>p</sup>; Karin E. Dill, MD.<sup>q</sup>

### Summary of Literature Review

#### **Introduction/Background**

Noncerebral systemic arterial embolism is an important cause of patient morbidity and mortality [1]. Arterial emboli can originate from a variety of cardiac and noncardiac sources. Cardiac sources include thrombus within the left atrium and left ventricle, valvular disease such as endocarditis, and cardiac neoplasms. Noncardiac sources of arterial embolism include thrombus and atherosclerosis within the aorta and peripheral arteries.

Intracardiac thrombus has been thoroughly described in the cardiology and neurology literature with several factors that predispose its formation and the potential for arterial embolization. For example, atrial fibrillation has been shown to be a significant risk factor for atrial thrombogenesis [2,3]. Complex left atrial appendage morphology also confers increased likelihood of thrombus development [4]. Myocardial infarction often results in focal hypokinesis or akinesis of the left ventricular myocardium, which predisposes to thrombus formation [5,6]. Aortic and mitral valve endocarditis as well as valvular neoplasms are other potential sources for arterial embolism detectable with imaging [7,8]. Aortic thrombi tend to be associated with aortic pathology including dissection, aneurysm, or ulcerative lesions [9,10]. Thrombus formation can also occur in the aorta secondary to hypercoagulable states such as malignancy, trauma, postoperative states, hormonal therapy, and inherited hypercoagulable disorders [1,11,12].

When a cardiac or noncardiac embolic source dislodges, the resulting embolus can occlude a variety of peripheral and visceral arteries causing ischemia [1,9,11,12]. Characteristic locations for noncerebral arterial occlusion include the upper extremities, abdominal viscera, and lower extremities [1,9,11]. Ischemia in these regions can progress to tissue infarction resulting in limb amputation, bowel resection, or nephrectomy. Determining the source of arterial embolism is essential in order to direct treatment decisions. Treatment options vary and include anti-coagulation, endovascular or surgical embolectomy, and peripheral arterial angioplasty with or without stenting to maintain long-term vascular patency [1,9,11-14].

The variants in this document assume that the diagnosis of an arterial occlusion has already been established by other means. For example, in the setting of an acute onset cold painful leg, the use of lower extremity arteriography, CT angiography (CTA), or MR angiography (MRA) could be employed to demonstrate arterial occlusion. This document specifically pertains to the workup of a suspected embolic etiology of the already known arterial occlusion.

#### **Special Imaging Considerations**

For the purposes of distinguishing between CT and CT angiography (CTA), ACR Appropriateness Criteria topics use the definition in the [ACR–NASCI–SIR–SPR Practice Parameter for the Performance and Interpretation of Body Computed Tomography Angiography \(CTA\)](#) [15]:

---

<sup>a</sup>Research Author, University of Texas Southwestern Medical Center, Dallas, Texas. <sup>b</sup>UT Southwestern Medical Center, Dallas, Texas. <sup>c</sup>Naval Medical Center Portsmouth, Portsmouth, Virginia. <sup>d</sup>Panel Chair, Emory Healthcare, Atlanta, Georgia. <sup>e</sup>Panel Chair, Duke University Medical Center, Durham, North Carolina. <sup>f</sup>Panel Vice-Chair, University of Michigan, Ann Arbor, Michigan. <sup>g</sup>Panel Vice-Chair, Massachusetts General Hospital, Boston, Massachusetts. <sup>h</sup>University of Michigan, Ann Arbor, Michigan. <sup>i</sup>Allegheny Health Network, Pittsburgh, Pennsylvania. <sup>j</sup>University of North Carolina School of Medicine, Chapel Hill, North Carolina. <sup>k</sup>Sentara Norfolk General Hospital/Eastern Virginia Medical School, Norfolk, Virginia; American College of Emergency Physicians. <sup>l</sup>VA Palo Alto Health Care System, Palo Alto, California and Stanford University, Stanford, California. <sup>m</sup>The University of Texas MD Anderson Cancer Center, Houston, Texas; Commission on Nuclear Medicine and Molecular Imaging. <sup>n</sup>Duke University Medical Center, Durham, North Carolina, Primary care physician. <sup>o</sup>Mayo Clinic, Rochester, Minnesota; Society of Cardiovascular Computed Tomography. <sup>p</sup>Specialty Chair, UT Southwestern Medical Center, Dallas, Texas. <sup>q</sup>Specialty Chair, Emory University Hospital, Atlanta, Georgia.

The American College of Radiology seeks and encourages collaboration with other organizations on the development of the ACR Appropriateness Criteria through representation of such organizations on expert panels. Participation on the expert panel does not necessarily imply endorsement of the final document by individual contributors or their respective organization.

Reprint requests to: [publications@acr.org](mailto:publications@acr.org)

*“CTA uses a thin-section CT acquisition that is timed to coincide with peak arterial or venous enhancement. The resultant volumetric dataset is interpreted using primary transverse reconstructions as well as multiplanar reformations and 3-D renderings.”*

All elements are essential: 1) timing, 2) reconstructions/reformats, and 3) 3-D renderings. Standard CTs with contrast also include timing issues and reconstructions/reformats. Only in CTA, however, is 3-D rendering a **required** element. This corresponds to the definitions that the CMS has applied to the Current Procedural Terminology codes.

### **Initial Imaging Definition**

Initial imaging is defined as imaging at the beginning of the care episode for the medical condition defined by the variant. More than one procedure can be considered usually appropriate in the initial imaging evaluation when:

- There are procedures that are equivalent alternatives (ie, only one procedure will be ordered to provide the clinical information to effectively manage the patient’s care)

OR

- There are complementary procedures (ie, more than one procedure is ordered as a set or simultaneously where each procedure provides unique clinical information to effectively manage the patient’s care).

### **Discussion of Procedures by Variant**

#### **Variant 1: Known upper extremity arterial occlusion. Suspected embolic etiology. Next imaging study to determine source.**

The variant assumes that an upper extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

#### **Aortography Chest**

Conventional catheter aortography has largely been replaced by noninvasive imaging modalities such as CTA and MRA given their high sensitivity/specificity for detecting aortic pathologies such as mural thrombus [16,17]. Aortography is typically used as an alternative diagnostic strategy following initial noninvasive imaging and when therapeutic interventions are being considered [11,17].

#### **CT Heart Function and Morphology With IV Contrast**

The primary role of cardiac CT in the initial evaluation of upper extremity arterial embolic occlusion is in the workup of cardiac thrombus as a source. Multiple studies have established high rates of atrial thrombus detection by cardiac CT compared to transesophageal echocardiography (TEE) [18-27]. Meta-analyses have found sensitivities of 96% to 99% and specificities of 92% to 94% for detection of left atrial or left atrial appendage thrombus with cardiac CT compared to a TEE reference standard [28-30]. When compared with intraoperative findings, cardiac CT was 100% sensitive and 85% specific for finding left atrial thrombus [31]. Complex left atrial appendage morphologies which predispose to thrombus formation can also be characterized by cardiac CT [32-34]. Additionally, cardiac CT can differentiate left ventricular thrombus from the myocardial wall with 1 study demonstrating a sensitivity, specificity, and positive and negative predictive values of 94%, 97%, 94%, and 97%, respectively [35]. Studies have also demonstrated cardiac CT to have comparable accuracy to TEE for identification of vegetations in the setting of infective endocarditis, another potential source of arterial embolism [36-38]. Cardiac CT can identify cardiac neoplasms, both benign and malignant, which have the potential to shed and embolize to distal arterial beds [39,40].

#### **CTA Chest With IV Contrast**

Multidetector chest CTA with intravenous (IV) contrast can be used to evaluate for at-risk atherosclerotic plaque or the presence of thrombus in the thoracic aorta. CTA is useful in the assessment of the size, extent, and location of an embolic source in the aorta, which can aid in management decisions [13,41]. A number of small studies have used chest CTA to detect aortic mural thrombus that was suspected of embolization [1,12-14,42]. Specific data on the sensitivity and specificity of this imaging modality are lacking.

### **MRA Chest Without and With IV Contrast**

Chest MRA without and with IV contrast can be used to evaluate for the presence of thrombus in the thoracic aorta. In 1 study, detection of thoracic aorta pathology by contrast-enhanced chest MRA was equivalent in sensitivity, specificity, and diagnostic accuracy compared to noncontrast MRA, although only a single case of thrombus was included in the analysis [43]. On the other hand, in a small analysis which included 9 patients with aortic thrombus, contrast-enhanced MRA had a lower thrombus detection rate compared to a noncontrast examination, although this finding was not statistically significant [44]. Data comparing MRA of the chest to other imaging modalities are lacking.

### **MRA Chest Without IV Contrast**

Chest MRA without IV contrast is an imaging study, which can detect the presence of thrombus in the thoracic aorta. One small study found this examination to have a higher detection rate for aortic thrombus when compared with contrast-enhanced MRA, although the difference was not statistically significant [44]. In another study, sensitivity, specificity, and diagnostic accuracy of unenhanced steady-state free precession MRA were 100% for the detection of thoracic aorta pathology compared to a contrast-enhanced MRA reference standard; however, this analysis only included 1 case of mural thrombus [43]. Data comparing MRA of the chest to other imaging modalities are lacking.

### **MRI Heart Function and Morphology Without and With IV Contrast**

Cardiac MR is a noninvasive imaging study that can detect intracardiac thrombus as well as valvular and neoplastic pathologies. A meta-analysis of 7 studies showed that delayed contrast-enhanced cardiac MR had a pooled sensitivity of 100% and a specificity of 99% for detecting left atrial and left atrial appendage thrombus in patients with atrial fibrillation [45]. In another meta-analysis, there was no significant difference in sensitivity and specificity between cardiac CT and cardiac MR in the detection of left atrial appendage thrombus [29]. Contrast-enhanced cardiac MR had a sensitivity of 88% and a specificity of 99% compared to surgical or pathological confirmation of left ventricular thrombus [46]. Cardiac MR is also an accurate imaging modality for the evaluation of valvular disease, including aortic and mitral valve vegetations, which can dislodge and result in arterial embolism [37,47]. Additionally, cardiac MR offers detailed soft tissue characterization for the analysis of benign and malignant intracardiac neoplasms [39,48].

### **MRI Heart Function and Morphology Without IV Contrast**

Cardiac MR without contrast provides a detailed anatomic evaluation of the heart chambers. In the workup of embolic sources, the primary role of cardiac MR is in the identification of intracardiac thrombus. A meta-analysis of 7 studies showed that cine cardiac MR had a pooled sensitivity of 91% and a specificity of 93% for detecting left atrial and left atrial appendage thrombus in patients with atrial fibrillation [45]. Furthermore, cine cardiac MR had an 82% sensitivity and a 100% specificity in detecting left ventricle thrombus in postmyocardial infarction patients compared with a standard delayed enhancement cardiac MR [49]. Cardiac MR without contrast is also capable of identifying valvular pathology and cardiac neoplasms, although data on its applicability in the setting of systemic arterial embolism are lacking.

### **US Duplex Doppler Abdomen**

There is no relevant literature to support the use of Doppler ultrasound (US) of the abdomen as an initial imaging modality in the evaluation of the source of known embolic upper extremity arterial occlusion.

### **US Echocardiography Transesophageal**

TEE is an invasive diagnostic study with the ability to detect cardiac pathology predisposed to embolism. TEE has a sensitivity of 93% to 100% and a specificity of 95% to 99% for detecting left atrial appendage thrombus when compared to intraoperative findings [31,50,51]. Furthermore, TEE can evaluate left ventricular systolic dysfunction, spontaneous echo contrast, slow left atrial appendage peak flow velocities, and complex left atrial appendage morphologies, which are all associated with left atrial thrombus and thromboembolic risk [2,4]. In addition, TEE can detect left ventricular thrombus with 1 study reporting a 40% sensitivity and a 96% specificity for the modality compared to findings at surgery or pathology [46]. Proximal aortic thrombus can also be assessed using TEE, although evaluation is limited by blind spots (distal ascending aorta and proximal aortic arch) owing to air in the trachea [10,13]. Detection of valvular disease and intracardiac neoplasms can also be accomplished with TEE.

### **US Echocardiography Transthoracic Resting**

Transthoracic echocardiography (TTE) is a noninvasive imaging modality capable of detecting cardiac pathology susceptible to embolism. TTE is inferior to TEE in the assessment of left atrial appendage thrombus because the

transducer is distant from the left atrium when placed on the chest [52]. In 1 study, a cardiac embolic source was detected by TEE in about 40% of patients with normal TTE [53]. In another study, a cardiac embolic source was identified by TTE in 15% of the study group compared with 57% by TEE [54]. Sensitivity and specificity were 23% and 96%, respectively, for the detection of left ventricular thrombus compared to findings at surgery or pathology [46]. In the detection of left ventricle thrombus, contrast-enhanced TTE had a 64% sensitivity and a 99% specificity compared to a delayed enhancement cardiac MR standard [49]. TTE can also be applied for the diagnosis of valvular disease and cardiac neoplasms. There is no evidence to support the use of TTE in the evaluation of aortic thrombus.

**Variant 2: Known arterial occlusion in the mesenteric or renal arterial system or renal infarcts. Suspected embolic etiology. Next imaging study to determine source.**

The variant assumes that a mesenteric/renal arterial occlusion or renal infarct has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on “[Imaging of Mesenteric Ischemia](#)” [55]).

**Aortography Chest and Abdomen**

Conventional catheter aortography has largely been replaced by noninvasive imaging modalities such as CTA and MRA given their high sensitivity/specificity for detecting aortic pathologies such as mural thrombus [16,17]. Aortography is typically used as an alternative diagnostic strategy following initial noninvasive imaging and when therapeutic interventions are being considered [11,17].

**CT Heart Function and Morphology With IV Contrast**

The primary role of cardiac CT in the initial evaluation of mesenteric or renal arterial embolic occlusion is in the workup of cardiac thrombus as a source. Multiple studies have established high rates of atrial thrombus detection by cardiac CT compared to TEE [18-27]. Meta-analyses have found sensitivities of 96% to 99% and specificities of 92% to 94% for detection of left atrial or left atrial appendage thrombus with cardiac CT compared to a TEE reference standard [28-30]. When compared to intraoperative findings, cardiac CT was 100% sensitive and 85% specific for finding left atrial thrombus [31]. Complex left atrial appendage morphologies that predispose to thrombus formation can also be characterized by cardiac CT [32-34]. Additionally, cardiac CT can differentiate left ventricular thrombus from the myocardial wall, with 1 study demonstrating a sensitivity, specificity, and positive and negative predictive values of 94%, 97%, 94%, and 97%, respectively [35]. Studies have also demonstrated cardiac CT to have comparable accuracy to TEE for identification of vegetations in the setting of infective endocarditis, another potential source of arterial embolism [36-38]. Cardiac CT can identify cardiac neoplasms, both benign and malignant, which have the potential to shed and embolize to distal arterial beds [39,40].

**CTA Chest With IV Contrast**

In some conditions or clinical scenarios, there may be a high suspicion that the embolic source is in the thoracic aorta, and a CTA limited to the chest may be diagnostic. As such, multidetector chest CTA with IV contrast can be used to evaluate for at-risk atherosclerotic plaque or the presence of thrombus in the thoracic aorta. CTA is useful in the assessment of the size, extent, and location of an embolic source in the aorta, which can aid in management decisions [13,41]. A number of small studies have used chest CTA to detect aortic mural thrombus that was suspected of embolization [1,12-14,42]. Specific data on the sensitivity and specificity of this imaging modality are lacking.

**CTA Chest and Abdomen With IV Contrast**

Multidetector CTA with IV contrast can be used to evaluate for the presence of at-risk atherosclerotic plaque or thrombus in the aorta in its entirety. CTA is useful in the assessment of the size, extent, and location of an embolic source in the aorta, which can aid in management decisions [13,41]. Aortic intraluminal thrombus is oftentimes associated with aneurysm, particularly in the abdomen, which is readily detected by CTA [56,57]. A number of small studies have used CTA to detect aortic mural thrombus that was suspected of embolization [1,12-14,42]. Specific data on the sensitivity and specificity of this imaging modality are lacking.

**MRA Chest Without and With IV Contrast**

In some conditions or clinical scenarios, there may be a high suspicion that the embolic source is in the thoracic aorta and an MRA limited to the chest may be diagnostic. In 1 study, detection of thoracic aorta pathology by contrast-enhanced chest MRA was equivalent in sensitivity, specificity, and diagnostic accuracy compared to noncontrast MRA, although only a single case of thrombus was included in the analysis [43]. On the other hand, in

a small analysis which included 9 patients with aortic thrombus, contrast-enhanced MRA had a lower thrombus detection rate compared to a noncontrast examination, although this finding was not statistically significant [44]. Data comparing MRA of the chest to other imaging modalities are lacking.

#### **MRA Chest Without IV Contrast**

In some conditions or clinical scenarios, there may be a high suspicion that the embolic source is in the thoracic aorta and an MRA limited to the chest may be diagnostic. One small study found this examination to have a higher detection rate for aortic thrombus when compared with contrast-enhanced MRA, although the difference was not statistically significant [44]. In another study, sensitivity, specificity, and diagnostic accuracy of unenhanced steady-state free precession MRA were 100% for the detection of thoracic aorta pathology compared to a contrast-enhanced MRA reference standard; however, this analysis only included 1 case of mural thrombus [43]. Data comparing MRA of the chest to other imaging modalities is lacking.

#### **MRA Chest and Abdomen Without and With IV Contrast**

MRA of the chest and abdomen without and with IV contrast can be used to evaluate for presence of an embolic source in the aorta in its entirety. In 1 study, detection of thoracic aorta pathology by contrast-enhanced chest MRA was equivalent in sensitivity, specificity, and diagnostic accuracy compared to noncontrast MRA, although only a single case of thrombus was included in the analysis [43]. On the other hand, in a small analysis which included 9 patients with aortic thrombus, contrast-enhanced MRA had a lower thrombus detection rate compared to a noncontrast examination, although this finding was not statistically significant [44]. Contrast-enhanced MRA of the abdomen has been used for intraluminal thrombus detection in the setting of aneurysms, although comparative data are insufficient [56-58]. Data comparing MRA of the chest and abdomen to other imaging modalities are lacking.

#### **MRA Chest and Abdomen Without IV Contrast**

Chest and abdomen MRA without IV contrast can be used to evaluate for the presence of an embolic source in the aorta in its entirety. One small study found this examination to have a higher detection rate for aortic thrombus when compared with contrast-enhanced MRA, although the difference was not statistically significant [44]. In another study, sensitivity, specificity, and diagnostic accuracy of unenhanced steady-state free precession MRA were 100% for the detection of thoracic aorta pathology compared to a contrast-enhanced MRA reference standard; however, this analysis only included 1 case of mural thrombus [43]. Noncontrast MRA has been used for the detection of abdominal aortic intraluminal thrombus, although there are insufficient data comparing it to contrast-enhanced MRA [56-58]. Data comparing MRA of the chest and abdomen to other imaging modalities are lacking.

#### **MRI Heart Function and Morphology Without and With IV Contrast**

Cardiac MR is a noninvasive imaging study that can reliably detect intracardiac thrombus as well as valvular and neoplastic pathologies. A meta-analysis of 7 studies showed that delayed contrast-enhanced cardiac MR had a pooled sensitivity of 100% and a specificity of 99% for detecting left atrial and left atrial appendage thrombus in patients with atrial fibrillation [45]. In another meta-analysis, there was no significant difference in sensitivity and specificity between cardiac CT and cardiac MR in the detection of left atrial appendage thrombus [29]. Contrast-enhanced cardiac MR had a sensitivity of 88% and a specificity of 99% compared to surgical or pathological confirmation of left ventricular thrombus [46]. Cardiac MR is also an accurate imaging modality for the evaluation of valvular disease, including aortic and mitral valve vegetations, which can dislodge and result in arterial embolism [37,47]. Additionally, cardiac MR offers detailed soft tissue characterization for the analysis of benign and malignant intracardiac neoplasms [39,48].

#### **MRI Heart Function and Morphology Without IV Contrast**

Cardiac MR without contrast provides a detailed anatomic evaluation of the heart chambers. In the workup of embolic sources, the primary role of cardiac MR is in the identification of intracardiac thrombus. A meta-analysis of 7 studies showed that cine cardiac MR had a pooled sensitivity of 91% and a specificity of 93% for detecting left atrial and left atrial appendage thrombus in patients with atrial fibrillation [45]. Furthermore, cine cardiac MR had an 82% sensitivity and 100% specificity in detecting left ventricle thrombus in postmyocardial infarction patients compared to a standard delayed enhancement cardiac MR [49]. Cardiac MR without contrast is also capable of identifying valvular pathology and cardiac neoplasms, although data on its applicability in the setting of systemic arterial embolism are lacking.

#### **US Duplex Doppler Abdomen**

There is no relevant literature to support the use of Doppler US of the abdomen as an initial imaging modality in the evaluation of the source of known embolic mesenteric/renal arterial occlusion. However, some imaging

protocols may include limited views of the abdominal aorta that may detect intraluminal aortic thrombus or significant atherosclerotic disease [56].

### **US Echocardiography Transesophageal**

TEE is an invasive diagnostic study with the ability to detect cardiac pathology predisposed to embolism. TEE has a sensitivity of 93% to 100% and a specificity of 95% to 99% for detecting left atrial appendage thrombus when compared to intraoperative findings [31,50,51]. Furthermore, TEE can evaluate left ventricular systolic dysfunction, spontaneous echo contrast, slow left atrial appendage peak flow velocities, and complex left atrial appendage morphologies, which are all associated with left atrial thrombus and thromboembolic risk [2,4]. In addition, TEE can detect left ventricular thrombus, with 1 study reporting a 40% sensitivity and a 96% specificity for the modality compared to findings at surgery or pathology [46]. Proximal aortic thrombus can also be assessed using TEE, although evaluation is limited by blind spots (distal ascending aorta and proximal aortic arch) owing to air in the trachea [10,13]. Detection of valvular disease and intracardiac neoplasms can also be accomplished with TEE.

### **US Echocardiography Transthoracic Resting**

TTE is a noninvasive imaging modality capable of detecting cardiac pathology susceptible to embolism. TTE is inferior to TEE in the assessment of left atrial appendage thrombus because the transducer is distant from the left atrium when placed on the chest [52]. In 1 study, a cardiac embolic source was detected by TEE in about 40% of patients with normal TTE [53]. In another study, a cardiac embolic source was identified by TTE in 15% of the study group compared with 57% by TEE [54]. Sensitivity and specificity were 23% and 96%, respectively, for the detection of left ventricular thrombus compared to findings at surgery or pathology [46]. In the detection of left ventricle thrombus, contrast-enhanced TTE had a 64% sensitivity and 99% specificity compared to a delayed enhancement cardiac MR standard [49]. TTE can also be applied for the diagnosis of valvular disease and cardiac neoplasms. There is no evidence to support the use of TTE in the evaluation of aortic thrombus.

### **Variant 3: Known lower extremity arterial occlusion. Suspected embolic etiology. Next imaging study to determine source.**

The variant assumes that a lower extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on “[Sudden Onset of Cold, Painful Leg](#)” [59]).

### **Aortography Chest, Abdomen, and Pelvis**

Conventional catheter aortography has largely been replaced by noninvasive imaging modalities such as CTA and MRA given their high sensitivity/specificity for detecting aortic pathologies such as mural thrombus [16,17]. Aortography is typically used as an alternative diagnostic strategy following initial noninvasive imaging and when therapeutic interventions are being considered [11,17].

### **CT Heart Function and Morphology With IV Contrast**

The primary role of cardiac CT in the initial evaluation of lower extremity arterial embolic occlusion is in the workup of cardiac thrombus as a source. Multiple studies have established high rates of atrial thrombus detection by cardiac CT compared to TEE [18-27]. Meta-analyses have found sensitivities of 96% to 99% and specificities of 92% to 94% for detection of left atrial or left atrial appendage thrombus with cardiac CT compared to a TEE reference standard [28-30]. When compared to intraoperative findings, cardiac CT was 100% sensitive and 85% specific for finding left atrial thrombus [31]. Complex left atrial appendage morphologies, which predispose to thrombus formation, can also be characterized by cardiac CT [32-34]. Additionally, cardiac CT can differentiate left ventricular thrombus from the myocardial wall with 1 study demonstrating a sensitivity, specificity, and positive and negative predictive values of 94%, 97%, 94%, and 97%, respectively [35]. Studies have also demonstrated cardiac CT to have comparable accuracy to TEE for identification of vegetations in the setting of infective endocarditis, another potential source of arterial embolism [36-38]. Cardiac CT can identify cardiac neoplasms, both benign and malignant, which have the potential to shed and embolize to distal arterial beds [39,40].

### **CTA Chest With IV Contrast**

In some conditions or clinical scenarios, there may be a high suspicion that the embolic source is in the thoracic aorta and a CTA limited to the chest may be diagnostic. As such, multidetector chest CTA with IV contrast can be used to evaluate for at-risk atherosclerotic plaque or the presence of thrombus in the thoracic aorta. CTA is useful in the assessment of the size, extent, and location of an embolic source in the aorta, which can aid in management decisions [13,41]. A number of small studies have used chest CTA to detect aortic mural thrombus that was



suspected of embolization [1,12-14,42]. Specific data on the sensitivity and specificity of this imaging modality are lacking.

#### **CTA Chest, Abdomen, and Pelvis With IV Contrast**

Multidetector CTA with IV contrast can be used to evaluate for the presence of at-risk atherosclerotic plaque or thrombus in the aorta in its entirety. CTA is useful in the assessment of the size, extent, and location of an embolic source in the aorta, which can aid in management decisions [13,41]. Aortic intraluminal thrombus is oftentimes associated with aneurysm, particularly in the abdomen, which is readily detected by CTA [56,57]. A number of small studies have used CTA to detect aortic mural thrombus that was suspected of embolization [1,12-14,42]. Specific data on the sensitivity and specificity of this imaging modality are lacking.

#### **MRA Chest Without and With IV Contrast**

In some conditions or clinical scenarios, there may be a high suspicion that the embolic source is in the thoracic aorta and an MRA limited to the chest may be diagnostic. In 1 study, detection of thoracic aorta pathology by contrast-enhanced chest MRA was equivalent in sensitivity, specificity, and diagnostic accuracy compared to noncontrast MRA, although only a single case of thrombus was included in the analysis [43]. On the other hand, in a small analysis which included 9 patients with aortic thrombus, contrast-enhanced MRA had a lower thrombus detection rate compared to a noncontrast examination, although this finding was not statistically significant [44]. Data comparing MRA of the chest to other imaging modalities are lacking.

#### **MRA Chest Without IV Contrast**

In some conditions or clinical scenarios, there may be a high suspicion that the embolic source is in the thoracic aorta and an MRA limited to the chest may be diagnostic. One small study found this examination to have a higher detection rate for aortic thrombus when compared with contrast-enhanced MRA, although the difference was not statistically significant [44]. In another study, sensitivity, specificity, and diagnostic accuracy of unenhanced steady-state free precession MRA were 100% for the detection of thoracic aorta pathology compared to a contrast-enhanced MRA reference standard; however, this analysis only included 1 case of mural thrombus [43]. Data comparing MRA of the chest to other imaging modalities is lacking.

#### **MRA Chest, Abdomen, and Pelvis Without and With IV Contrast**

MRA of the chest, abdomen, and pelvis without and with IV contrast can be used to evaluate for the presence of an embolic source in the aorta in its entirety. In 1 study, detection of thoracic aorta pathology by contrast-enhanced chest MRA was equivalent in sensitivity, specificity, and diagnostic accuracy compared to noncontrast MRA, although only a single case of thrombus was included in the analysis [43]. On the other hand, in a small analysis which included 9 patients with aortic thrombus, contrast-enhanced MRA had a lower thrombus detection rate compared to a noncontrast examination, although this finding was not statistically significant [44]. Contrast-enhanced MRA of the abdomen has been used for intraluminal thrombus detection in the setting of aneurysms, although comparative data is insufficient [56-58]. Data comparing MRA of the chest, abdomen, and pelvis to other imaging modalities is lacking.

#### **MRA Chest, Abdomen, and Pelvis Without IV Contrast**

Chest, abdomen, and pelvis MRA without IV contrast can be used to evaluate for the presence of an embolic source in the aorta in its entirety. One small study found this examination to have a higher detection rate for aortic thrombus when compared with contrast-enhanced MRA, although the difference was not statistically significant [44]. In another study, sensitivity, specificity, and diagnostic accuracy of unenhanced steady-state free precession MRA were 100% for the detection of thoracic aorta pathology compared to a contrast-enhanced MRA reference standard, however this analysis only included 1 case of mural thrombus [43]. Noncontrast MRA has been used for the detection of abdominal aortic intraluminal thrombus, although there is insufficient data comparing it to contrast-enhanced MRA [56-58]. Data comparing MRA of the chest, abdomen, and pelvis to other imaging modalities is lacking.

#### **MRI Heart Function and Morphology Without and With IV Contrast**

Cardiac MR is a noninvasive imaging study that can reliably detect intracardiac thrombus as well as valvular and neoplastic pathologies. A meta-analysis of 7 studies showed that delayed contrast-enhanced cardiac MR had a pooled sensitivity of 100% and a specificity of 99% for detecting left atrial and left atrial appendage thrombus in patients with atrial fibrillation [45]. In another meta-analysis, there was no significant difference in sensitivity and specificity between cardiac CT and cardiac MR in the detection of left atrial appendage thrombus [29]. Contrast-enhanced cardiac MR had a sensitivity of 88% and a specificity of 99% compared to surgical or pathological

confirmation of left ventricular thrombus [46]. Cardiac MR is also an accurate imaging modality for the evaluation of valvular disease, including aortic and mitral valve vegetations, which can dislodge and result in arterial embolism [37,47]. Additionally, cardiac MR offers detailed soft tissue characterization for the analysis of benign and malignant intracardiac neoplasms [39,48].

### **MRI Heart Function and Morphology Without IV Contrast**

Cardiac MR without contrast provides a detailed anatomic evaluation of the heart chambers. In the workup of embolic sources, the primary role of cardiac MR is in the identification of intracardiac thrombus. A meta-analysis of 7 studies showed that cine cardiac MR had a pooled sensitivity of 91% and a specificity of 93% for detecting left atrial and left atrial appendage thrombus in patients with atrial fibrillation [45]. Furthermore, cine cardiac MR had an 82% sensitivity and a 100% specificity in detecting left ventricle thrombus in postmyocardial infarction patients compared with a standard delayed enhancement cardiac MR [49]. Cardiac MR without contrast is also capable of identifying valvular pathology and cardiac neoplasms, although data on its applicability in the setting of systemic arterial embolism are lacking.

### **US Duplex Doppler Abdomen**

There is no relevant literature to support the use of Doppler US of the abdomen as an initial imaging modality in the evaluation of the source of known embolic lower extremity arterial occlusion. However, some imaging protocols may include limited views of the abdominal aorta, which may detect intraluminal aortic thrombus or significant atherosclerotic disease [56].

### **US Echocardiography Transesophageal**

TEE is an invasive diagnostic study with the ability to detect cardiac pathology predisposed to embolism. TEE has a sensitivity of 93% to 100% and a specificity of 95% to 99% for detecting left atrial appendage thrombus when compared to intraoperative findings [31,50,51]. Furthermore, TEE can evaluate left ventricular systolic dysfunction, spontaneous echo contrast, slow left atrial appendage peak flow velocities, and complex left atrial appendage morphologies, which are all associated with left atrial thrombus and thromboembolic risk [2,4]. In addition, TEE can detect left ventricular thrombus, with 1 study reporting a 40% sensitivity and a 96% specificity for the modality compared to findings at surgery or pathology [46]. Proximal aortic thrombus can also be assessed using TEE, although evaluation is limited by blind spots (distal ascending aorta and proximal aortic arch) owing to air in the trachea [10,13]. Detection of valvular disease and intracardiac neoplasms can also be accomplished with TEE.

### **US Echocardiography Transthoracic Resting**

TTE is a noninvasive imaging modality capable of detecting cardiac pathology susceptible to embolism. TTE is inferior to TEE in the assessment of left atrial appendage thrombus because the transducer is distant from the left atrium when placed on the chest [52]. In 1 study, a cardiac embolic source was detected by TEE in about 40% of patients with normal TTE [53]. In another study, a cardiac embolic source was identified by TTE in 15% of the study group compared with 57% by TEE [54]. Sensitivity and specificity were 23% and 96%, respectively, for the detection of left ventricular thrombus compared to findings at surgery or pathology [46]. In the detection of left ventricle thrombus, contrast-enhanced TTE had a 64% sensitivity and a 99% specificity compared to a delayed enhancement cardiac MR standard [49]. TTE can also be applied for the diagnosis of valvular disease and cardiac neoplasms. There is no evidence to support the use of TTE in the evaluation of aortic thrombus.

### **Variant 4: Known multiorgan system arterial occlusions. Suspected embolic etiology. Next imaging study to determine source.**

The variant assumes that multiorgan arterial occlusions have already been established. Typically, these diagnoses are made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

### **CT Heart Function and Morphology With IV Contrast**

The primary role of cardiac CT in the initial evaluation of multiorgan system arterial embolic occlusion is in the workup of cardiac thrombus as a source. Multiple studies have established high rates of atrial thrombus detection by cardiac CT compared to TEE [18-27]. Meta-analyses have found sensitivities of 96% to 99% and specificities of 92% to 94% for detection of left atrial or left atrial appendage thrombus with cardiac CT compared to a TEE reference standard [28-30]. When compared with intraoperative findings, cardiac CT was 100% sensitive and 85% specific for finding left atrial thrombus [31]. Complex left atrial appendage morphologies, which predispose to thrombus formation, can also be characterized by cardiac CT [32-34]. Additionally, cardiac CT can differentiate left ventricular thrombus from the myocardial wall, with 1 study demonstrating a sensitivity, specificity, and

positive and negative predictive values of 94%, 97%, 94%, and 97%, respectively [35]. Studies have also demonstrated cardiac CT to have comparable accuracy to TEE for identification of vegetations in the setting of infective endocarditis, another potential source of arterial embolism [36-38]. Cardiac CT can identify cardiac neoplasms, both benign and malignant, which have the potential to shed and embolize to distal arterial beds [39,40].

#### **CTA Chest With IV Contrast**

In some conditions or clinical scenarios, there may be a high suspicion that the embolic source is in the thoracic aorta, and a CTA limited to the chest may be diagnostic. As such, multidetector chest CTA with IV contrast can be used to evaluate for at-risk atherosclerotic plaque or the presence of thrombus in the thoracic aorta. CTA is useful in the assessment of the size, extent, and location of an embolic source in the aorta, which can aid in management decisions [13,41]. A number of small studies have used chest CTA to detect aortic mural thrombus that was suspected of embolization [1,12-14,42]. Specific data on the sensitivity and specificity of this imaging modality are lacking.

#### **CTA Chest, Abdomen, and Pelvis With IV contrast**

Multidetector CTA with IV contrast can be used to evaluate for the presence of at-risk atherosclerotic plaque or thrombus in the aorta in its entirety. CTA is useful in the assessment of the size, extent, and location of an embolic source in the aorta, which can aid in management decisions [13,41]. Aortic intraluminal thrombus is oftentimes associated with aneurysm, particularly in the abdomen, which is readily detected by CTA [56,57]. A number of small studies have used CTA to detect aortic mural thrombus that was suspected of embolization [1,12-14,42]. Specific data on the sensitivity and specificity of this imaging modality are lacking.

#### **MRA Chest Without and With IV Contrast**

In some conditions or clinical scenarios, there may be a high suspicion that the embolic source is in the thoracic aorta and an MRA limited to the chest may be diagnostic. In 1 study, detection of thoracic aorta pathology by contrast-enhanced chest MRA was equivalent in sensitivity, specificity, and diagnostic accuracy compared to noncontrast MRA, although only a single case of thrombus was included in the analysis [43]. On the other hand, in a small analysis which included 9 patients with aortic thrombus, contrast-enhanced MRA had a lower thrombus detection rate compared to a noncontrast examination, although this finding was not statistically significant [44]. Data comparing MRA of the chest to other imaging modalities are lacking.

#### **MRA Chest Without IV Contrast**

In some conditions or clinical scenarios, there may be a high suspicion that the embolic source is in the thoracic aorta and an MRA limited to the chest may be diagnostic. One small study found this examination to have a higher detection rate for aortic thrombus when compared with contrast-enhanced MRA, although the difference was not statistically significant [44]. In another study, sensitivity, specificity, and diagnostic accuracy of unenhanced steady-state free precession MRA were 100% for the detection of thoracic aorta pathology compared to a contrast-enhanced MRA reference standard; however, this analysis only included 1 case of mural thrombus [43]. Data comparing MRA of the chest to other imaging modalities is lacking.

#### **MRA Chest, Abdomen, and Pelvis Without and With IV Contrast**

MRA of the chest, abdomen, and pelvis without and with IV contrast can be used to evaluate for the presence of an embolic source in the aorta in its entirety. In 1 study, detection of thoracic aorta pathology by contrast-enhanced chest MRA was equivalent in sensitivity, specificity, and diagnostic accuracy compared to noncontrast MRA, although only a single case of thrombus was included in the analysis [43]. On the other hand, in a small analysis which included 9 patients with aortic thrombus, contrast-enhanced MRA had a lower thrombus detection rate compared to a noncontrast examination, although this finding was not statistically significant [44]. Contrast-enhanced MRA of the abdomen has been used for intraluminal thrombus detection in the setting of aneurysms, although comparative data is insufficient [56-58]. Data comparing MRA of the chest, abdomen, and pelvis to other imaging modalities is lacking.

#### **MRA Chest, Abdomen, and Pelvis Without IV Contrast**

Chest, abdomen, and pelvis MRA without IV contrast can be used to evaluate for the presence of an embolic source in the aorta in its entirety. One small study found this examination to have a higher detection rate for aortic thrombus when compared with contrast-enhanced MRA, although the difference was not statistically significant [44]. In another study, sensitivity, specificity, and diagnostic accuracy of unenhanced steady-state free precession MRA were 100% for the detection of thoracic aorta pathology compared to a contrast-enhanced MRA reference standard, however this analysis only included 1 case of mural thrombus [43]. Noncontrast MRA has been used for the

detection of abdominal aortic intraluminal thrombus, although there is insufficient data comparing it to contrast-enhanced MRA [56-58]. Data comparing MRA of the chest, abdomen, and pelvis to other imaging modalities is lacking.

### **MRI Heart Function and Morphology Without and With IV Contrast**

Cardiac MR is a noninvasive imaging study that can reliably detect intracardiac thrombus as well as valvular and neoplastic pathologies. A meta-analysis of 7 studies showed that delayed contrast-enhanced cardiac MR had a pooled sensitivity of 100% and a specificity of 99% for detecting left atrial and left atrial appendage thrombus in patients with atrial fibrillation [45]. In another meta-analysis, there was no significant difference in sensitivity and specificity between cardiac CT and cardiac MR in the detection of left atrial appendage thrombus [29]. Contrast-enhanced cardiac MR had a sensitivity of 88% and a specificity of 99% compared to surgical or pathological confirmation of left ventricular thrombus [46]. Cardiac MR is also an accurate imaging modality for the evaluation of valvular disease, including aortic and mitral valve vegetations, which can dislodge and result in arterial embolism [37,47]. Additionally, cardiac MR offers detailed soft tissue characterization for the analysis of benign and malignant intracardiac neoplasms [39,48].

### **MRI Heart Function and Morphology Without IV Contrast**

Cardiac MR without contrast provides a detailed anatomic evaluation of the heart chambers. In the workup of embolic sources, the primary role of cardiac MR is in the identification of intracardiac thrombus. A meta-analysis of 7 studies showed that cine cardiac MR had a pooled sensitivity of 91% and a specificity of 93% for detecting left atrial and left atrial appendage thrombus in patients with atrial fibrillation [45]. Furthermore, cine cardiac MR had an 82% sensitivity and a 100% specificity in detecting left ventricle thrombus in postmyocardial infarction patients compared with a standard delayed enhancement cardiac MR [49]. Cardiac MR without contrast is also capable of identifying valvular pathology and cardiac neoplasms, although data on its applicability in the setting of systemic arterial thromboembolism are lacking.

### **US Duplex Doppler Abdomen**

There is no relevant literature to support the use of Doppler US of the abdomen as an initial imaging modality in the evaluation of the source of known embolic multiorgan arterial occlusion. However, some imaging protocols may include limited views of the abdominal aorta, which may detect intraluminal aortic thrombus or significant atherosclerotic disease [56].

### **US Echocardiography Transesophageal**

TEE is an invasive diagnostic study with the ability to detect cardiac pathology predisposed to embolism. TEE has a sensitivity of 93% to 100% and a specificity of 95% to 99% for detecting left atrial appendage thrombus when compared with intraoperative findings [31,50,51]. Furthermore, TEE can evaluate left ventricular systolic dysfunction, spontaneous echo contrast, slow left atrial appendage peak flow velocities, and complex left atrial appendage morphologies, which are all associated with left atrial thrombus and thromboembolic risk [2,4]. In addition, TEE can detect left ventricular thrombus with 1 study reporting a 40% sensitivity and a 96% specificity for the modality compared to findings at surgery or pathology [46]. Proximal aortic thrombus can also be assessed using TEE, although evaluation is limited by blind spots (distal ascending aorta and proximal aortic arch) owing to air in the trachea [10,13]. Detection of valvular disease and intracardiac neoplasms can also be accomplished with TEE.

### **US Echocardiography Transthoracic Resting**

TTE is a noninvasive imaging modality capable of detecting cardiac pathology susceptible to embolism. TTE is inferior to TEE in the assessment of left atrial appendage thrombus because the transducer is distant from the left atrium when placed on the chest [52]. In 1 study, a cardiac embolic source was detected by TEE in about 40% of patients with normal TTE [53]. In another study, a cardiac embolic source was identified by TTE in 15% of the study group compared with 57% by TEE [54]. Sensitivity and specificity were 23% and 96%, respectively, for the detection of left ventricular thrombus compared to findings at surgery or pathology [46]. In the detection of left ventricle thrombus, contrast-enhanced TTE had a 64% sensitivity and a 99% specificity compared to a delayed enhancement cardiac MR standard [49]. TTE can also be applied for the diagnosis of valvular disease and cardiac neoplasms. There is no evidence to support the use of TTE in the evaluation of aortic thrombus.

### **Summary of Recommendations**

- **Variation 1:** TEE, TTE, MRI heart function and morphology without and with IV contrast, MRI heart function and morphology without IV contrast, and CT heart function and morphology with IV contrast are usually

appropriate for determining the cardiac source of a known upper extremity arterial occlusion with a suspected embolic etiology. These procedures are equivalent alternatives (ie, only one procedure will be ordered to provide the clinical information to effectively manage the patient's care). Furthermore, CTA of the chest with IV contrast and MRA of the chest without and with IV contrast are equivalent alternatives that are usually appropriate for determining a central arterial source of a known upper extremity arterial occlusion with a suspected embolic etiology. During the workup of an upper extremity embolic source, cardiac and cross-sectional angiographic imaging may be complementary (both are performed) in certain clinical scenarios/conditions.

- **Variation 2:** TEE, TTE, MRI heart function and morphology without and with IV contrast, MRI heart function and morphology without IV contrast, and CT heart function and morphology with IV contrast are usually appropriate for determining the cardiac source of a known visceral arterial occlusion with a suspected embolic etiology. These procedures are equivalent alternatives (ie, only one procedure will be ordered to provide the clinical information to effectively manage the patient's care). Furthermore, CTA of the chest and abdomen with IV contrast and MRA of the chest and abdomen without and with IV contrast are equivalent alternatives that are usually appropriate for determining a central arterial source of a known visceral arterial occlusion with a suspected embolic etiology. CTA of the chest with IV contrast is usually appropriate in some clinical situations in which there is a high suspicion of the embolic source originating in the thoracic aorta. During the workup of a visceral embolic source, cardiac and cross-sectional angiographic imaging may be complementary (both are performed) in certain clinical scenarios/conditions.
- **Variation 3:** TEE, TTE, MRI heart function and morphology without and with IV contrast, MRI heart function and morphology without IV contrast, and CT heart function and morphology with IV contrast are usually appropriate for determining the cardiac source of a known lower extremity arterial occlusion with a suspected embolic etiology. These procedures are equivalent alternatives (ie, only one procedure will be ordered to provide the clinical information to effectively manage the patient's care). Furthermore, CTA of the chest, abdomen, and pelvis with IV contrast and MRA of the chest, abdomen, and pelvis without and with IV contrast are equivalent alternatives that are usually appropriate for determining a central arterial source of a known lower extremity arterial occlusion with a suspected embolic etiology. CTA of the chest with IV contrast or MRA of the chest without and with IV contrast are usually appropriate in some clinical situations in which there is a high suspicion of the embolic source originating in the thoracic aorta. During the workup of a lower extremity embolic source, cardiac and cross-sectional angiographic imaging may be complementary (both are performed) in certain clinical scenarios/conditions.
- **Variation 4:** TEE, TTE, MRI heart function and morphology without and with IV contrast, MRI heart function and morphology without IV contrast, and CT heart function and morphology with IV contrast are usually appropriate for determining the cardiac source of known multiorgan system arterial occlusions with a suspected embolic etiology. These procedures are equivalent alternatives (ie, only one procedure will be ordered to provide the clinical information to effectively manage the patient's care). Furthermore, CTA of the chest, abdomen, and pelvis with IV contrast, MRA of the chest, abdomen, and pelvis without and with IV contrast, and MRA of the chest, abdomen, and pelvis without IV contrast are equivalent alternatives that are usually appropriate for determining a central arterial source of known multiorgan system arterial occlusions with a suspected embolic etiology. CTA of the chest with IV contrast or MRA of the chest without and with IV contrast are usually appropriate in some clinical situations when there is a high suspicion of the embolic source originating in the thoracic aorta. During the workup of a multiorgan system embolic source, cardiac and cross-sectional angiographic imaging may be complementary (both are performed) in certain clinical scenarios/conditions.

### Supporting Documents

The evidence table, literature search, and appendix for this topic are available at <https://acsearch.acr.org/list>. The appendix includes the strength of evidence assessment and the final rating round tabulations for each recommendation.

For additional information on the Appropriateness Criteria methodology and other supporting documents go to [www.acr.org/ac](http://www.acr.org/ac).

## Appropriateness Category Names and Definitions

Appropriateness Category Name	Appropriateness Rating	Appropriateness Category Definition
Usually Appropriate	7, 8, or 9	The imaging procedure or treatment is indicated in the specified clinical scenarios at a favorable risk-benefit ratio for patients.
May Be Appropriate	4, 5, or 6	The imaging procedure or treatment may be indicated in the specified clinical scenarios as an alternative to imaging procedures or treatments with a more favorable risk-benefit ratio, or the risk-benefit ratio for patients is equivocal.
May Be Appropriate (Disagreement)	5	The individual ratings are too dispersed from the panel median. The different label provides transparency regarding the panel's recommendation. "May be appropriate" is the rating category and a rating of 5 is assigned.
Usually Not Appropriate	1, 2, or 3	The imaging procedure or treatment is unlikely to be indicated in the specified clinical scenarios, or the risk-benefit ratio for patients is likely to be unfavorable.

## Relative Radiation Level Information

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Patients in the pediatric age group are at inherently higher risk from exposure, because of both organ sensitivity and longer life expectancy (relevant to the long latency that appears to accompany radiation exposure). For these reasons, the RRL dose estimate ranges for pediatric examinations are lower as compared with those specified for adults (see Table below). Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® [Radiation Dose Assessment Introduction](#) document [60].

Relative Radiation Level Designations		
Relative Radiation Level*	Adult Effective Dose Estimate Range	Pediatric Effective Dose Estimate Range
○	0 mSv	0 mSv
⊕	<0.1 mSv	<0.03 mSv
⊕⊕	0.1-1 mSv	0.03-0.3 mSv
⊕⊕⊕	1-10 mSv	0.3-3 mSv
⊕⊕⊕⊕	10-30 mSv	3-10 mSv
⊕⊕⊕⊕⊕	30-100 mSv	10-30 mSv

\*RRL assignments for some of the examinations cannot be made, because the actual patient doses in these procedures vary as a function of a number of factors (eg, region of the body exposed to ionizing radiation, the imaging guidance that is used). The RRLs for these examinations are designated as "Varies."

## References

1. Weiss S, Buhlmann R, von Allmen RS, et al. Management of floating thrombus in the aortic arch. *J Thorac Cardiovasc Surg* 2016;152:810-7.

2. Kumagai T, Matsuura Y, Yamamoto T, Ugawa Y, Fukushima T. Risk factors for left atrial thrombus from transesophageal echocardiography findings in ischemic stroke patients. *Fukushima J Med Sci* 2014;60:154-8.
3. Wysokinski WE, Ammash N, Sobande F, Kalsi H, Hodge D, McBane RD. Predicting left atrial thrombi in atrial fibrillation. *Am Heart J* 2010;159:665-71.
4. Yamamoto M, Seo Y, Kawamatsu N, et al. Complex left atrial appendage morphology and left atrial appendage thrombus formation in patients with atrial fibrillation. *Circ Cardiovasc Imaging* 2014;7:337-43.
5. Gianstefani S, Douiri A, Delithanasis I, et al. Incidence and predictors of early left ventricular thrombus after ST-elevation myocardial infarction in the contemporary era of primary percutaneous coronary intervention. *Am J Cardiol* 2014;113:1111-6.
6. McCarthy CP, Vaduganathan M, McCarthy KJ, Januzzi JL, Jr., Bhatt DL, McEvoy JW. Left Ventricular Thrombus After Acute Myocardial Infarction: Screening, Prevention, and Treatment. *JAMA Cardiol* 2018;3:642-49.
7. Sordelli C, Fele N, Mocerino R, et al. Infective Endocarditis: Echocardiographic Imaging and New Imaging Modalities. *J Cardiovasc Echogr* 2019;29:149-55.
8. Wintersperger BJ, Becker CR, Gulbins H, et al. Tumors of the cardiac valves: imaging findings in magnetic resonance imaging, electron beam computed tomography, and echocardiography. *Eur Radiol* 2000;10:443-9.
9. O'Connell JB, Quinones-Baldrich WJ. Proper evaluation and management of acute embolic versus thrombotic limb ischemia. *Semin Vasc Surg* 2009;22:10-6.
10. Yoo SM, Lee HY, White CS. MDCT evaluation of acute aortic syndrome. *Radiol Clin North Am* 2010;48:67-83.
11. Tsilimparis N, Hanack U, Pisimisis G, Yousefi S, Wintzer C, Ruckert RI. Thrombus in the non-aneurysmal, non-atherosclerotic descending thoracic aorta--an unusual source of arterial embolism. *Eur J Vasc Endovasc Surg* 2011;41:450-7.
12. Klang E, Kerpel A, Soffer S, et al. CT imaging features of symptomatic and asymptomatic floating aortic thrombus. *Clin Radiol* 2018;73:323 e9-23 e14.
13. Pagni S, Trivedi J, Ganzel BL, et al. Thoracic aortic mobile thrombus: is there a role for early surgical intervention? *Ann Thorac Surg* 2011;91:1875-81.
14. Boufi M, Mameli A, Compes P, Hartung O, Alimi YS. Elective stent-graft treatment for the management of thoracic aorta mural thrombus. *Eur J Vasc Endovasc Surg* 2014;47:335-41.
15. American College of Radiology. ACR–NASCI–SIR–SPR Practice Parameter for the Performance and Interpretation of Body Computed Tomography Angiography (CTA). Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/body-cta.pdf>. Accessed September 30, 2022.
16. Fleischmann D, Mitchell RS, Miller DC. Acute aortic syndromes: new insights from electrocardiographically gated computed tomography. *Semin Thorac Cardiovasc Surg* 2008;20:340-7.
17. Holloway BJ, Rosewarne D, Jones RG. Imaging of thoracic aortic disease. *Br J Radiol* 2011;84 Spec No 3:S338-54.
18. Hur J, Kim YJ, Lee HJ, et al. Left atrial appendage thrombi in stroke patients: detection with two-phase cardiac CT angiography versus transesophageal echocardiography. *Radiology* 2009;251:683-90.
19. Hur J, Kim YJ, Lee HJ, et al. Dual-enhanced cardiac CT for detection of left atrial appendage thrombus in patients with stroke: a prospective comparison study with transesophageal echocardiography. *Stroke* 2011;42:2471-7.
20. Hur J, Kim YJ, Lee HJ, et al. Cardioembolic stroke: dual-energy cardiac CT for differentiation of left atrial appendage thrombus and circulatory stasis. *Radiology* 2012;263:688-95.
21. Teunissen C, Habets J, Velthuis BK, Cramer MJ, Loh P. Double-contrast, single-phase computed tomography angiography for ruling out left atrial appendage thrombus prior to atrial fibrillation ablation. *Int J Cardiovasc Imaging* 2017;33:121-28.
22. Kapa S, Martinez MW, Williamson EE, et al. ECG-gated dual-source CT for detection of left atrial appendage thrombus in patients undergoing catheter ablation for atrial fibrillation. *J Interv Card Electrophysiol* 2010;29:75-81.
23. Zhai Z, Tang M, Zhang S, et al. Transoesophageal echocardiography prior to catheter ablation could be avoided in atrial fibrillation patients with a low risk of stroke and without filling defects in the late-phase MDCT scan: A retrospective analysis of 783 patients. *Eur Radiol* 2018;28:1835-43.
24. Ikegami Y, Tanimoto K, Inagawa K, et al. Identification of Left Atrial Appendage Thrombi in Patients With Persistent and Long-Standing Persistent Atrial Fibrillation Using Intra-Cardiac Echocardiography and Cardiac Computed Tomography. *Circ J* 2017;82:46-52.

25. Kantarci M, Ogul H, Sade R, Aksakal E, Colak A, Tanboga IH. Circulatory Stasis or Thrombus in Left Atrial Appendage, An Easy Diagnostic Solution. *J Comput Assist Tomogr* 2019;43:406-09.
26. Bilchick KC, Mealor A, Gonzalez J, et al. Effectiveness of integrating delayed computed tomography angiography imaging for left atrial appendage thrombus exclusion into the care of patients undergoing ablation of atrial fibrillation. *Heart Rhythm* 2016;13:12-9.
27. Martinez MW, Kirsch J, Williamson EE, et al. Utility of nongated multidetector computed tomography for detection of left atrial thrombus in patients undergoing catheter ablation of atrial fibrillation. *JACC Cardiovasc Imaging* 2009;2:69-76.
28. Romero J, Husain SA, Kelesidis I, Sanz J, Medina HM, Garcia MJ. Detection of left atrial appendage thrombus by cardiac computed tomography in patients with atrial fibrillation: a meta-analysis. *Circ Cardiovasc Imaging* 2013;6:185-94.
29. Vira T, Pechlivanoglou P, Connelly K, Wijeyesundera HC, Roifman I. Cardiac computed tomography and magnetic resonance imaging vs. transoesophageal echocardiography for diagnosing left atrial appendage thrombi. *Europace* 2019;21:e1-e10.
30. Zou H, Zhang Y, Tong J, Liu Z. Multidetector computed tomography for detecting left atrial/left atrial appendage thrombus: a meta-analysis. *Intern Med J* 2015;45:1044-53.
31. Choi BH, Ko SM, Hwang HK, et al. Detection of left atrial thrombus in patients with mitral stenosis and atrial fibrillation: retrospective comparison of two-phase computed tomography, transoesophageal echocardiography and surgical findings. *Eur Radiol* 2013;23:2944-53.
32. Korhonen M, Muuronen A, Arponen O, et al. Left atrial appendage morphology in patients with suspected cardiogenic stroke without known atrial fibrillation. *PLoS One* 2015;10:e0118822.
33. Hozawa M, Morino Y, Matsumoto Y, et al. 3D-computed tomography to compare the dimensions of the left atrial appendage in patients with normal sinus rhythm and those with paroxysmal atrial fibrillation. *Heart Vessels* 2018;33:777-85.
34. Dieker W, Behnes M, Fastner C, et al. Impact of left atrial appendage morphology on thrombus formation after successful left atrial appendage occlusion: Assessment with cardiac-computed-tomography. *Sci Rep* 2018;8:1670.
35. Bittencourt MS, Achenbach S, Marwan M, et al. Left ventricular thrombus attenuation characterization in cardiac computed tomography angiography. *J Cardiovasc Comput Tomogr* 2012;6:121-6.
36. Kim IC, Chang S, Hong GR, et al. Comparison of Cardiac Computed Tomography With Transesophageal Echocardiography for Identifying Vegetation and Intracardiac Complications in Patients With Infective Endocarditis in the Era of 3-Dimensional Images. *Circ Cardiovasc Imaging* 2018;11:e006986.
37. Erba PA, Pizzi MN, Roque A, et al. Multimodality Imaging in Infective Endocarditis: An Imaging Team Within the Endocarditis Team. *Circulation* 2019;140:1753-65.
38. Feuchtner GM, Stolzmann P, Dichtl W, et al. Multislice computed tomography in infective endocarditis: comparison with transesophageal echocardiography and intraoperative findings. *J Am Coll Cardiol* 2009;53:436-44.
39. Araoz PA, Mulvagh SL, Tazelaar HD, Julsrud PR, Breen JF. CT and MR imaging of benign primary cardiac neoplasms with echocardiographic correlation. *Radiographics* 2000;20:1303-19.
40. Kassop D, Donovan MS, Cheezum MK, et al. Cardiac Masses on Cardiac CT: A Review. *Curr Cardiovasc Imaging Rep* 2014;7:9281.
41. Mesurole B, Qanadli SD, Merad M, El Hajjam M, Mignon F, Lacombe P. Dual-slice helical CT of the thoracic aorta. *J Comput Assist Tomogr* 2000;24:548-56.
42. Ryoo S, Chung JW, Lee MJ, et al. An Approach to Working Up Cases of Embolic Stroke of Undetermined Source. *J Am Heart Assoc* 2016;5:e002975.
43. Krishnam MS, Tomasian A, Malik S, Desphande V, Laub G, Ruehm SG. Image quality and diagnostic accuracy of unenhanced SSFP MR angiography compared with conventional contrast-enhanced MR angiography for the assessment of thoracic aortic diseases. *Eur Radiol* 2010;20:1311-20.
44. Gebker R, Gomaa O, Schnackenburg B, Rebakowski J, Fleck E, Nagel E. Comparison of different MRI techniques for the assessment of thoracic aortic pathology: 3D contrast enhanced MR angiography, turbo spin echo and balanced steady state free precession. *Int J Cardiovasc Imaging* 2007;23:747-56.
45. Chen J, Zhang H, Zhu D, Wang Y, Byanju S, Liao M. Cardiac MRI for detecting left atrial/left atrial appendage thrombus in patients with atrial fibrillation : Meta-analysis and systematic review. *Herz* 2019;44:390-97.



46. Srichai MB, Junor C, Rodriguez LL, et al. Clinical, imaging, and pathological characteristics of left ventricular thrombus: a comparison of contrast-enhanced magnetic resonance imaging, transthoracic echocardiography, and transesophageal echocardiography with surgical or pathological validation. *Am Heart J* 2006;152:75-84.
47. Rustemli A, Bhatti TK, Wolff SD. Evaluating cardiac sources of embolic stroke with MRI. *Echocardiography* 2007;24:301-8; discussion 08.
48. Motwani M, Kidambi A, Herzog BA, Uddin A, Greenwood JP, Plein S. MR imaging of cardiac tumors and masses: a review of methods and clinical applications. *Radiology* 2013;268:26-43.
49. Weinsaft JW, Kim J, Medicherla CB, et al. Echocardiographic Algorithm for Post-Myocardial Infarction LV Thrombus: A Gatekeeper for Thrombus Evaluation by Delayed Enhancement CMR. *JACC Cardiovasc Imaging* 2016;9:505-15.
50. He YQ, Liu L, Zhang MC, Zeng H, Yang P. Dual-Energy Computed Tomography-Enabled Material Separation in Diagnosing Left Atrial Appendage Thrombus. *Tex Heart Inst J* 2019;46:107-14.
51. Kumar V, Nanda NC. Is it time to move on from two-dimensional transesophageal to three-dimensional transthoracic echocardiography for assessment of left atrial appendage? Review of existing literature. *Echocardiography* 2012;29:112-6.
52. Nakanishi K, Homma S. Role of echocardiography in patients with stroke. *J Cardiol* 2016;68:91-9.
53. de Bruijn SF, Agema WR, Lammers GJ, et al. Transesophageal echocardiography is superior to transthoracic echocardiography in management of patients of any age with transient ischemic attack or stroke. *Stroke* 2006;37:2531-4.
54. Pearson AC, Labovitz AJ, Tatineni S, Gomez CR. Superiority of transesophageal echocardiography in detecting cardiac source of embolism in patients with cerebral ischemia of uncertain etiology. *J Am Coll Cardiol* 1991;17:66-72.
55. Ginsburg M, Obara P, Lambert DL, et al. ACR Appropriateness Criteria® Imaging of Mesenteric Ischemia. *J Am Coll Radiol* 2018;15:S332-S40.
56. Labruto F, Blomqvist L, Swedenborg J. Imaging the intraluminal thrombus of abdominal aortic aneurysms: techniques, findings, and clinical implications. *J Vasc Interv Radiol* 2011;22:1069-75; quiz 75.
57. Litmanovich D, Bankier AA, Cantin L, Raptopoulos V, Boiselle PM. CT and MRI in diseases of the aorta. *AJR Am J Roentgenol* 2009;193:928-40.
58. Nguyen VL, Leiner T, Hellenthal FA, et al. Abdominal aortic aneurysms with high thrombus signal intensity on magnetic resonance imaging are associated with high growth rate. *Eur J Vasc Endovasc Surg* 2014;48:676-84.
59. Weiss CR, Azene EM, Majdalany BS, et al. ACR Appropriateness Criteria® Sudden Onset of Cold, Painful Leg. *J Am Coll Radiol* 2017;14:S307-S13.
60. American College of Radiology. ACR Appropriateness Criteria® Radiation Dose Assessment Introduction. Available at: <https://www.acr.org/-/media/ACR/Files/Appropriateness-Criteria/RadiationDoseAssessmentIntro.pdf>. Accessed September 30, 2022.

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.