

American College of Radiology ACR Appropriateness Criteria®

Clinical Condition: Chronic Elbow Pain

Variant 1: Evaluation for chronic elbow pain. First test.

Radiologic Procedure	Rating	Comments	RRL*
X-ray elbow	9		☼
MRI elbow without IV contrast	1		○
MRI elbow without and with IV contrast	1		○
MR arthrography elbow	1		○
CT elbow without IV contrast	1		☼☼
CT elbow with IV contrast	1		☼☼
CT elbow without and with IV contrast	1		☼☼
CT arthrography elbow	1		☼☼
US elbow	1		○
3-phase bone scan elbow	1		☼☼☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 2: Mechanical symptoms (locking, clicking, limited motion); suspect intra-articular osteocartilaginous body or synovial abnormality; radiographs nondiagnostic.

Radiologic Procedure	Rating	Comments	RRL*
MRI elbow without IV contrast	9	Either routine MRI or MR arthrogram is appropriate. Use of this procedure depends on availability and expertise. If effusion is present, procedure without contrast is preferred.	○
MR arthrography elbow	9	Either routine MRI or MR arthrogram is appropriate. Use of this procedure depends on availability, expertise, and local conditions.	○
CT elbow without IV contrast	8	This is an alternative procedure.	☼☼
CT arthrography elbow	8	This is an alternative procedure.	☼☼
US elbow	3		○
MRI elbow without and with IV contrast	1		○
CT elbow with IV contrast	1		☼☼
CT elbow without and with IV contrast	1		☼☼
3-phase bone scan elbow	1		☼☼☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Clinical Condition: Chronic Elbow Pain

Variant 3: Suspect occult fracture or other bone abnormality; radiographs nondiagnostic.

Radiologic Procedure	Rating	Comments	RRL*
MRI elbow without IV contrast	9		○
CT elbow without IV contrast	5		☢☢
3-phase bone scan elbow	4		☢☢☢
MR arthrography elbow	2		○
CT arthrography elbow	2		☢☢
MRI elbow without and with IV contrast	1		○
CT elbow with IV contrast	1		☢☢
CT elbow without and with IV contrast	1		☢☢
US elbow	1		○
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 4: Assess stability of osteochondral injury; radiographs nondiagnostic.

Radiologic Procedure	Rating	Comments	RRL*
MRI elbow without IV contrast	9	This is an alternative procedure.	○
MR arthrography elbow	9	This is an alternative procedure.	○
CT arthrography elbow	8	This is an alternative procedure.	☢☢
MRI elbow without and with IV contrast	5	This procedure may be appropriate but there was disagreement among panel members on the appropriateness rating as defined by the panel's median rating.	○
CT elbow without IV contrast	3		☢☢
CT elbow with IV contrast	1		☢☢
CT elbow without and with IV contrast	1		☢☢
US elbow	1		○
3-phase bone scan elbow	1		☢☢☢
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Clinical Condition: Chronic Elbow Pain

Variant 5: Palpable soft-tissue mass; radiographs nondiagnostic.

Radiologic Procedure	Rating	Comments	RRL*
MRI elbow without and with IV contrast	9	Contrast may not be necessary in all cases.	○
MRI elbow without IV contrast	7	This is an alternative procedure for some masses.	○
US elbow	7	This is an alternative procedure.	○
CT elbow with IV contrast	3		☢☢
CT elbow without IV contrast	3		☢☢
CT elbow without and with IV contrast	2		☢☢
3-phase bone scan elbow	1		☢☢☢
CT arthrography elbow	1		☢☢
MR arthrography elbow	1		○
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 6: Suspect chronic epicondylitis, refractory to empirical treatment; radiographs nondiagnostic.

Radiologic Procedure	Rating	Comments	RRL*
MRI elbow without IV contrast	9	This is an alternative procedure.	○
US elbow	8	This is an alternative procedure.	○
MRI elbow without and with IV contrast	2		○
MR arthrography elbow	2		○
3-phase bone scan elbow	2		☢☢☢
CT elbow without IV contrast	1		☢☢
CT elbow with IV contrast	1		☢☢
CT elbow without and with IV contrast	1		☢☢
CT arthrography elbow	1		☢☢
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Clinical Condition: Chronic Elbow Pain

Variant 7: Suspect collateral ligament tear; radiographs nondiagnostic.

Radiologic Procedure	Rating	Comments	RRL*
MRI elbow without IV contrast	9	Either routine MRI or MR arthrogram is appropriate. Use of this procedure depends on availability, expertise, and local conditions.	○
MR arthrography elbow	9	Either routine MRI or MR arthrogram is appropriate. Use of this procedure depends on availability, expertise, and local conditions.	○
X-ray elbow stress views	6		☼
US elbow	6		○
CT arthrography elbow	5		☼☼
CT elbow without IV contrast	2		☼☼
MRI elbow without and with IV contrast	1		○
CT elbow with IV contrast	1		☼☼
CT elbow without and with IV contrast	1		☼☼
3-phase bone scan elbow	1		☼☼☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 8: Suspect biceps tendon tear; radiographs nondiagnostic.

Radiologic Procedure	Rating	Comments	RRL*
MRI elbow without IV contrast	9	This is an alternative procedure.	○
US elbow	8	This is an alternative procedure.	○
MRI elbow without and with IV contrast	1		○
MR arthrography elbow	1		○
CT elbow without IV contrast	1		☼☼
CT elbow with IV contrast	1		☼☼
CT elbow without and with IV contrast	1		☼☼
CT arthrography elbow	1		☼☼
3-phase bone scan elbow	1		☼☼☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Clinical Condition: Chronic Elbow Pain

Variant 9: Suspect nerve abnormality; radiographs nondiagnostic.

Radiologic Procedure	Rating	Comments	RRL*
MRI elbow without IV contrast	9	This is an alternative procedure.	○
US elbow	8	This is an alternative procedure.	○
MRI elbow without and with IV contrast	2		○
MR arthrography elbow	1		○
CT elbow without IV contrast	1		☢☢
CT elbow with IV contrast	1		☢☢
CT elbow without and with IV contrast	1		☢☢
CT arthrography elbow	1		☢☢
3-phase bone scan elbow	1		☢☢☢
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 10: Elbow stiffness; suspect heterotopic ossification/osteophytosis by radiography. Next test.

Radiologic Procedure	Rating	Comments	RRL*
CT elbow without IV contrast	9		☢☢
MRI elbow without IV contrast	5		○
3-phase bone scan elbow	5	This procedure may be appropriate but there was disagreement among panel members on the appropriateness rating as defined by the panel's median rating.	☢☢☢
CT elbow with IV contrast	1		☢☢
CT elbow without and with IV contrast	1		☢☢
MRI elbow without and with IV contrast	1		○
US elbow	1		○
MR arthrography elbow	1		○
CT arthrography elbow	1		☢☢
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Clinical Condition: Chronic Elbow Pain

Variant 11: Suspect inflammatory arthritis or bursitis; radiographs obtained.

Radiologic Procedure	Rating	Comments	RRL*
MRI elbow without and with IV contrast	9	This is an alternative procedure.	○
MRI elbow without IV contrast	8	This is an alternative procedure.	○
US elbow	7	This is an alternative procedure.	○
3-phase bone scan elbow	3		☢☢☢
CT elbow without IV contrast	1		☢☢
CT elbow with IV contrast	1		☢☢
CT elbow without and with IV contrast	1		☢☢
MR arthrography elbow	1		○
CT arthrography elbow	1		☢☢
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

CHRONIC ELBOW PAIN

Expert Panel on Musculoskeletal Imaging: Curtis W. Hayes, MD¹; Catherine C. Roberts, MD²; Jenny T. Bencardino, MD³; Marc Appel, MD⁴; Erin Arnold, MD⁵; Steven J. Bacceti, MD⁶; Eric Y. Chang, MD⁷; Michael G. Fox, MD⁸; Ian Blair Fries, MD⁹; Bennett S. Greenspan, MD, MS¹⁰; Mary G. Hochman, MD¹¹; Jon A. Jacobson, MD¹²; Douglas N. Mintz, MD¹³; Gary Mlady, MD¹⁴; Mark D. Murphey, MD¹⁵; Joel S. Newman, MD¹⁶; Zehava Sadka Rosenberg, MD¹⁷; David A. Rubin, MD¹⁸; Kirstin M. Small, MD¹⁹; Barbara N. Weissman, MD.²⁰

Summary of Literature Review

Introduction/Overview of Imaging Modalities

Chronic elbow pain may be caused by a variety of osseous abnormalities, soft-tissue abnormalities, or both. Radiography is most helpful for assessing bony structures, calcification, or ossification. Computed tomography (CT), magnetic resonance imaging (MRI), and ultrasonography (US) are the most commonly used additional imaging studies. Although CT and US can be used for specific indications, MRI effectively demonstrates most abnormalities in the elbow. MRI provides important diagnostic information for evaluating the elbow in many different conditions, including collateral ligament injury; epicondylitis; injury to the biceps and triceps tendons; and abnormality of the ulnar, radial, or median nerve; and for evaluating masses about the elbow joint [1-13]. There are a lack of studies showing the sensitivity and specificity of MRI in many of these conditions; most studies demonstrate MRI findings in patients either known or highly likely to have a specific condition. Imaging choices will be considered for a variety of specific clinical variants in the following sections.

Discussion of Imaging Modalities by Variant

Variant 1: Evaluation for chronic elbow pain. First test.

Radiographs are the best initial imaging study for a patient with chronic elbow pain. In some cases, radiographs may reveal the definitive cause of the problem (eg, osseocartilaginous intra-articular body (IAB), osteophytes, heterotopic ossification, osteochondritis dissecans, or calcification in and around the joint in the form of hydroxyapatite deposition or calcium pyrophosphate crystal deposition). Exclusion of an osseous abnormality with radiographs may be helpful when conservative therapy is planned. Radiographs are also a useful adjunct to interpretation if musculoskeletal MRI is subsequently performed [14].

Instability may be difficult to diagnose clinically and some authors suggest that radiographic stress views can be helpful in diagnosing valgus instability [15]. Specific techniques for obtaining these radiographs have been recommended [15] and comparison to the unaffected side may also be of help [16].

Variant 2: Mechanical symptoms (locking, clicking, limited motion); suspect intra-articular osteocartilaginous body or synovial abnormality; radiographs nondiagnostic.

CT and CT arthrography with single-contrast (iodinated contrast or air) and double-contrast (iodinated contrast and air) techniques are superior to radiography for detecting a chondral or osteochondral lesion or IAB [17]. Both of these studies have limitations; a small IAB may be obscured by contrast or confused with air bubbles (double-contrast arthrography). A CT air arthrogram can avoid confusion of air bubbles with IABs. Regardless of method, detection of an IAB is limited by its size and location within the elbow joint, although detection is enhanced by the presence of joint effusion [5].

¹Principal Author, VCU Health System, Richmond, Virginia. ²Panel Chair, Mayo Clinic, Phoenix, Arizona. ³Panel Vice-chair, NYU Hospital for Joint Diseases, New York, New York. ⁴James J. Peters VA Medical Center, Bronx, New York, American Academy of Orthopaedic Surgeons. ⁵Orthopaedics and Rheumatology of the North Shore, Skokie, Illinois, American College of Rheumatology. ⁶UMass Memorial Medical Center, Worcester, Massachusetts. ⁷VA San Diego Healthcare System, San Diego, California. ⁸University of Virginia Health System, Charlottesville, Virginia. ⁹Bone, Spine and Hand Surgery, Chartered, Brick, NJ, American Academy of Orthopaedic Surgeons. ¹⁰Medical College of Georgia, Augusta, Georgia. ¹¹Beth Israel Deaconess Medical Center, Boston, Massachusetts. ¹²University of Michigan Medical Center, Ann Arbor, Michigan. ¹³Hospital for Special Surgery, New York, New York. ¹⁴University of New Mexico, Albuquerque, New Mexico. ¹⁵Uniformed Services University of the Health Sciences, Bethesda, Maryland. ¹⁶New England Baptist Hospital, Boston, Massachusetts. ¹⁷Hospital for Joint Diseases, New York, New York. ¹⁸Washington University School of Medicine, Saint Louis, Missouri. ¹⁹Brigham & Women's Hospital, Boston, Massachusetts. ²⁰Specialty Chair, Brigham & Women's Hospital, Boston, Massachusetts.

The American College of Radiology seeks and encourages collaboration with other organizations on the development of the ACR Appropriateness Criteria through society representation on expert panels. Participation by representatives from collaborating societies on the expert panel does not necessarily imply individual or society endorsement of the final document.

Reprint requests to: publications@acr.org

Some authors recommend MRI as the initial study [10,12]. MRI following direct intra-articular contrast administration (MR arthrography) is preferred to routine MRI for diagnosing an IAB [18,19]. MRI and MR arthrography are limited in their ability to detect purely cartilaginous lesions in the elbow [20].

Variante 3: Suspect occult fracture or other bone abnormality; radiographs nondiagnostic.

Initial evaluation should begin with radiography, but if the radiographs are nondiagnostic, both traumatic and stress fractures may be identified with MRI and 3-phase bone scan [21-23]. CT may have utility in detection and further evaluation of acute traumatic elbow fractures [24], as well as limited usefulness in identifying stress fractures at the elbow [21]. US is helpful in demonstrating lipohemarthrosis in cases of occult elbow fractures in children [25].

Variante 4: Assess stability of osteochondral injury; radiographs nondiagnostic.

Some authors advocate MRI as the initial study for suspected osteochondral fracture [10,12]. Intravenous (IV) or intra-articular contrast has some limited usefulness in the diagnosis of chondral injuries on MRI [20]. Both CT and MRI can assess for osteochondral fragment stability [26]. MRI following direct intra-articular contrast administration (MR arthrography) can play a role in improving evaluation of stability of an osteochondral lesion [18,19]. Similarly, CT arthrography provides more diagnostic information than standard CT in this setting. Although US may show osteochondral abnormalities in some situations [27], MRI offers a more comprehensive evaluation. IV contrast (indirect arthrography) used in conjunction with MRI can be useful in situations where direct arthrography is not an option.

Variante 5: Palpable soft-tissue mass; radiographs nondiagnostic.

Both MRI and US effectively evaluate suspected soft tissue masses around the elbow. IV contrast is needed for MR evaluation of some, but not all, soft-tissue masses. For example, simple lipoma has a characteristic appearance without contrast. CT can be used as an alternative procedure, particularly for patients who have a contraindication to MRI, but is a second-tier imaging study. As with MRI, IV contrast can be a useful adjunct for diagnosis but is not necessary in all cases. Since palpable soft tissue masses are typically located outside the joint, arthrography is not appropriate for initial evaluation of a mass. Masses with hyperemia or inflammatory masses may be detected with 3-phase bone scan [28-30]. For more detailed information and references, see the ACR Appropriateness Criteria® [“Soft-Tissue Masses”](#) [31].

Variante 6: Suspect chronic epicondylitis, refractory to empirical treatment; radiographs nondiagnostic.

Epicondylitis, caused by tendon degeneration and tear of the common extensor tendon laterally (“tennis elbow”) or the common flexor tendon medially (in pitchers, golfers, and tennis players), is a common clinical diagnosis. Imaging is usually not necessary [32]. MRI or US may be useful for confirming the diagnosis in refractory cases and to exclude associated tendon and ligament tears [4,5,8,33-37]. IV contrast does not significantly aid in diagnosis of this entity [38]. Epicondylitis may be detected on 3-phase bone scan [39].

Variante 7: Suspect collateral ligament tear; radiographs nondiagnostic.

Radiographs can be useful to identify heterotopic calcification (ossification) of the ulnar collateral ligament [40]. This finding may be associated with partial or complete tears of that structure. Avulsion of the ulnar collateral ligament at the insertion site on the ulna is a source of chronic medial elbow pain in the throwing athlete. Although US has been shown to detect medial epicondylar fragmentation of the humerus in throwing athletes [27], this finding is optimally evaluated with a combination of radiographs and MRI [41]. MR arthrography has been advocated to distinguish complete tears from partial tears of the ulnar collateral ligament [7,42]. At 3T, MR arthrography is more accurate than conventional MRI in detection of ligament tears [43]. With the use of appropriate pulse sequences, MRI is an effective tool in the preoperative diagnosis of posterolateral rotatory instability. This includes assessment of the ulnar band of the lateral collateral ligament [9]. Stress views may be useful in assessing medial or lateral stability.

Variante 8: Suspect biceps tendon tear; radiographs nondiagnostic.

US has been shown to be helpful for diagnosing abnormalities of the distal biceps and triceps tendons, flexor and extensor tendons, ligaments, and nerves, providing an alternative to MRI although MRI is typically used in this clinical scenario [33-35,44-53]. Neither IV nor intra-articular contrast assists in MR diagnosis. Evaluation of the biceps tendon with CT is limited and not usually appropriate.

Variant 9: Suspect nerve abnormality; radiographs nondiagnostic.

The ulnar nerve is particularly vulnerable to trauma from a direct blow in the region of the superficially located restricted space of the cubital tunnel. Anatomic variations of the cubital tunnel retinaculum can contribute to ulnar neuropathy [54]. Axial T1-weighted MR images have been shown to depict the size and shape of the nerve, and axial T2-weighted or short tau inversion recovery images may show an increased signal in the presence of neuritis [1,44,55]. Both are more sensitive than conventional nerve conduction studies [56]. US may also show ulnar nerve enlargement and increased vascularity [47] and, when added to electrodiagnostic tests, increases sensitivity for the diagnosis of ulnar neuropathy at the elbow from 78% to 98% [57]. A snapping of the medial head of the triceps can cause recurrent dislocation of the ulnar nerve. This diagnosis can be confirmed with MRI or CT using axial images with the elbow in flexion and extension [55,58,59]. US is ideal for dynamic assessment of ulnar nerve subluxation and dislocation, as well as for confirmation of snapping triceps syndrome [60,61]. Radial nerve, median nerve, and other entrapment syndromes can also be evaluated with MRI [5,32,55,62].

Variant 10: Elbow stiffness; suspect heterotopic ossification/osteophytosis by radiography. Next test.

CT is superior to radiography in the preoperative assessment of osteophytosis or heterotopic ossification in the patient with symptomatic stiff elbow [17,63]. The inflammatory component of heterotopic ossification can be detected on the early phases of a 3-phase bone scan, and the delayed images will reveal increased uptake due to the bone formation [29,64]. Bone scintigraphy has been used in early detection of heterotopic ossification and in assessing its metabolic activity to appropriately time surgical removal [64]. MRI can also be used for this diagnosis, but osseous detail is inferior to CT. IV contrast does not contribute to this diagnosis. The use of US is limited because of shadowing from bone formation.

Variant 11: Suspect inflammatory arthritis or bursitis; radiographs obtained.

Chronic elbow pain can also be caused by a number of joint-related processes, such as inflammatory arthritis and synovial proliferative disorders. Evaluation begins with radiography to assess for joint distention and erosions. MRI can also show erosions and is effective in characterizing synovitis (a low signal suggests hemosiderin) and the extent and activity of disease [65]. In the setting of rheumatoid arthritis, US can also be used to detect joint effusion, synovitis, and erosions [66]. Bicipitoradial and interosseous bursitis around the distal biceps tendon is a source of elbow pain that can be assessed with MRI or US [52,67]. MRI also demonstrates the effects of the bursa on adjacent structures, including the posterior interosseous and median nerves [67]. Inflammatory arthritis or bursitis can be detected by the early phases of a 3-phase bone scan, as well as on the delayed images, by increased uptake [28-30].

For imaging of osseous tumors see the ACR Appropriateness Criteria® “[Primary Bone Tumors](#)” [68] and ACR Appropriateness Criteria® “[Metastatic Bone Disease](#)” [69].

Summary of Recommendations

- Initial evaluation of chronic elbow pain should begin with radiography.
- Chondral and osteochondral abnormalities can be further evaluated with MRI or CT. The addition of arthrography is helpful, especially for detecting intra-articular bodies.
- Radiographically occult bone abnormalities can be detected with MRI, CT, or bone scintigraphy.
- Soft-tissue abnormalities (tendons, ligaments, nerves, joint recesses, and masses) are well demonstrated with MRI or US.
- Dynamic assessment with US is effective for diagnosing nerve or muscle subluxation.

Summary of Evidence

Of the 69 references cited in the *ACR Appropriateness Criteria® Chronic Elbow Pain* document, all of them are categorized as diagnostic references including 1 well designed study, 8 good quality studies, and 14 quality studies that may have design limitations. There are 46 references that may not be useful as primary evidence.

The 69 references cited in the *ACR Appropriateness Criteria® Chronic Elbow Pain* document were published from 1982-2015.

While there are references that report on studies with design limitations, 9 well designed or good quality studies provide good evidence.

Relative Radiation Level Information

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Patients in the pediatric age group are at inherently higher risk from exposure, both because of organ sensitivity and longer life expectancy (relevant to the long latency that appears to accompany radiation exposure). For these reasons, the RRL dose estimate ranges for pediatric examinations are lower as compared to those specified for adults (see Table below). Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® [Radiation Dose Assessment Introduction](#) document.

Relative Radiation Level Designations		
Relative Radiation Level*	Adult Effective Dose Estimate Range	Pediatric Effective Dose Estimate Range
○	0 mSv	0 mSv
☼	<0.1 mSv	<0.03 mSv
☼☼	0.1-1 mSv	0.03-0.3 mSv
☼☼☼	1-10 mSv	0.3-3 mSv
☼☼☼☼	10-30 mSv	3-10 mSv
☼☼☼☼☼	30-100 mSv	10-30 mSv

*RRL assignments for some of the examinations cannot be made, because the actual patient doses in these procedures vary as a function of a number of factors (eg, region of the body exposed to ionizing radiation, the imaging guidance that is used). The RRLs for these examinations are designated as “Varies”.

Supporting Documents

For additional information on the Appropriateness Criteria methodology and other supporting documents go to www.acr.org/ac.

References

1. Keen NN, Chin CT, Engstrom JW, Saloner D, Steinbach LS. Diagnosing ulnar neuropathy at the elbow using magnetic resonance neurography. *Skeletal Radiol*. 2012;41(4):401-407.
2. Bredella MA, Tirman PF, Fritz RC, Feller JF, Wischer TK, Genant HK. MR imaging findings of lateral ulnar collateral ligament abnormalities in patients with lateral epicondylitis. *AJR Am J Roentgenol*. 1999;173(5):1379-1382.
3. Chung CB, Chew FS, Steinbach L. MR imaging of tendon abnormalities of the elbow. *Magn Reson Imaging Clin N Am*. 2004;12(2):233-245, vi.
4. Coel M, Yamada CY, Ko J. MR imaging of patients with lateral epicondylitis of the elbow (tennis elbow): importance of increased signal of the anconeus muscle. *AJR Am J Roentgenol*. 1993;161(5):1019-1021.
5. Fritz RC, Steinbach LS. Magnetic resonance imaging of the musculoskeletal system: Part 3. The elbow. *Clin Orthop Relat Res*. 1996(324):321-339.
6. Kijowski R, De Smet AA. Magnetic resonance imaging findings in patients with medial epicondylitis. *Skeletal Radiol*. 2005;34(4):196-202.
7. Kijowski R, Tuite M, Sanford M. Magnetic resonance imaging of the elbow. Part II: Abnormalities of the ligaments, tendons, and nerves. *Skeletal Radiol*. 2005;34(1):1-18.
8. Potter HG, Hannafin JA, Morwessel RM, DiCarlo EF, O'Brien SJ, Altchek DW. Lateral epicondylitis: correlation of MR imaging, surgical, and histopathologic findings. *Radiology*. 1995;196(1):43-46.
9. Potter HG, Weiland AJ, Schatz JA, Paletta GA, Hotchkiss RN. Posterolateral rotatory instability of the elbow: usefulness of MR imaging in diagnosis. *Radiology*. 1997;204(1):185-189.
10. Quinn SF, Haberman JJ, Fitzgerald SW, Traugher PD, Belkin RI, Murray WT. Evaluation of loose bodies in the elbow with MR imaging. *J Magn Reson Imaging*. 1994;4(2):169-172.

11. Schwartz ML, al-Zahrani S, Morwessel RM, Andrews JR. Ulnar collateral ligament injury in the throwing athlete: evaluation with saline-enhanced MR arthrography. *Radiology*. 1995;197(1):297-299.
12. Sonin AH, Tutton SM, Fitzgerald SW, Peduto AJ. MR imaging of the adult elbow. *Radiographics*. 1996;16(6):1323-1336.
13. Del Grande F, Aro M, Farahani SJ, Wilckens J, Cosgarea A, Carrino JA. Three-Tesla MR imaging of the elbow in non-symptomatic professional baseball pitchers. *Skeletal Radiol*. 2015;44(1):115-123.
14. Taljanovic MS, Hunter TB, Fitzpatrick KA, Krupinski EA, Pope TL. Musculoskeletal magnetic resonance imaging: importance of radiography. *Skeletal Radiol*. 2003;32(7):403-411.
15. O'Driscoll SW. Stress radiographs are important in diagnosing valgus instability of the elbow. *J Bone Joint Surg Am*. 2002;84-A(4):686; author reply 686-687.
16. Lee GA, Katz SD, Lazarus MD. Elbow valgus stress radiography in an uninjured population. *Am J Sports Med*. 1998;26(3):425-427.
17. Zubler V, Saupe N, Jost B, Pfirrmann CW, Hodler J, Zanetti M. Elbow stiffness: effectiveness of conventional radiography and CT to explain osseous causes. *AJR Am J Roentgenol*. 2010;194(6):W515-520.
18. Grainger AJ, Elliott JM, Campbell RS, Tirman PF, Steinbach LS, Genant HK. Direct MR arthrography: a review of current use. *Clin Radiol*. 2000;55(3):163-176.
19. Steinbach LS, Palmer WE, Schweitzer ME. Special focus session. MR arthrography. *Radiographics*. 2002;22(5):1223-1246.
20. Theodoropoulos JS, Dwyer T, Wolin PM. Correlation of preoperative MRI and MRA with arthroscopically proven articular cartilage lesions of the elbow. *Clin J Sport Med*. 2012;22(5):403-407.
21. Anderson MW. Imaging of upper extremity stress fractures in the athlete. *Clin Sports Med*. 2006;25(3):489-504, vii.
22. Ammann W, Matheson GO. Radionuclide Bone Imaging in the Detection of Stress Fractures. *Clinical Journal of Sport Medicine*. 1991;1(2):115-122.
23. Querellou S, Moineau G, Le Duc-Pennec A, et al. Detection of occult wrist fractures by quantitative radiosintigraphy: a prospective study on selected patients. *Nucl Med Commun*. 2009;30(11):862-867.
24. Haapamaki VV, Kiuru MJ, Koskinen SK. Multidetector computed tomography diagnosis of adult elbow fractures. *Acta Radiol*. 2004;45(1):65-70.
25. Zuazo I, Bonnefoy O, Tauzin C, et al. Acute elbow trauma in children: role of ultrasonography. *Pediatr Radiol*. 2008;38(9):982-988.
26. Ouellette H, Kassarijian A, Tetreault P, Palmer W. Imaging of the overhead throwing athlete. *Semin Musculoskelet Radiol*. 2005;9(4):316-333.
27. Harada M, Takahara M, Sasaki J, Mura N, Ito T, Ogino T. Using sonography for the early detection of elbow injuries among young baseball players. *AJR Am J Roentgenol*. 2006;187(6):1436-1441.
28. Alazraki NP. Radionuclide imaging in the evaluation of infections and inflammatory disease. *Radiol Clin North Am*. 1993;31(4):783-794.
29. Freed JH, Hahn H, Menter R, Dillon T. The use of the three-phase bone scan in the early diagnosis of heterotopic ossification (HO) and in the evaluation of Didronel therapy. *Paraplegia*. 1982;20(4):208-216.
30. Palestro CJ. Radionuclide imaging of osteomyelitis. *Semin Nucl Med*. 2015;45(1):32-46.
31. American College of Radiology. ACR Appropriateness Criteria®: Soft-Tissue Masses. Available at: <https://acsearch.acr.org/docs/69434/Narrative/>. Accessed September 30, 2015.
32. Beltran J, Rosenberg ZS. Diagnosis of compressive and entrapment neuropathies of the upper extremity: value of MR imaging. *AJR Am J Roentgenol*. 1994;163(3):525-531.
33. Obradov M, Anderson PG. Ultra sonographic findings for chronic lateral epicondylitis. *JBR-BTR*. 2012;95(2):66-70.
34. Poltawski L, Ali S, Jayaram V, Watson T. Reliability of sonographic assessment of tendinopathy in tennis elbow. *Skeletal Radiol*. 2012;41(1):83-89.
35. Martinoli C, Bianchi S, Giovagnorio F, Pugliese F. Ultrasound of the elbow. *Skeletal Radiol*. 2001;30(11):605-614.
36. Park GY, Lee SM, Lee MY. Diagnostic value of ultrasonography for clinical medial epicondylitis. *Arch Phys Med Rehabil*. 2008;89(4):738-742.
37. van Kollenburg JA, Brouwer KM, Jupiter JB, Ring D. Magnetic resonance imaging signal abnormalities in enthesopathy of the extensor carpi radialis longus origin. *J Hand Surg Am*. 2009;34(6):1094-1098.
38. Herber S, Kalden P, Kreitner KF, Riedel C, Rompe JD, Thelen M. [MRI in chronic epicondylitis humeri radialis using 1.0 T equipment--contrast medium administration necessary?]. *Rofö*. 2001;173(5):454-459.

39. Pienimäki TT, Takalo RJ, Ahonen AK, Karppinen JI. Three-phase bone scintigraphy in chronic epicondylitis. *Arch Phys Med Rehabil.* 2008;89(11):2180-2184.
40. Mulligan SA, Schwartz ML, Broussard MF, Andrews JR. Heterotopic calcification and tears of the ulnar collateral ligament: radiographic and MR imaging findings. *AJR Am J Roentgenol.* 2000;175(4):1099-1102.
41. Glajchen N, Schwartz ML, Andrews JR, Gladstone J. Avulsion fracture of the sublime tubercle of the ulna: a newly recognized injury in the throwing athlete. *AJR Am J Roentgenol.* 1998;170(3):627-628.
42. Steinbach LS, Schwartz M. Elbow arthrography. *Radiol Clin North Am.* 1998;36(4):635-649.
43. Magee T. Accuracy of 3-T MR arthrography versus conventional 3-T MRI of elbow tendons and ligaments compared with surgery. *AJR Am J Roentgenol.* 2015;204(1):W70-75.
44. Baumer P, Dombert T, Staub F, et al. Ulnar neuropathy at the elbow: MR neurography--nerve T2 signal increase and caliber. *Radiology.* 2011;260(1):199-206.
45. Downey R, Jacobson JA, Fessell DP, Tran N, Morag Y, Kim SM. Sonography of partial-thickness tears of the distal triceps brachii tendon. *J Ultrasound Med.* 2011;30(10):1351-1356.
46. Festa A, Mulieri PJ, Newman JS, Spitz DJ, Leslie BM. Effectiveness of magnetic resonance imaging in detecting partial and complete distal biceps tendon rupture. *J Hand Surg Am.* 2010;35(1):77-83.
47. Frijlink DW, Brekelmans GJ, Visser LH. Increased nerve vascularization detected by color Doppler sonography in patients with ulnar neuropathy at the elbow indicates axonal damage. *Muscle Nerve.* 2013;47(2):188-193.
48. Tagliafico A, Gandolfo N, Michaud J, Perez MM, Palmieri F, Martinoli C. Ultrasound demonstration of distal triceps tendon tears. *Eur J Radiol.* 2012;81(6):1207-1210.
49. Tagliafico A, Michaud J, Perez MM, Martinoli C. Ultrasound of distal brachialis tendon attachment: normal and abnormal findings. *Br J Radiol.* 2013;86(1025):20130004.
50. Miller TT, Adler RS, Friedman L. Sonography of injury of the ulnar collateral ligament of the elbow-initial experience. *Skeletal Radiol.* 2004;33(7):386-391.
51. Miller TT, Shapiro MA, Schultz E, Kalish PE. Comparison of sonography and MRI for diagnosing epicondylitis. *J Clin Ultrasound.* 2002;30(4):193-202.
52. Sofka CM, Adler RS. Sonography of cubital bursitis. *AJR Am J Roentgenol.* 2004;183(1):51-53.
53. Lobo Lda G, Fessell DP, Miller BS, et al. The role of sonography in differentiating full versus partial distal biceps tendon tears: correlation with surgical findings. *AJR Am J Roentgenol.* 2013;200(1):158-162.
54. Sofka CM, Potter HG. Imaging of elbow injuries in the child and adult athlete. *Radiol Clin North Am.* 2002;40(2):251-265.
55. Bordalo-Rodrigues M, Rosenberg ZS. MR imaging of entrapment neuropathies at the elbow. *Magn Reson Imaging Clin N Am.* 2004;12(2):247-263, vi.
56. Vucic S, Cordato DJ, Yiannikas C, Schwartz RS, Shnier RC. Utility of magnetic resonance imaging in diagnosing ulnar neuropathy at the elbow. *Clin Neurophysiol.* 2006;117(3):590-595.
57. Beekman R, Van Der Plas JP, Uitdehaag BM, Schellens RL, Visser LH. Clinical, electrodiagnostic, and sonographic studies in ulnar neuropathy at the elbow. *Muscle Nerve.* 2004;30(2):202-208.
58. Spinner RJ, Goldner RD, Fada RA, Sotereanos DG. Snapping of the triceps tendon over the lateral epicondyle. *J Hand Surg Am.* 1999;24(2):381-385.
59. Spinner RJ, Hayden FR, Jr., Hipps CT, Goldner RD. Imaging the snapping triceps. *AJR Am J Roentgenol.* 1996;167(6):1550-1551.
60. Park GY, Kim JM, Lee SM. The ultrasonographic and electrodiagnostic findings of ulnar neuropathy at the elbow. *Arch Phys Med Rehabil.* 2004;85(6):1000-1005.
61. Jacobson JA, Jebson PJ, Jeffers AW, Fessell DP, Hayes CW. Ulnar nerve dislocation and snapping triceps syndrome: diagnosis with dynamic sonography--report of three cases. *Radiology.* 2001;220(3):601-605.
62. Bodner G, Harpf C, Meirer R, Gardetto A, Kovacs P, Gruber H. Ultrasonographic appearance of supinator syndrome. *J Ultrasound Med.* 2002;21(11):1289-1293.
63. Lindenhovius AL, Jupiter JB. The posttraumatic stiff elbow: a review of the literature. *J Hand Surg Am.* 2007;32(10):1605-1623.
64. Shehab D, Elgazzar AH, Collier BD. Heterotopic ossification. *J Nucl Med.* 2002;43(3):346-353.
65. Jbara M, Patnana M, Kazmi F, Beltran J. MR imaging: Arthropathies and infectious conditions of the elbow, wrist, and hand. *Radiol Clin North Am.* 2006;44(4):625-642, ix.
66. Lerch K, Borisch N, Paetzl C, Grifka J, Hartung W. Proposal for a sonographic classification of target joints in rheumatoid arthritis. *Rheumatol Int.* 2005;25(3):215-219.

67. Skaf AY, Boutin RD, Dantas RW, et al. Bicipitoradial bursitis: MR imaging findings in eight patients and anatomic data from contrast material opacification of bursae followed by routine radiography and MR imaging in cadavers. *Radiology*. 1999;212(1):111-116.
68. American College of Radiology. ACR Appropriateness Criteria®: Primary Bone Tumors. Available at: <https://acsearch.acr.org/docs/69421/Narrative/>. Accessed September 30, 2015.
69. American College of Radiology. ACR Appropriateness Criteria®: Metastatic Bone Disease. Available at: <https://acsearch.acr.org/docs/69431/Narrative/>. Accessed September 30, 2015.

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.