

American College of Radiology ACR Appropriateness Criteria®

Clinical Condition: Fever Without Source or Unknown Origin—Child

Variant 1: Neonate younger than 1 month of age with fever without source (FWS) and no respiratory symptoms.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	5	This procedure may be appropriate but there was disagreement among panel members on the appropriateness rating as defined by the panel's median rating.	⊕
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 2: Neonate younger than 1 month of age with FWS and respiratory symptoms.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	8		⊕
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 3: Infant or child aged 1 to 36 months with FWS with low risk for occult pneumonia (no respiratory signs or symptoms, fever <39°C, leukocytosis <20,000/mm³).

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	2		⊕
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 4: Infant or child aged 1 to 36 months with FWS with any of the following: respiratory signs or symptoms, fever ≥39°C, or white blood cell count ≥20,000/mm³.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	9		⊕
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Clinical Condition: Fever Without Source or Unknown Origin—Child

Variant 5: Child with FWS and neutropenia.

Radiologic Procedure	Rating	Comments	RRL*
CT chest with IV contrast	6	This procedure may be appropriate if patient has respiratory symptoms or has had stem cell transplant.	☼☼☼☼
X-ray chest	5	This procedure may be appropriate but there was disagreement among panel members on the appropriateness rating as defined by the panel’s median rating. This procedure may be appropriate if patient has respiratory symptoms.	☼
CT chest without IV contrast	5	This procedure may be appropriate if patient has respiratory symptoms or has had stem cell transplant.	☼☼☼☼
CT abdomen with IV contrast	5	Consider in patients who have had stem cell transplant.	☼☼☼☼
CT paranasal sinuses with IV contrast	4	Contrast and brain imaging are essential if central nervous system invasion is a concern.	☼☼☼
CT paranasal sinuses without IV contrast	4	Consider in patients who have had stem cell transplant.	☼☼☼
CT abdomen without IV contrast	2		☼☼☼☼
CT chest without and with IV contrast	1		☼☼☼☼
CT abdomen without and with IV contrast	1		☼☼☼☼☼
CT paranasal sinuses without and with IV contrast	1		☼☼☼☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Clinical Condition: Fever Without Source or Unknown Origin—Child

Variant 6: Infant or child more than 1 month of age with fever of unknown origin (FUO).

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	7		☼
US abdomen	6		○
CT chest with IV contrast	5		☼☼☼☼
CT chest without IV contrast	4		☼☼☼☼
CT abdomen with IV contrast	4		☼☼☼☼
CT paranasal sinuses without IV contrast	4		☼☼☼
FDG-PET/CT skull base to mid-thigh	4	This procedure should not be used as the initial study. Consider if extensive clinical and imaging workup is negative.	☼☼☼☼☼
CT paranasal sinuses with IV contrast	3		☼☼☼
MRI whole body without IV contrast	3		○
CT abdomen without IV contrast	2		☼☼☼☼
MRI whole body without and with IV contrast	2	This procedure should not be used as the initial study. Consider if extensive clinical and imaging workup is negative.	○
CT chest without and with IV contrast	1		☼☼☼☼
CT abdomen without and with IV contrast	1		☼☼☼☼☼
CT paranasal sinuses without and with IV contrast	1		☼☼☼☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

FEVER WITHOUT SOURCE OR UNKNOWN ORIGIN—CHILD

Expert Panel on Pediatric Imaging: Sjikr J. Westra, MD¹; Boaz K. Karmazyn, MD²; Adina L. Alazraki, MD³; Molly E. Dempsey, MD⁴; Jonathan R. Dillman, MD⁵; Matthew Garber, MD⁶; Sheila G. Moore, MD⁷; Molly E. Raske, MD⁸; Henry E. Rice, MD⁹; Cynthia K. Rigsby, MD¹⁰; Nabile Safdar, MD, MPH¹¹; Stephen F. Simoneaux, MD¹²; Peter J. Strouse, MD¹³; Andrew T. Trout, MD¹⁴; Sandra L. Wootton-Gorges, MD¹⁵; Brian D. Coley, MD.¹⁶

Summary of Literature Review

Introduction/Background

The febrile pediatric patient, especially an infant, represents a dilemma for the primary care physician. The definition of fever is generally regarded as a rectal temperature of $\geq 38^{\circ}\text{C}$ [1-3]. Oral temperatures are less reliable in infants and young children, although they are the usual method of measuring temperature in older children and adults. The cause of fever in the pediatric patient can often be determined from the history, physical examination, and laboratory tests [4-9]. Prior medical conditions, medications, foreign travel, and immunization history are all important in directing subsequent investigations [2,10-14]. However, 20% of cases will have no apparent source and thus are defined as having fever without source (FWS) [5]. FWS is therefore defined as an acute illness in which the origin of the fever is not apparent after initial careful history and examination [2,5,6,8,9,14,15]. Most FWS are caused by infections [2,6,8,9,14,16]. The approach to a febrile child is generally divided into the infant younger than 3 months and the older infant and child between 3 and 36 months of age [3,5-7]. Infants younger than 1 month deserve more aggressive evaluation because these children have more immature immune systems, are more difficult to evaluate, and do not have the protection afforded by the *Hemophilus influenzae* (*H. flu*) and *Streptococcus pneumoniae* (*S. pneumoniae*) vaccines [2,3,6,9,17].

Although FWS is mostly self-limited and of little clinical concern, the burden on clinicians is to decide which children actually have a serious bacterial infection (SBI) that requires antibiotic treatment and even hospitalization [18,19]. Febrile neonates are at higher risk; the reported incidence of SBI in all febrile neonates presenting to emergency departments varies between 6% and 28% [17,20]. In children, the usual sources/causes of SBI are urinary tract infection, pneumonia, bloodstream infection, and meningitis. With the advent of vaccines for the most common pathogenic serotypes of *H. flu* and *S. pneumoniae*, the incidence of SBI has dropped significantly [2,3,6,10,13,14]. Although it is implied by the definition of FWS that the etiology of fever is unknown, many studies and guidelines include children with respiratory symptoms [2,3,5,6,8,16,21-24]. For this reason, we were compelled to include in our guidelines for FWS children with respiratory symptoms.

Although the terms are sometimes used interchangeably, FWS is different from fever of unknown origin (FUO). Pediatric FUO refers to a fever of $\geq 38.3^{\circ}\text{C}$ with no apparent diagnosis after initial outpatient or hospital evaluation that includes a careful history and physical exam and initial laboratory assessment. There is much variability in published studies of FUO, with required duration of fever ranging from 1 to 3 weeks [9,20,25]. The majority of children with FUO have infectious causes, although inflammatory, neoplastic, and autoimmune conditions are also in the differential [9,20,24-27]. The distinction between FWS and FUO is more than just academic because the clinical and imaging approaches to these conditions can differ.

Overview of Imaging Modalities

A detailed and thorough history and physical examination is the most important component of the diagnostic evaluation of a child with FWS or FUO. Chest radiographs have a role in evaluation of occult pneumonia and should be performed in neonates with FWS and respiratory symptoms and in selected older children with high

¹Principal Author, Massachusetts General Hospital, Boston, Massachusetts. ²Co-Author and Panel Chair, Riley Hospital for Children, Indiana University, Indianapolis, Indiana. ³Children's Healthcare of Atlanta, Atlanta, Georgia. ⁴Texas Scottish Rite Hospital for Children, Dallas, Texas. ⁵C. S. Mott Children's Hospital, Ann Arbor, Michigan. ⁶Division of General and Hospital Pediatrics, Columbia, South Carolina, American Academy of Pediatrics. ⁷Children's Hospital of Wisconsin, Milwaukee, Wisconsin. ⁸Children's Hospitals and Clinics of Minnesota, Minneapolis, Minnesota. ⁹Duke University Medical Center, Durham, North Carolina, American Pediatric Surgical Association. ¹⁰Ann & Robert H. Lurie Children's Hospital of Chicago, Chicago, Illinois. ¹¹Emory University Hospital, Atlanta, Georgia. ¹²Children's Healthcare of Atlanta, Atlanta, Georgia. ¹³C. S. Mott Children's Hospital, Ann Arbor, Michigan. ¹⁴Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio. ¹⁵University of California Davis Medical Center, Sacramento, California. ¹⁶Specialty Chair, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio.

The American College of Radiology seeks and encourages collaboration with other organizations on the development of the ACR Appropriateness Criteria through society representation on expert panels. Participation by representatives from collaborating societies on the expert panel does not necessarily imply individual or society endorsement of the final document.

Reprint requests to: publications@acr.org

fever, leukocytosis, and respiratory signs and symptoms [28]. However, chest radiographs are limited in evaluation of pneumonia in young children due to poor differentiation between viral and bacterial pneumonia and considerable intraobserver and interobserver variability in interpretation [29-34]. Computed tomography (CT) of the paranasal sinus, chest, and abdomen are important for evaluation of fungal infection in neutropenic patients, especially after bone marrow transplantation, and in patients who do not respond to broad-spectrum antibiotics [35,36]. There are small series on the use of fluorine-18-2-fluoro-2-deoxy-D-glucose positron emission tomography (FDG-PET)/CT in evaluation of FUO that show its potential to detect occult infection, inflammatory processes, and malignancy. It can be used in selected children with FUO with negative extensive workup [37-39].

Discussion of Imaging Modalities by Variant

Variants 1 and 2: Neonate younger than 1 month of age with FWS.

In febrile neonates younger than 28 days, history and physical examination alone may not be able to completely exclude SBI, even in children who appear clinically well or mildly ill [40]. Therefore, a full sepsis workup is frequently performed. This includes complete blood count, blood culture, urinalysis and urine culture, lumbar puncture with evaluation of cerebrospinal fluid, and administration of antibiotics in the emergency department, followed by hospitalization pending results of cultures [2,3,8,9,22,28].

A chest radiograph is indicated in neonates with FWS and respiratory symptoms [3,21]. In addition, a chest radiograph in a septic-appearing neonate with FWS may disclose an occult pneumonia [6,8,16,28]. Some investigators advocate routine chest radiographs in all neonates with FWS because these infants are relatively immunocompromised compared with older infants and children, and the consequences of a missed SBI or occult infection are felt to be greater [6]. A chest radiograph can help exclude congenital or acquired cardiac disease in a child who is febrile and ill. However, the benefit of routine use of chest radiography in neonates without respiratory symptoms has not been established [3,21].

Variant 3: Infant or child aged 1 to 36 months with FWS with low risk for occult pneumonia (no respiratory signs or symptoms, fever <39°C, leukocytosis <20,000/mm³).

In the absence of a “toxic” appearance, respiratory distress, poor peripheral perfusion, high fever, and leukocytosis, the risk for SBI is low in children with FWS and there is no indication for routine chest radiography. Patterson et al [41] retrospectively studied 105 infants who had fever. Of the 37 patients who had no respiratory symptoms or signs, only 1 had a chest radiograph showing a focal parenchymal airspace disease. Hyperinflation and peribronchial thickening were not classified as abnormal. In a prospective study, the same authors included 121 infants who were free of signs or symptoms of lower respiratory tract infection but who had fever [41]. None had chest radiographs that showed an abnormality. Even in younger children at the age of 1 to 3 months that are of increased risk of SBI, there is no role of routine chest radiography. Heulitt et al [42] showed that only 6% of infants with fever without respiratory manifestations developed pneumonia and that all of those infants did well. A meta-analysis of 361 febrile infants younger than 3 months without clinical evidence of pulmonary disease on history or physical examination showed that none of these children had pneumonia [43]. Baraff et al [44] reported a 3.3% incidence of positive chest radiographs based on collected reviews of infants and children from birth to 36 months of age with fever and no respiratory symptoms or signs. McCarthy [45], summarizing a number of clinical series dealing with acute episodes of fever in infants, also believes that chest radiographs should be obtained only when there are clinical symptoms or signs of pneumonia. A later study by Baraff reports that occult pneumonia is seen in only 3% of infants without respiratory findings on physical examination. Given that the risk of SBI in febrile infants and children has dropped in the era of pneumococcal vaccination and that most FWS cases will be related to urinary tract or viral infections, some authors recommend obtaining urinalysis first and considering chest imaging only if this is negative [2,9,46].

Bramson et al [43] combined their data with those of 2 prior studies [42,47] and subjected these to a statistical meta-analysis. The larger number of patients in the combined study allowed more valid conclusions concerning the accepted practice of performing chest radiographs in febrile infants as part of the sepsis workup. These 3 series had 671 infants. In 361 infants with no clinical evidence of pulmonary disease on history and physical examination, all had normal chest radiographs. A finding of only hyperinflation on a chest radiograph was interpreted as normal because it was felt that the infants would likely have a viral illness or reactive airway disease and would not usually be receiving antibiotics, unlike older children and adults [48]. Bramson et al [43] indicated that a chest radiograph in a patient with no pulmonary symptoms or signs would be positive <1.2% of the time. In the current era of *S. pneumoniae* and *H. flu* vaccine use, this rate might fall even further. Murphy et al [49] found an incidence of radiographic pneumonia in 5.3% of 2,128 children under 10 years of age with no lower

respiratory symptoms (other than cough) and concluded that there was a low utility in obtaining chest radiographs in febrile children without cough. A longer duration of cough, fever, and leukocytosis increased the likelihood of radiographic pneumonia in these children.

Variant 4: Infant or child aged 1 to 36 months with FWS with any of the following: respiratory signs or symptoms, fever $\geq 39^{\circ}\text{C}$, or white blood cell count $\geq 20,000/\text{mm}^3$.

Patients with high fever, signs of respiratory distress, or white blood cell (WBC) count $>20,000/\text{mm}^3$ are at increased risk of pneumonia; therefore, chest radiographs are indicated [2,5,6,8,16,22-24]. The presence of rales is the single best clinical indicator of pneumonia in infants and children. Tachypnea, intercostal retractions, and nasal flaring are also predictive of pneumonia in the pediatric population [50-53]. Other clinical factors that can be predictive of pneumonia in children of all ages, such as degree of fever, WBC count, and pulse oximetry, have been studied [16,52,54-56].

In a meta-analysis study [43], it was found that 33.2% (85/256) of the infants younger than 3 months with at least 1 clinical finding of pulmonary disease (ie, tachypnea >50 breaths/min, cyanosis, O_2 saturation $<95\%$, rales, rhonchi, retractions, wheezing, coryza, grunting, stridor, nasal flaring, or cough) had positive chest radiographs. In a study by Mahabee-Gittens et al [52] of 510 children from 2 to 59 months of age presenting with symptoms of lower respiratory infection who had chest radiographs, 8.6% showed pneumonia. Clinical variables found to correlate with positive radiographic findings included age >12 months, respiratory rate >50 breaths per minute, oxygen saturation $\leq 96\%$, and nasal flaring in children younger than 12 months. Combinations of these clinical variables produced likelihood ratios of radiographic pneumonia from 3.6 to 11.0. In a similar study by Lynch et al [53] of 570 children over 1 year of age, the presence of decreased breath sounds, crackles, and tachypnea in various combinations had a high sensitivity but a poor specificity to predict pneumonia, defined as “focal infiltrates on chest radiographs.” Bachur et al [56] found that 26% of children with fever $\geq 39^{\circ}\text{C}$ and a WBC count $\geq 20,000/\text{mm}^3$ had pneumonia on chest radiographs. The use of polyvalent *S. pneumonia* vaccine has been shown to reduce pneumonia with radiographic consolidation by 73% [57]. This led Baraff [5] to suggest that a chest radiograph should be obtained in patients with high fever and elevated WBC count who have not received the pneumococcal vaccine, regardless of respiratory findings. Rutman et al [14] reported that since the institution of pneumococcal vaccination, the incidence of radiographically evident pneumonia has dropped by 20% to 39%. In their study of 355 children younger than 5 years with fever $>39^{\circ}\text{C}$, WBC count $>20,000/\text{mm}^3$, and respiratory symptoms, pneumonia was present in 18%, making chest radiography a reasonable study under those circumstances. Brook [16] also recommends obtaining a chest radiograph in all patients younger than 36 months with an oxygen saturation $<95\%$, although there is no supporting evidence given nor are there data as to the diagnostic yield of such radiographs.

The American College of Emergency Physicians states that a chest radiograph should be considered in patients older than 3 months with fever $\geq 39^{\circ}\text{C}$ and a WBC count $\geq 20,000/\text{mm}^3$ [58]. Similar recommendations have been made by the British Thoracic Society for children younger than 5 years [59].

Variant 5: Child with FWS and neutropenia.

Fever is a cause of great concern in a child with cancer or immunodeficiency who is neutropenic. In neutropenic patients, a significant fever is usually defined as a single oral temperature of $\geq 38.3^{\circ}\text{C}$ or 2 measurements of $\geq 38.0^{\circ}\text{C}$ at least 1 hour apart [60]. Neutropenia is an absolute neutrophil count of $<500/\text{mm}^3$, or $<1000/\text{mm}^3$ with the expectation of rapid decrease [60]. Such children are more susceptible to the common infections facing all children; gram-positive organisms are responsible for 70% of SBI in these patients, but gram-negative organisms are responsible for most SBI-related fatalities [60]. These patients are also at risk for viral and other atypical infections, and invasive fungal infections are a particular concern for high-risk patients with persistent febrile neutropenia [60]. Because of the heightened clinical concern, a chest radiograph is usually obtained in addition to other assessments, including cultures of the blood and urine.

The practice of routinely obtaining a chest radiograph has been challenged. Korones et al [61] evaluated 54 children with cancer who were hospitalized for hundreds of episodes of fever and neutropenia. They found an incidence of radiographic pneumonia of only 3% to 6%. The children without respiratory findings had no evidence of pneumonia on chest radiographs, and children who did not have chest radiographs showed no significant outcome differences from those who did. Philips et al [62] confirmed in a meta-analysis the low use of routine chest radiographs in this setting but stated that in those with a predisposition to pneumonia and those not responding to a short empiric course of antibiotics, chest radiographs should be performed despite the absence of clinical signs of a lower respiratory tract infection.

Children with neutropenia and FWS often undergo advanced imaging, but there is little evidence-based data about which studies are most efficacious. In their 2002 guidelines (not pediatric specific), the Infectious Diseases Society of America noted that one-half of febrile neutropenic patients with normal chest radiographs will have evidence of pneumonia on chest CT [63]. Archibald et al [36] evaluated the performance of CT in 83 neutropenic pediatric cancer patients who had 109 instances of fever lasting 4 days or more. Rates of positive CT findings varied by body region: head and neck, 8%; paranasal sinus, 41%; chest, 49%; and abdomen, 19%. Findings on paranasal sinus and chest CT led to changes in therapy in 24% and 30% of cases, respectively. However, they added that CT was rarely abnormal in the absence of localizing signs or symptoms and that in the absence of symptoms, CT findings rarely lead to therapeutic changes. In a more recent study, Agrawal et al [35] demonstrated a similar distribution of positive findings among body regions but found that only 2 of the initial positive CT scans led to a change in management (6.5% of positive scans, 0.8% of all initial scans). They therefore recommend limiting initial empiric CT imaging to the chest only in patients without localizing signs or symptoms. Regarding the use of FDG-PET/CT, Blokhuis et al [39] found a 78% sensitivity and 67% specificity in 12 of such children.

An important subset of neutropenic children is those who have undergone hematopoietic stem cell transplant. Cox et al [64] studied 81 hematopoietic stem cell transplant patients who had a chest radiograph performed as part of the evaluation of initial fever during transplant. None of the chest radiographs provided sufficient information for further management. In 2 of 14 episodes in patients with normal chest radiographs and in 9 of 22 episodes in patients with nonspecific chest radiographs, CT scanning resulted in a change in clinical management. Findings of large lung nodules and “halo sign” are suggestive of fungal infection [65].

Variant 6: Infant or child more than 1 month of age with fever of unknown origin (FUO).

Occult infection is the usual cause of FUO in children and is less commonly due to rheumatologic, autoimmune, neoplastic, or other inflammatory conditions [9,20,24,27,66,67]. Some children never have a specific diagnosis reached [9,24]. Evaluation of FUO in children is mainly based on thorough physical examination, history, and laboratory studies such as a complete blood cell count and peripheral smear, erythrocyte sedimentation rate, C-reactive protein, aerobic blood cultures, urinalysis, urine culture, tuberculin skin test, electrolytes, blood urea nitrogen, creatinine, hepatic enzymes, and human immunodeficiency virus serology [9,20,24,27,68,69]. Chest radiographs are usually obtained to evaluate for occult pneumonia and lymphadenopathy. Although many studies describe the clinical course of such patients, few of them examine the utility of diagnostic imaging modalities in these difficult patients. In general, if a detailed review of the history, physical examination, and screening evaluation fail to suggest a diagnosis, more extensive imaging can be considered. This includes abdominal ultrasound (US) and CT studies of the chest, abdomen, and paranasal sinus [4]. Steele et al [70] evaluated 109 children with FUO with conventional radionuclide techniques. These studies were often positive but rarely led to an unsuspected diagnosis.

FDG-PET/CT is sensitive in detection of infection, inflammatory diseases, vasculitis, arthritis, and malignancies and was found helpful on evaluation of FUO in several adult series [71,72] and meta-analysis studies [73,74]. The highest yield of FDG-PET can be expected in patients with adenopathy, low hemoglobin, and increased C-reactive protein levels [75]. There are only a few small series on the use of FDG-PET/CT in children with FUO [37]. Blokhuis et al [39] found a sensitivity of 80% and a specificity of 78% in 16 of such children. In the largest series to date by Jasper et al [38], FDG-PET/CT was used in evaluation of 44 children with FUO and 33 children with unexplained signs of inflammation without fever. According to the authors, the PET findings contributed to the final diagnosis in 35 patients (45%). The study methodology is limited, with no defined selection criteria for children with FUO in whom the PET study was performed and a vague definition on the benefits of FDG-PET/CT in reaching the final diagnosis.

Whole body magnetic resonance imaging (MRI) based on inversion recovery, T1, and diffusion sequences is an evolving technique that was used in several small series in the evaluation of systemic diseases, such as multifocal osteomyelitis and tumors, but there is not yet sufficient evidence for its use for evaluation of FUO [76-78]. The same applies to PET/MRI [79].

Summary of Recommendations

- Neonates younger than 1 month with FWS are a high-risk group; however, the yield of routine chest radiography is low in the absence of respiratory symptoms.

- In a child with FWS, a chest radiograph should be obtained when there is clinical evidence of a respiratory illness and for those with fever $\geq 39^{\circ}\text{C}$ or WBC count $\geq 20,000/\text{mm}^3$.
- In children with neutropenia and FWS, especially those after bone marrow transplant with persistent fever despite the administration of antibiotics, CT of the chest should be considered even if the chest radiograph is negative. There is lower yield for CT of the abdomen and paranasal sinus.
- Imaging studies in children with FUO have a low yield.
- More studies are needed to evaluate the potential role of FDG-PET/CT and whole-body MRI in management of children with fever.

Summary of Evidence

Of the 79 references cited in the *ACR Appropriateness Criteria® Fever without Source-Child* document, 73 are categorized as diagnostic references including 1 well designed study, 3 good quality studies, and 16 quality studies that may have design limitations. Additionally, 2 references are categorized as therapeutic references including 1 well designed study. There are 54 references that may not be useful as primary evidence. There are 4 references that are meta-analysis studies.

The 79 references cited in the *ACR Appropriateness Criteria® Fever without Source-Child* document were published from 1972-2014.

While there are references that report on studies with design limitations, 5 well designed or good quality studies provide good evidence.

Relative Radiation Level Information

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Patients in the pediatric age group are at inherently higher risk from exposure, both because of organ sensitivity and longer life expectancy (relevant to the long latency that appears to accompany radiation exposure). For these reasons, the RRL dose estimate ranges for pediatric examinations are lower as compared to those specified for adults (see Table below). Additional information regarding radiation dose assessment for imaging examinations can be found in the *ACR Appropriateness Criteria® Radiation Dose Assessment Introduction* document.

Relative Radiation Level Designations		
Relative Radiation Level*	Adult Effective Dose Estimate Range	Pediatric Effective Dose Estimate Range
○	0 mSv	0 mSv
⊕	<0.1 mSv	<0.03 mSv
⊕⊕	0.1-1 mSv	0.03-0.3 mSv
⊕⊕⊕	1-10 mSv	0.3-3 mSv
⊕⊕⊕⊕	10-30 mSv	3-10 mSv
⊕⊕⊕⊕⊕	30-100 mSv	10-30 mSv

*RRL assignments for some of the examinations cannot be made, because the actual patient doses in these procedures vary as a function of a number of factors (eg, region of the body exposed to ionizing radiation, the imaging guidance that is used). The RRLs for these examinations are designated as “Varies”.

Supporting Documents

For additional information on the Appropriateness Criteria methodology and other supporting documents go to www.acr.org/ac.

References

1. Chinnock R, Butto J, Fernando N. Hot tots: current approach to the young febrile infant. *Compr Ther*. 1995;21(3):109-114.
2. Baraff LJ. Management of infants and young children with fever without source. *Pediatr Ann*. 2008;37(10):673-679.
3. Ishimine P. The evolving approach to the young child who has fever and no obvious source. *Emerg Med Clin North Am*. 2007;25(4):1087-1115, vii.
4. Arnow PM, Flaherty JP. Fever of unknown origin. *Lancet*. 1997;350(9077):575-580.
5. Baraff LJ. Management of fever without source in infants and children. *Ann Emerg Med*. 2000;36(6):602-614.
6. Ishimine P. Fever without source in children 0 to 36 months of age. *Pediatr Clin North Am*. 2006;53(2):167-194.
7. Kourtis AP, Sullivan DT, Sathian U. Practice guidelines for the management of febrile infants less than 90 days of age at the ambulatory network of a large pediatric health care system in the United States: summary of new evidence. *Clin Pediatr (Phila)*. 2004;43(1):11-16.
8. Massin MM, Montesanti J, Lepage P. Management of fever without source in young children presenting to an emergency room. *Acta Paediatr*. 2006;95(11):1446-1450.
9. Tolan RW, Jr. Fever of unknown origin: a diagnostic approach to this vexing problem. *Clin Pediatr (Phila)*. 2010;49(3):207-213.
10. Gabriel ME, Aiuto L, Kohn N, Barone SR. Management of febrile children in the conjugate pneumococcal vaccine era. *Clin Pediatr (Phila)*. 2004;43(1):75-82.
11. Lohr JA, Hendley JO. Prolonged fever of unknown origin: a record of experiences with 54 childhood patients. *Clin Pediatr (Phila)*. 1977;16(9):768-773.
12. McClung HJ. Prolonged fever of unknown origin in children. *Am J Dis Child*. 1972;124(4):544-550.
13. Mintegi S, Benito J, Sanchez J, Azkunaga B, Iturralde I, Garcia S. Predictors of occult bacteremia in young febrile children in the era of heptavalent pneumococcal conjugated vaccine. *Eur J Emerg Med*. 2009;16(4):199-205.
14. Rutman MS, Bachur R, Harper MB. Radiographic pneumonia in young, highly febrile children with leukocytosis before and after universal conjugate pneumococcal vaccination. *Pediatr Emerg Care*. 2009;25(1):1-7.
15. Gartner JC, Jr. Fever of unknown origin. *Adv Pediatr Infect Dis*. 1992;7:1-24.
16. Brook I. Unexplained fever in young children: how to manage severe bacterial infection. *Bmj*. 2003;327(7423):1094-1097.
17. Aronson PL. Evaluation of the febrile young infant: an update. *Pediatr Emerg Med Pract*. 2013;10(2):1-17.
18. Lacour AG, Zamora SA, Gervaix A. A score identifying serious bacterial infections in children with fever without source. *Pediatr Infect Dis J*. 2008;27(7):654-656.
19. Semeraro M, Thomee C, Rolland E, et al. A predictor of unfavourable outcome in neutropenic paediatric patients presenting with fever of unknown origin. *Pediatr Blood Cancer*. 2010;54(2):284-290.
20. Cogulu O, Koturoglu G, Kurugol Z, Ozkinay F, Vardar F, Ozkinay C. Evaluation of 80 children with prolonged fever. *Pediatr Int*. 2003;45(5):564-569.
21. Arora R, Mahajan P. Evaluation of child with fever without source: review of literature and update. *Pediatr Clin North Am*. 2013;60(5):1049-1062.
22. Goldman RD, Scolnik D, Chauvin-Kimoff L, et al. Practice variations in the treatment of febrile infants among pediatric emergency physicians. *Pediatrics*. 2009;124(2):439-445.
23. Machado BM, Cardoso DM, de Paulis M, Escobar AM, Gilio AE. Fever without source: evaluation of a guideline. *J Pediatr (Rio J)*. 2009;85(5):426-432.
24. Pasic S, Minic A, Djuric P, et al. Fever of unknown origin in 185 paediatric patients: a single-centre experience. *Acta Paediatr*. 2006;95(4):463-466.
25. McCarthy P. Fever without apparent source on clinical examination. *Curr Opin Pediatr*. 2005;17(1):93-110.
26. Hofer M, Mahlaoui N, Prieur AM. A child with a systemic febrile illness - differential diagnosis and management. *Best Pract Res Clin Rheumatol*. 2006;20(4):627-640.
27. Palazzi DL. Fever of unknown origin in children: Evaluation. 2014; Available at: <http://www.uptodate.com/contents/fever-of-unknown-origin-in-children-evaluation>. Accessed September 30, 2015.
28. Moher D, Hui C, Neto G, et al. Diagnosis and Management of Febrile Infants (0–3 Months). Evidence Report/Technology Assessment No. 205. 2012; 1-18. Available at:

<http://www.effectivehealthcare.ahrq.gov/search-for-guides-reviews-and-reports/?pageaction=displayproduct&productid=1015>. Accessed September 30, 2015.

29. Albaum MN, Hill LC, Murphy M, et al. Interobserver reliability of the chest radiograph in community-acquired pneumonia. PORT Investigators. *Chest*. 1996;110(2):343-350.
30. Bettenay FA, de Campo JF, McCrossin DB. Differentiating bacterial from viral pneumonias in children. *Pediatr Radiol*. 1988;18(6):453-454.
31. Johnson J, Kline JA. Intraobserver and interobserver agreement of the interpretation of pediatric chest radiographs. *Emerg Radiol*. 2010;17(4):285-290.
32. Korppi M, Don M, Valent F, Canciani M. The value of clinical features in differentiating between viral, pneumococcal and atypical bacterial pneumonia in children. *Acta Paediatr*. 2008;97(7):943-947.
33. Spottswood SE, Liaw K, Hernanz-Schulman M, et al. The clinical impact of the radiology report in wheezing and nonwheezing febrile children: a survey of clinicians. *Pediatr Radiol*. 2009;39(4):348-353.
34. Williams GJ, Macaskill P, Kerr M, et al. Variability and accuracy in interpretation of consolidation on chest radiography for diagnosing pneumonia in children under 5 years of age. *Pediatr Pulmonol*. 2013;48(12):1195-1200.
35. Agrawal AK, Saini N, Gildengorin G, Feusner JH. Is routine computed tomographic scanning justified in the first week of persistent febrile neutropenia in children with malignancies? *Pediatr Blood Cancer*. 2011;57(4):620-624.
36. Archibald S, Park J, Geyer JR, Hawkins DS. Computed tomography in the evaluation of febrile neutropenic pediatric oncology patients. *Pediatr Infect Dis J*. 2001;20(1):5-10.
37. del Rosal T, Goycochea WA, Mendez-Echevarria A, et al. (1)(8)F-FDG PET/CT in the diagnosis of occult bacterial infections in children. *Eur J Pediatr*. 2013;172(8):1111-1115.
38. Jasper N, Dabritz J, Frosch M, Loeffler M, Weckesser M, Foell D. Diagnostic value of [(18)F]-FDG PET/CT in children with fever of unknown origin or unexplained signs of inflammation. *Eur J Nucl Med Mol Imaging*. 2010;37(1):136-145.
39. Blokhuis GJ, Bleeker-Rovers CP, Diender MG, Oyen WJ, Draaisma JM, de Geus-Oei LF. Diagnostic value of FDG-PET/(CT) in children with fever of unknown origin and unexplained fever during immune suppression. *Eur J Nucl Med Mol Imaging*. 2014;41(10):1916-1923.
40. Pantell RH, Newman TB, Bernzweig J, et al. Management and outcomes of care of fever in early infancy. *JAMA*. 2004;291(10):1203-1212.
41. Patterson RJ, Bisset GS, 3rd, Kirks DR, Vanness A. Chest radiographs in the evaluation of the febrile infant. *AJR Am J Roentgenol*. 1990;155(4):833-835.
42. Heulitt MJ, Ablow RC, Santos CC, O'Shea TM, Hilfer CL. Febrile infants less than 3 months old: value of chest radiography. *Radiology*. 1988;167(1):135-137.
43. Bramson RT, Meyer TL, Silbiger ML, Blickman JG, Halpern E. The futility of the chest radiograph in the febrile infant without respiratory symptoms. *Pediatrics*. 1993;92(4):524-526.
44. Baraff LJ, Bass JW, Fleisher GR, et al. Practice guideline for the management of infants and children 0 to 36 months of age with fever without source. Agency for Health Care Policy and Research. *Ann Emerg Med*. 1993;22(7):1198-1210.
45. McCarthy PL. The pediatric clinical evaluation and pneumonia. *Curr Opin Pediatr*. 1996;8(5):427-429.
46. Antonyrajah B, Mukundan D. Fever without apparent source on clinical examination. *Curr Opin Pediatr*. 2008;20(1):96-102.
47. Crain EF, Bulas D, Bijur PE, Goldman HS. Is a chest radiograph necessary in the evaluation of every febrile infant less than 8 weeks of age? *Pediatrics*. 1991;88(4):821-824.
48. Bramson RT, Griscom NT, Cleveland RH. Interpretation of chest radiographs in infants with cough and fever. *Radiology*. 2005;236(1):22-29.
49. Murphy CG, van de Pol AC, Harper MB, Bachur RG. Clinical predictors of occult pneumonia in the febrile child. *Acad Emerg Med*. 2007;14(3):243-249.
50. Leventhal JM. Clinical predictors of pneumonia as a guide to ordering chest roentgenograms. *Clin Pediatr (Phila)*. 1982;21(12):730-734.
51. Bleeker SE, Derksen-Lubsen G, Grobbee DE, Donders AR, Moons KG, Moll HA. Validating and updating a prediction rule for serious bacterial infection in patients with fever without source. *Acta Paediatr*. 2007;96(1):100-104.
52. Mahabee-Gittens EM, Grupp-Phelan J, Brody AS, et al. Identifying children with pneumonia in the emergency department. *Clin Pediatr (Phila)*. 2005;44(5):427-435.

53. Lynch T, Platt R, Gouin S, Larson C, Patenaude Y. Can we predict which children with clinically suspected pneumonia will have the presence of focal infiltrates on chest radiographs? *Pediatrics*. 2004;113(3 Pt 1):e186-189.
54. Losek JD, Kishaba RG, Berens RJ, Bonadio WA, Wells RG. Indications for chest roentgenogram in the febrile young infant. *Pediatr Emerg Care*. 1989;5(3):149-152.
55. Zukin DD, Hoffman JR, Cleveland RH, Kushner DC, Herman TE. Correlation of pulmonary signs and symptoms with chest radiographs in the pediatric age group. *Ann Emerg Med*. 1986;15(7):792-796.
56. Bachur R, Perry H, Harper MB. Occult pneumonias: empiric chest radiographs in febrile children with leukocytosis. *Ann Emerg Med*. 1999;33(2):166-173.
57. Black S, Shinefield H, Fireman B, et al. Efficacy, safety and immunogenicity of heptavalent pneumococcal conjugate vaccine in children. Northern California Kaiser Permanente Vaccine Study Center Group. *Pediatr Infect Dis J*. 2000;19(3):187-195.
58. Clinical policy for children younger than three years presenting to the emergency department with fever. *Ann Emerg Med*. 2003;42(4):530-545.
59. British Thoracic Society Guidelines for the Management of Community Acquired Pneumonia in Childhood. *Thorax*. 2002;57 Suppl 1:i1-24.
60. Paulus S, Dobson S. Febrile neutropenia in children with cancer. *Adv Exp Med Biol*. 2009;634:185-204.
61. Korones DN, Hussong MR, Gullace MA. Routine chest radiography of children with cancer hospitalized for fever and neutropenia: is it really necessary? *Cancer*. 1997;80(6):1160-1164.
62. Phillips B, Wade R, Westwood M, Riley R, Sutton AJ. Systematic review and meta-analysis of the value of clinical features to exclude radiographic pneumonia in febrile neutropenic episodes in children and young people. *J Paediatr Child Health*. 2012;48(8):641-648.
63. Hughes WT, Armstrong D, Bodey GP, et al. 2002 guidelines for the use of antimicrobial agents in neutropenic patients with cancer. *Clin Infect Dis*. 2002;34(6):730-751.
64. Cox JA, DeMasi J, McCollom S, Jackson G, Scothorn D, Aquino VM. The diagnostic utility of routine chest radiography in the evaluation of the initial fever in patients undergoing hematopoietic stem cell. *Pediatr Blood Cancer*. 2011;57(4):666-668.
65. Gasparetto TD, Escuissato DL, Marchiori E. Pulmonary infections following bone marrow transplantation: high-resolution CT findings in 35 paediatric patients. *Eur J Radiol*. 2008;66(1):117-121.
66. Abdulsalam AM, Al-Jahdali HH, Memish ZA, Ahmad AH. Fever of unknown origin. Experience of a large tertiary care hospital in Saudi Arabia. *Saudi Med J*. 2005;26(2):352-354.
67. Arce-Salinas CA, Morales-Velazquez JL, Villasenor-Ovies P, Muro-Cruz D. Classical fever of unknown origin (FUO): current causes in Mexico. *Rev Invest Clin*. 2005;57(6):762-769.
68. Ciftci E, Ince E, Dogru U. Pyrexia of unknown origin in children: a review of 102 patients from Turkey. *Ann Trop Paediatr*. 2003;23(4):259-263.
69. Tsukahara M, Tsuneoka H, Iino H, Murano I, Takahashi H, Uchida M. Bartonella henselae infection as a cause of fever of unknown origin. *J Clin Microbiol*. 2000;38(5):1990-1991.
70. Steele RW, Jones SM, Lowe BA, Glasier CM. Usefulness of scanning procedures for diagnosis of fever of unknown origin in children. *J Pediatr*. 1991;119(4):526-530.
71. Sheng JF, Sheng ZK, Shen XM, et al. Diagnostic value of fluorine-18 fluorodeoxyglucose positron emission tomography/computed tomography in patients with fever of unknown origin. *Eur J Intern Med*. 2011;22(1):112-116.
72. Simons KS, Pickkers P, Bleeker-Rovers CP, Oyen WJ, van der Hoeven JG. F-18-fluorodeoxyglucose positron emission tomography combined with CT in critically ill patients with suspected infection. *Intensive Care Med*. 2010;36(3):504-511.
73. Dong MJ, Zhao K, Liu ZF, Wang GL, Yang SY, Zhou GJ. A meta-analysis of the value of fluorodeoxyglucose-PET/PET-CT in the evaluation of fever of unknown origin. *Eur J Radiol*. 2011;80(3):834-844.
74. Hao R, Yuan L, Kan Y, Li C, Yang J. Diagnostic performance of 18F-FDG PET/CT in patients with fever of unknown origin: a meta-analysis. *Nucl Med Commun*. 2013;34(7):682-688.
75. Crouzet J, Boudousq V, Lechiche C, et al. Place of (18)F-FDG-PET with computed tomography in the diagnostic algorithm of patients with fever of unknown origin. *Eur J Clin Microbiol Infect Dis*. 2012;31(8):1727-1733.
76. Darge K, Jaramillo D, Siegel MJ. Whole-body MRI in children: current status and future applications. *Eur J Radiol*. 2008;68(2):289-298.

77. Korchi AM, Hanquinet S, Anooshiravani M, Merlini L. Whole-body magnetic resonance imaging: an essential tool for diagnosis and work up of non-oncological systemic diseases in children. *Minerva Pediatr.* 2014;66(3):169-176.
78. Ley S, Ley-Zaporozhan J, Schenk JP. Whole-body MRI in the pediatric patient. *Eur J Radiol.* 2009;70(3):442-451.
79. Purz S, Sabri O, Viehweger A, et al. Potential Pediatric Applications of PET/MR. *J Nucl Med.* 2014;55(Supplement 2):32S-39S.

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.