

**American College of Radiology  
ACR Appropriateness Criteria®  
Second and Third Trimester Vaginal Bleeding**

**Variant 1:                    Second and third trimester vaginal bleeding. Painless bleeding. Initial imaging.**

Procedure	Appropriateness Category	Relative Radiation Level
US duplex Doppler pelvis	Usually Appropriate	0
US pregnant uterus transabdominal	Usually Appropriate	0
US pregnant uterus transvaginal	Usually Appropriate	0
US cervix transperineal	May Be Appropriate	0

**Variant 2:                    Second and third trimester vaginal bleeding. Painful bleeding. Initial imaging.**

Procedure	Appropriateness Category	Relative Radiation Level
US duplex Doppler pelvis	Usually Appropriate	0
US pregnant uterus transabdominal	Usually Appropriate	0
US pregnant uterus transvaginal	Usually Appropriate	0
US cervix transperineal	May Be Appropriate	0

**Variant 3:                    Second and third trimester vaginal bleeding. Suspicion of or known placental previa, low-lying placental, or vasa previa. Initial imaging.**

Procedure	Appropriateness Category	Relative Radiation Level
US duplex Doppler pelvis	Usually Appropriate	0
US pregnant uterus transabdominal	Usually Appropriate	0
US pregnant uterus transvaginal	Usually Appropriate	0
US cervix transperineal	Usually Not Appropriate	0

## SECOND AND THIRD TRIMESTER VAGINAL BLEEDING

Expert Panel on GYN and OB Imaging: Thomas D. Shipp, MD, RDMS<sup>a</sup>; Liina Poder, MD<sup>b</sup>; Vickie A. Feldstein, MD<sup>c</sup>; Edward R. Oliver, MD, PhD<sup>d</sup>; Susan B. Promes, MD, MBA<sup>e</sup>; Loretta M. Strachowski, MD<sup>f</sup>; Betsy L. Sussman, MD<sup>g</sup>; Eileen Y. Wang, MD<sup>h</sup>; Therese M. Weber, MD<sup>i</sup>; Tom Winter, MD, MA<sup>j</sup>; Phyllis Glanc, MD.<sup>k</sup>

### Summary of Literature Review

#### **Introduction/Background**

Vaginal bleeding, or bleeding per vaginam, in the second and third trimesters of pregnancy is less common than in the first trimester but can similarly arise from an array of etiologies [1,2]. These diagnoses can widely range from having vaginal bleeding for no identifiable reason—seen in approximately half of those with vaginal bleeding in pregnancy [3]—to being the normal and expected bleeding with term labor, to having serious potential consequences for both the mother and fetus [4]. Although history and physical examination can identify many causes of vaginal bleeding, imaging, particularly with ultrasound (US), is the mainstay for making accurate diagnoses and for emergent guidance of management, which in the second and third trimesters may lead to hospitalization and/or delivery. Imaging is further critical to the evaluation of those with second and third trimester vaginal bleeding as digital pelvic examination should be avoided until diagnoses such as placenta previa, low-lying placenta, and vasa previa have been excluded.

The outcome of pregnancies with vaginal bleeding in the second and third trimesters depends on the precise etiology for the bleeding. Expected physiologic bleeding is seen with miscarriage and cervical change with labor. Bleeding accompanying miscarriage, which occurs in the first half of pregnancy, is common. As the cervix prepares for labor, there is frequently vaginal bleeding, termed the “bloody show” that typically precedes labor. Pathologic vaginal bleeding can be seen in a variety of cases including bleeding from the cervix with preterm labor, bleeding from placenta previa, vasa previa, placental abruption, or uterine rupture. The bleeding from any of these entities can lead to adverse perinatal and maternal outcomes, including the risk of death. Rare causes of bleeding include those arising from the cervix, such as cervicitis or cervical polyp, and the vagina, such as a laceration; these typically can be elucidated by the history and physical examination, though at times ultrasound may identify a polyp in the endocervical canal. Placental abruption affects approximately 1% of pregnancies, but US diagnosis is much less common and likely identifies, at most, 50% of cases of abruption though likely many fewer [5]. US imaging is vitally important as a central abruption is associated with worse perinatal outcome than a marginal separation of the placenta [6]. The most common diagnosis made is that of placenta previa, affecting approximately 1 in 200 pregnancies at delivery [7,8]. Vasa previa is much less frequent, affecting approximately 1 in 2,500 to 1 in 5,000 deliveries [8-10]. Uterine rupture is rare but can occur most commonly in patients with prior cesarean delivery or other uterine surgery. Vaginal bleeding can also occur with placental attachment disorder/placenta accreta spectrum; however, this is covered further in the ACR Appropriateness Criteria<sup>®</sup> topic on “[Suspected Placenta Accreta Spectrum Disorder](#)” [11].

Transabdominal US is the mainstay of obstetrical imaging; however, transvaginal US is particularly helpful given the not uncommon frequency with which cervical-related causes of second and third trimester vaginal bleeding occurs [12-14]. Furthermore, transvaginal US has been shown to be both accurate and safe for the diagnosis of placenta previa, as covered in more detail in the [ACR-ACOG-AIUM-SMFM-SRU Practice Parameter for the Performance of Standard Diagnostic Obstetrical Ultrasound](#) [15]. There has been literature documenting the utility of transperineal US (or translabial US) especially for evaluation of the cervix [16]; however, optimal visualization of the cervix occurs with closer approximation of the cervix, such as with real-time evaluation using transvaginal US [17] and as covered further in the ACR Appropriateness Criteria<sup>®</sup> topic on “[Assessment of Gravid Cervix](#)” [18].

<sup>a</sup>Brigham & Women’s Hospital, Boston, Massachusetts; American College of Obstetricians and Gynecologists. <sup>b</sup>Panel Chair, University of California San Francisco, San Francisco, California. <sup>c</sup>University of California San Francisco, San Francisco, California. <sup>d</sup>Children’s Hospital of Philadelphia and Perelman School of Medicine at the University of Pennsylvania, Philadelphia, Pennsylvania. <sup>e</sup>Penn State Health, Hershey, Pennsylvania; American College of Emergency Physicians. <sup>f</sup>University of California San Francisco, San Francisco, California. <sup>g</sup>The University of Vermont Medical Center, Burlington, Vermont. <sup>h</sup>Hospital of the University of Pennsylvania, Philadelphia, Pennsylvania; American College of Obstetricians and Gynecologists. <sup>i</sup>University of Alabama at Birmingham, Birmingham, Alabama. <sup>j</sup>University of Utah, Salt Lake City, Utah. <sup>k</sup>Specialty Chair, University of Toronto and Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada.

The American College of Radiology seeks and encourages collaboration with other organizations on the development of the ACR Appropriateness Criteria through society representation on expert panels. Participation by representatives from collaborating societies on the expert panel does not necessarily imply individual or society endorsement of the final document.

Reprint requests to: [publications@acr.org](mailto:publications@acr.org)

In the setting of premature rupture of membranes, there is a paucity of evidence on whether the introduction of the transvaginal probe may be associated with an increased risk of chorioamnionitis. Carlan et al [19] in a randomized trial of 92 patients did not find any associated increased risk. Nonetheless, caution should be applied in this setting.

### **Special Imaging Considerations**

Transrectal US is a useful technique and is generally well tolerated when transvaginal and transperineal US are either unsuccessful or declined [20,21].

### **Initial Imaging Definition**

Imaging at the beginning of the care episode for the medical condition defined by the variant. More than one procedure can be considered usually appropriate in the initial imaging evaluation when:

- There are procedures that are equivalent alternatives (ie, only one procedure will be ordered to provide the clinical information to effectively manage the patient's care)

OR

- There are complementary procedures (ie, more than one procedure is ordered as a set or simultaneously in which each procedure provides unique clinical information to effectively manage the patient's care).

### **Discussion of Procedures by Variant**

#### **Variant 1: Second and third trimester vaginal bleeding. Painless bleeding. Initial imaging.**

##### **US Cervix Transperineal**

The literature is sparse with regard to defined indications for the use of transperineal US or translabial US among those with painless vaginal bleeding. In current US use, if transabdominal US demonstrates an open cervix with bulging membranes or if the history and physical examination suggests ruptured membranes, then transperineal US might be considered. If transabdominal US is inconclusive and transvaginal US is declined by the patient, transperineal US or transrectal US may be useful for making an accurate diagnosis.

##### **US Duplex Doppler Pelvis**

Although Doppler velocimetry US is an adjunct to any of the other procedures for evaluation of painless vaginal bleeding, it can be invaluable for identification of pathologic entities such as vasa previa [22]. Vessels overlying the internal cervical os can be subtle and challenging to diagnose without optimal imaging conditions, and the use of color and spectral Doppler velocimetry US can be essential for making accurate diagnoses and distinguishing fetal from maternal vessels especially with the presence of fetal arterial vessels.

##### **US Pregnant Uterus Transabdominal**

The primary goal of transabdominal US in the setting of painless bleeding is to exclude pathologic etiologies for the bleeding. Impending miscarriage or early labor can typically be excluded via history and physical examination. The most common etiologies for painless vaginal bleeding would be placenta previa, and, less commonly, vasa previa [23]. Transabdominal US is the preferred initial imaging procedure as a screening tool for placenta previa [24]. This evaluation should include visualization of the placenta, the inferior placental margin, the placental umbilical cord insertion, and the cervix from the external os to the internal os. Any vessels overlying the internal cervical os should be specifically excluded. Transabdominal visualization of the lower uterine segment and cervix can be suboptimal because of inadequate urine in the bladder or from shadowing from the maternal symphysis pubis.

##### **US Pregnant Uterus Transvaginal**

Transvaginal US is frequently necessary for evaluation of painless vaginal bleeding, especially if transabdominal US is inconclusive or inadequate. As with transabdominal US, the evaluation should include visualization of the placenta if identifiable, the inferior placental margin, the cervix from the external os to the internal os, and the identification of any vessels overlying the internal cervical os. The evaluation should also include a measurement of the distance from the leading placental margin to the internal cervical os or the distance that the placenta crosses the internal cervical os [13]. In fact, in the second trimester, transvaginal US is predictive of placenta previa at delivery if the placenta crosses the internal cervical os by  $\geq 15$  mm [25], and an overlap of 25 mm or more at 20 to 23 weeks' gestation was associated with an exceptional risk for cesarean delivery [25,26]; a repeat sonographic evaluation later in pregnancy was suggested in these cases [21].

### **Variant 2: Second and third trimester vaginal bleeding. Painful bleeding. Initial imaging.**

Painful vaginal bleeding in the second and third trimesters includes an additional range of etiologies than that seen with painless vaginal bleeding, such as placental abruption, which is more commonly associated with abdominal pain. History and physical examination are vital for assessing for miscarriage or labor, whether term or preterm. Term labor would not necessarily prompt the need for imaging if the history and physical examination are consistent with this diagnosis. Should history and physical examination suggest miscarriage or preterm labor, evaluation would include an evaluation of the viability of the fetus and a detailed assessment of the cervix from external os to internal os, especially to assess cervical length. The bladder should not be overfilled to minimize the likelihood of falsely elongating the cervical length. Also, knowing the placental location will allow the clinician to know if it is safe to examine the patient's cervix digitally to assess for preterm labor.

#### **US Cervix Transperineal**

The literature is sparse in regard to defined indications for the use of transperineal US among those with painful vaginal bleeding. In current US use, if transabdominal US demonstrates an open cervix with bulging membranes or if the history and physical examination suggests ruptured membranes, then transperineal US might be considered. If transabdominal US is inconclusive and transvaginal US is declined by the patient, transperineal US may be useful for making an accurate diagnosis.

#### **US Duplex Doppler Pelvis**

Although Doppler velocimetry US is an adjunct to any of the other procedures for evaluation of painful vaginal bleeding, it can be invaluable for identification of pathologic entities like placental abruption. An acute clot within the uterus can have an echogenicity similar to that of the placenta; therefore, a placental abruption can be difficult to diagnose, especially prior to organization of the clot [5]. Identification of an area of interest contiguous with the placenta but with no blood flow as documented using color or power Doppler US could suggest the presence of an acute clot as seen with placental abruption. Although color or power Doppler US appears to be helpful with the diagnosis of placental abruption, spectral Doppler US of the fetal arterial vessels appears to be less helpful [27].

#### **US Pregnant Uterus Transabdominal**

Transabdominal US assessment is the preferred initial procedure for this evaluation and for assessment of the placenta. As in those with painless vaginal bleeding, an evaluation for placenta previa and vasa previa is indicated. The placenta is also comprehensively evaluated to assess its location, the presence of any clot as seen with placental separation in placental abruption, the placental location, and the presence of any vessels overlying the internal cervical os [23,24]. Although US is not sensitive for the diagnosis of placental abruption, the identification of a placental abruption is associated with worse perinatal outcomes [28]. The uterus should be evaluated for signs of uterine rupture, especially among those with a history of prior cesarean delivery. Although uterine rupture is typically a clinical diagnosis, imaging might be helpful in certain cases. Any disruption of the myometrium would suggest a uterine rupture. A threshold of 2.5-mm thickness of the lower uterine segment has been shown to be predictive of uterine dehiscence [29].

#### **US Pregnant Uterus Transvaginal**

Transvaginal US is frequently necessary for evaluation of painful vaginal bleeding especially if transabdominal US is inconclusive or inadequate. As with transabdominal US, the evaluation should include visualization of the placenta if identifiable, the inferior placental margin, the presence of any clot within the uterus, the continuity of the myometrium, the cervix from the external os to the internal os, and the identification of any vessels overlying the internal cervical os. Although a transvaginally identified short cervix is well known to be related to preterm delivery, associated vaginal bleeding significantly increases the risk of preterm delivery [30]. Evaluation of the lower uterine segment, the uterine isthmus [31], during transvaginal US can predict bleeding during pregnancy and delivery among those with placenta previa [32].

### **Variant 3: Second and third trimester vaginal bleeding. Suspicion of or known placental previa, low-lying placental, or vasa previa. Initial imaging.**

#### **US Cervix Transperineal**

To our knowledge, there is currently a paucity of literature to support the use of transperineal US for the initial evaluation of placenta previa, low-lying placenta, or vasa previa [33].

#### **US Duplex Doppler Pelvis**

Although Doppler velocimetry US is an adjunct to transabdominal US and transvaginal US for evaluation of known or suspected placenta previa, low-lying placenta, or vasa previa, it can be invaluable for identification of pathologic

entities like vasa previa. Vessels overlying the internal cervical os can be subtle and challenging to diagnose without optimal imaging conditions, and the use of Doppler velocimetry US can be essential for making this diagnosis [7,34]. Furthermore, spectral Doppler may play an important role in distinguishing fetal arterial vessels from maternal vessels when overlying the internal cervical os.

### **US Pregnant Uterus Transabdominal**

Evaluation of second or third trimester vaginal bleeding with known or suspected placenta previa, low-lying placenta, or vasa previa initially involves transabdominal US. The placenta is comprehensively evaluated to assess its location, the inferior placental margin and how far the placenta lies from the internal os or how far the placenta overlaps the internal cervical os [35], the placental umbilical cord insertion, and the cervix, especially of the internal os. If present, any vessels overlying the internal cervical os should be searched for and identified. If, during transabdominal US at 18 to 20 weeks' gestation, the placenta is  $\geq 2$  cm from the internal cervical os, follow-up evaluation of the placenta is not necessary [36,37]. The later in gestation that a placenta previa is diagnosed, the more likely it will remain a placenta previa at delivery; however, even with a scan at 32 to 35 weeks' gestation, only 73% of placenta previas persist at delivery [38]. Twins are also at a 40% increased risk for placenta previa than singletons, highlighting the need to be vigilant in this population [39]. Other risk factors for placenta previa include increasing maternal age, parity, prior cesarean delivery, prior spontaneous abortion, and maternal tobacco use [40,41]. A low-lying placenta that is  $< 2$  cm from the internal cervical os has a  $< 2\%$  chance of remaining low-lying at delivery; nonetheless, these patients whose placentas remain low are at increased risk for cesarean delivery [42].

### **US Pregnant Uterus Transvaginal**

Transvaginal US is frequently necessary for evaluation of known or suspected placenta previa, low-lying placenta, or vasa previa, especially if transabdominal US is inconclusive or inadequately demonstrates the internal cervical os and inferior placental margin. As with transabdominal US, the evaluation should include visualization of the placenta if identifiable, the inferior placental margin, the cervix from the external os to the internal os, and the identification of any vessels overlying the internal cervical os. Among those with placenta previa, a shortening cervix is associated with an increased risk for emergency cesarean delivery [43], and a transvaginal cervical length of  $< 35$  mm was associated with an increased risk of preterm cesarean delivery due to massive hemorrhage [44]. As placenta previa and low-lying placenta can resolve over gestation, vasa previa has been reported to resolve in 5.8% to 23.8% of cases [45,46].

### **Summary of Recommendations**

- **Variant 1:** US pregnant uterus transabdominal, US duplex Doppler pelvis, and US pregnant uterus transvaginal are usually appropriate for the initial imaging of painless second and third trimester vaginal bleeding. These procedures are complementary (ie, more than one procedure is ordered as a set or simultaneously where each procedure provides unique clinical information to effectively manage the patient's care).
- **Variant 2:** US pregnant uterus transabdominal, US duplex Doppler pelvis, and US pregnant uterus transvaginal are usually appropriate for the initial imaging of painful second and third trimester vaginal bleeding. These procedures are complementary (ie, more than one procedure is ordered as a set or simultaneously where each procedure provides unique clinical information to effectively manage the patient's care).
- **Variant 3:** US pregnant uterus transabdominal, US duplex Doppler pelvis, and US pregnant uterus transvaginal are usually appropriate for the initial imaging of second and third trimester vaginal bleeding with suspicion of or known placenta previa, low-lying placental, or vasa previa. These procedures are complementary (ie, more than one procedure is ordered as a set or simultaneously where each procedure provides unique clinical information to effectively manage the patient's care).

### **Supporting Documents**

The evidence table, literature search, and appendix for this topic are available at <https://acsearch.acr.org/list>. The appendix includes the strength of evidence assessment and the final rating round tabulations for each recommendation.

For additional information on the Appropriateness Criteria methodology and other supporting documents go to [www.acr.org/ac](http://www.acr.org/ac).

## Safety Considerations in Pregnant Patients

Imaging of the pregnant patient can be challenging, particularly with respect to minimizing radiation exposure and risk. For further information and guidance, see the following ACR documents:

- [ACR–SPR Practice Parameter for the Safe and Optimal Performance of Fetal Magnetic Resonance Imaging \(MRI\)](#) [47]
- [ACR-SPR Practice Parameter for Imaging Pregnant or Potentially Pregnant Adolescents and Women with Ionizing Radiation](#) [48]
- [ACR-ACOG-AIUM-SMFM-SRU Practice Parameter for the Performance of Standard Diagnostic Obstetrical Ultrasound](#) [15]
- [ACR Manual on Contrast Media](#) [49]
- [ACR Guidance Document on MR Safe Practices: 2013](#) [50]

## Appropriateness Category Names and Definitions

Appropriateness Category Name	Appropriateness Rating	Appropriateness Category Definition
Usually Appropriate	7, 8, or 9	The imaging procedure or treatment is indicated in the specified clinical scenarios at a favorable risk-benefit ratio for patients.
May Be Appropriate	4, 5, or 6	The imaging procedure or treatment may be indicated in the specified clinical scenarios as an alternative to imaging procedures or treatments with a more favorable risk-benefit ratio, or the risk-benefit ratio for patients is equivocal.
May Be Appropriate (Disagreement)	5	The individual ratings are too dispersed from the panel median. The different label provides transparency regarding the panel’s recommendation. “May be appropriate” is the rating category and a rating of 5 is assigned.
Usually Not Appropriate	1, 2, or 3	The imaging procedure or treatment is unlikely to be indicated in the specified clinical scenarios, or the risk-benefit ratio for patients is likely to be unfavorable.

## Relative Radiation Level Information

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Patients in the pediatric age group are at inherently higher risk from exposure, because of both organ sensitivity and longer life expectancy (relevant to the long latency that appears to accompany radiation exposure). For these reasons, the RRL dose estimate ranges for pediatric examinations are lower as compared with those specified for adults (see Table below). Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® [Radiation Dose Assessment Introduction](#) document [51].

Relative Radiation Level Designations		
Relative Radiation Level*	Adult Effective Dose Estimate Range	Pediatric Effective Dose Estimate Range
○	0 mSv	0 mSv
⊕	<0.1 mSv	<0.03 mSv
⊕⊕	0.1-1 mSv	0.03-0.3 mSv
⊕⊕⊕	1-10 mSv	0.3-3 mSv
⊕⊕⊕⊕	10-30 mSv	3-10 mSv
⊕⊕⊕⊕⊕	30-100 mSv	10-30 mSv

\*RRL assignments for some of the examinations cannot be made, because the actual patient doses in these procedures vary as a function of a number of factors (eg, region of the body exposed to ionizing radiation, the imaging guidance that is used). The RRLs for these examinations are designated as “Varies.”

## References

- Norwitz ER, Park JS. Overview of the etiology and evaluation of vaginal bleeding in pregnant women. Available at: <https://www.uptodate.com/contents/overview-of-the-etiology-and-evaluation-of-vaginal-bleeding-in-pregnant-women>. Accessed March 27, 2020.
- Society for Maternal-Fetal Medicine . Electronic address pso, Gyamfi-Bannerman C. Society for Maternal-Fetal Medicine (SMFM) Consult Series #44: Management of bleeding in the late preterm period. Am J Obstet Gynecol 2018;218:B2-B8.
- Yang J, Hartmann KE, Savitz DA, et al. Vaginal bleeding during pregnancy and preterm birth. Am J Epidemiol 2004;160:118-25.
- Vahanian SA, Vintzileos AM. Placental implantation abnormalities: a modern approach. Curr Opin Obstet Gynecol 2016;28:477-84.
- Oyelese Y, Ananth CV. Placental abruption. Obstet Gynecol 2006;108:1005-16.
- Nkwabong E, Tiomela Goula G. Placenta abruption surface and perinatal outcome. J Matern Fetal Neonatal Med 2017;30:1456-59.
- Reddy UM, Abuhamad AZ, Levine D, Saade GR, Fetal Imaging Workshop Invited P. Fetal imaging: executive summary of a joint Eunice Kennedy Shriver National Institute of Child Health and Human Development, Society for Maternal-Fetal Medicine, American Institute of Ultrasound in Medicine, American College of Obstetricians and Gynecologists, American College of Radiology, Society for Pediatric Radiology, and Society of Radiologists in Ultrasound Fetal Imaging workshop. Obstet Gynecol 2014;123:1070-82.
- Silver RM. Abnormal Placentation: Placenta Previa, Vasa Previa, and Placenta Accreta. Obstet Gynecol 2015;126:654-68.
- Baumfeld Y, Gutvirtz G, Shoham I, Sheiner E. Fetal heart rate patterns of pregnancies with vasa previa and velamentous cord insertion. Arch Gynecol Obstet 2016;293:361-7.
- Society of Maternal-Fetal Publications C, Sinkey RG, Odibo AO, Dashe JS. #37: Diagnosis and management of vasa previa. Am J Obstet Gynecol 2015;213:615-9.
- American College of Radiology. ACR Appropriateness Criteria®: Suspected Placenta Accreta Spectrum Disorder. Available at: <https://acsearch.acr.org/docs/3102403/Narrative/>. Accessed March 27, 2020.
- Leerentveld RA, Gilberts EC, Arnold MJ, Wladimiroff JW. Accuracy and safety of transvaginal sonographic placental localization. Obstet Gynecol 1990;76:759-62.
- Oppenheimer LW, Farine D, Ritchie JW, Lewinsky RM, Telford J, Fairbanks LA. What is a low-lying placenta? Am J Obstet Gynecol 1991;165:1036-8.
- Timor-Tritsch IE, Monteagudo A. Diagnosis of placenta previa by transvaginal sonography. Ann Med 1993;25:279-83.
- American College of Radiology. ACR-ACOG-AIUM-SMFM-SRU Practice Parameter for the Performance of Standard Diagnostic Obstetrical Ultrasound. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/us-ob.pdf>. Accessed March 27, 2020.
- Hertzberg BS, Bowie JD, Carroll BA, Kliever MA, Weber TM. Diagnosis of placenta previa during the third trimester: role of transperineal sonography. AJR Am J Roentgenol 1992;159:83-7.



17. Vintzileos AM, Ananth CV, Smulian JC. Using ultrasound in the clinical management of placental implantation abnormalities. *Am J Obstet Gynecol* 2015;213:S70-7.
18. American College of Radiology. ACR Appropriateness Criteria®: Assessment of Gravid Cervix. Available at: <https://acsearch.acr.org/docs/69464/Narrative/>. Accessed March 27, 2020.
19. Carlan SJ, Richmond LB, O'Brien WF. Randomized trial of endovaginal ultrasound in preterm premature rupture of membranes. *Obstet Gynecol* 1997;89:458-61.
20. Timor-Tritsch IE, Monteagudo A, Rebarber A, Goldstein SR, Tsybal T. Transrectal scanning: an alternative when transvaginal scanning is not feasible. *Ultrasound Obstet Gynecol* 2003;21:473-9.
21. Wachsberg RH. Transrectal ultrasonography for problem solving after transvaginal ultrasonography of the female internal reproductive tract. *J Ultrasound Med* 2003;22:1349-56.
22. Nomiyama M, Toyota Y, Kawano H. Antenatal diagnosis of velamentous umbilical cord insertion and vasa previa with color Doppler imaging. *Ultrasound Obstet Gynecol* 1998;12:426-9.
23. D'Antonio F, Bhide A. Ultrasound in placental disorders. *Best Pract Res Clin Obstet Gynaecol* 2014;28:429-42.
24. Quant HS, Friedman AM, Wang E, Parry S, Schwartz N. Transabdominal ultrasonography as a screening test for second-trimester placenta previa. *Obstet Gynecol* 2014;123:628-33.
25. Taipale P, Hiilesmaa V, Ylostalo P. Transvaginal ultrasonography at 18-23 weeks in predicting placenta previa at delivery. *Ultrasound Obstet Gynecol* 1998;12:422-5.
26. Becker RH, Vonk R, Mende BC, Ragoesch V, Entezami M. The relevance of placental location at 20-23 gestational weeks for prediction of placenta previa at delivery: evaluation of 8650 cases. *Ultrasound Obstet Gynecol* 2001;17:496-501.
27. Morales-Rosello J, Khalil A, Akhoundova F, et al. Fetal cerebral and umbilical Doppler in pregnancies complicated by late-onset placental abruption. *J Matern Fetal Neonatal Med* 2017;30:1320-24.
28. Glantz C, Purnell L. Clinical utility of sonography in the diagnosis and treatment of placental abruption. *J Ultrasound Med* 2002;21:837-40.
29. Uharcsek P, Brestansky A, Ravinger J, Manova A, Zajacova M. Sonographic assessment of lower uterine segment thickness at term in women with previous cesarean delivery. *Arch Gynecol Obstet* 2015;292:609-12.
30. Ramaeker DM, Simhan HN. Sonographic cervical length, vaginal bleeding, and the risk of preterm birth. *Am J Obstet Gynecol* 2012;206:224 e1-4.
31. Hasegawa J, Kawabata I, Takeda Y, et al. Improving the Accuracy of Diagnosing Placenta Previa on Transvaginal Ultrasound by Distinguishing between the Uterine Isthmus and Cervix: A Prospective Multicenter Observational Study. *Fetal Diagn Ther* 2017;41:145-51.
32. Goto M, Hasegawa J, Arakaki T, et al. Placenta previa with early opening of the uterine isthmus is associated with high risk of bleeding during pregnancy, and massive haemorrhage during caesarean delivery. *Eur J Obstet Gynecol Reprod Biol* 2016;201:7-11.
33. Gouhar GK, Sadek SM, Siam S, Ahmad RA. Role of transperineal sonography in diagnosis of placenta previa/accreta: A prospective study. *The Egyptian Journal of Radiology and Nuclear Medicine* 2012;43:637-45.
34. Ruiters L, Kok N, Limpens J, et al. Systematic review of accuracy of ultrasound in the diagnosis of vasa previa. *Ultrasound Obstet Gynecol* 2015;45:516-22.
35. Oppenheimer L, Maternal Fetal Medicine C. Diagnosis and management of placenta previa. *J Obstet Gynaecol Can* 2007;29:261-66.
36. Kapoor S, Thomas JT, Petersen SG, Gardener GJ. Is the third trimester repeat ultrasound scan for placental localisation needed if the placenta is low lying but clear of the os at the mid-trimester morphology scan? *Aust N Z J Obstet Gynaecol* 2014;54:428-32.
37. Olive EC, Roberts CL, Nassar N, Algert CS. Test characteristics of placental location screening by transabdominal ultrasound at 18-20 weeks. *Ultrasound Obstet Gynecol* 2006;28:944-9.
38. Dashe JS, McIntire DD, Ramus RM, Santos-Ramos R, Twickler DM. Persistence of placenta previa according to gestational age at ultrasound detection. *Obstet Gynecol* 2002;99:692-7.
39. Ananth CV, Demissie K, Smulian JC, Vintzileos AM. Placenta previa in singleton and twin births in the United States, 1989 through 1998: a comparison of risk factor profiles and associated conditions. *Am J Obstet Gynecol* 2003;188:275-81.
40. Faiz AS, Ananth CV. Etiology and risk factors for placenta previa: an overview and meta-analysis of observational studies. *J Matern Fetal Neonatal Med* 2003;13:175-90.



41. Gilliam M, Rosenberg D, Davis F. The likelihood of placenta previa with greater number of cesarean deliveries and higher parity. *Obstet Gynecol* 2002;99:976-80.
42. Heller HT, Mullen KM, Gordon RW, Reiss RE, Benson CB. Outcomes of pregnancies with a low-lying placenta diagnosed on second-trimester sonography. *J Ultrasound Med* 2014;33:691-6.
43. Shin JE, Shin JC, Lee Y, Kim SJ. Serial Change in Cervical Length for the Prediction of Emergency Cesarean Section in Placenta Previa. *PLoS One* 2016;11:e0149036.
44. Sekiguchi A, Nakai A, Okuda N, Inde Y, Takeshita T. Consecutive cervical length measurements as a predictor of preterm cesarean section in complete placenta previa. *J Clin Ultrasound* 2015;43:17-22.
45. Rebarber A, Dolin C, Fox NS, Klauser CK, Saltzman DH, Roman AS. Natural history of vasa previa across gestation using a screening protocol. *J Ultrasound Med* 2014;33:141-7.
46. Swank ML, Garite TJ, Maurel K, et al. Vasa previa: diagnosis and management. *Am J Obstet Gynecol* 2016;215:223 e1-6.
47. American College of Radiology. ACR–SPR Practice Parameter for the Safe and Optimal Performance of Fetal Magnetic Resonance Imaging (MRI). Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/mr-fetal.pdf>. Accessed March 27, 2020.
48. American College of Radiology. ACR-SPR Practice Parameter for Imaging Pregnant or Potentially Pregnant Adolescents and Women with Ionizing Radiation. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/pregnant-pts.pdf>. Accessed March 27, 2020.
49. American College of Radiology. *Manual on Contrast Media*. Available at: <https://www.acr.org/Clinical-Resources/Contrast-Manual>. Accessed March 27, 2020.
50. Expert Panel on MRS, Kanal E, Barkovich AJ, et al. ACR guidance document on MR safe practices: 2013. *J Magn Reson Imaging* 2013;37:501-30.
51. American College of Radiology. ACR Appropriateness Criteria® Radiation Dose Assessment Introduction. Available at: <https://www.acr.org/-/media/ACR/Files/Appropriateness-Criteria/RadiationDoseAssessmentIntro.pdf>. Accessed March 27, 2020.

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.