

**American College of Radiology
ACR Appropriateness Criteria®
Hearing Loss and/or Vertigo**

**Variant 1: Acquired conductive hearing loss in absence of clinically evident mass in the middle ear.
Initial imaging.**

Procedure	Appropriateness Category	Relative Radiation Level
CT temporal bone without IV contrast	Usually Appropriate	☼☼☼
CT head with IV contrast	Usually Not Appropriate	☼☼☼
CT head without and with IV contrast	Usually Not Appropriate	☼☼☼
CT head without IV contrast	Usually Not Appropriate	☼☼☼
CT temporal bone without and with IV contrast	Usually Not Appropriate	☼☼☼
CT temporal bone with IV contrast	Usually Not Appropriate	☼☼☼
CTA head with IV contrast	Usually Not Appropriate	☼☼☼
MR venography head without IV contrast	Usually Not Appropriate	○
MRA head without and with IV contrast	Usually Not Appropriate	○
MRA head without IV contrast	Usually Not Appropriate	○
MRI head and internal auditory canal without and with IV contrast	Usually Not Appropriate	○
MRI head and internal auditory canal without IV contrast	Usually Not Appropriate	○

Variant 2: Acquired conductive hearing loss secondary to cholesteatoma or neoplasm with suspected intracranial or inner ear extension. Surgical planning.

Procedure	Appropriateness Category	Relative Radiation Level
CT temporal bone without IV contrast	Usually Appropriate	☼☼☼
MRI head and internal auditory canal without and with IV contrast	Usually Appropriate	○
MRI head and internal auditory canal without IV contrast	May Be Appropriate	○
CT temporal bone with IV contrast	May Be Appropriate	☼☼☼
CTA head with IV contrast	May Be Appropriate (Disagreement)	☼☼☼
CT head with IV contrast	Usually Not Appropriate	☼☼☼
CT head without and with IV contrast	Usually Not Appropriate	☼☼☼
CT head without IV contrast	Usually Not Appropriate	☼☼☼
CT temporal bone without and with IV contrast	Usually Not Appropriate	☼☼☼
MR venography head with IV contrast	Usually Not Appropriate	○
MR venography head without IV contrast	Usually Not Appropriate	○
MRA head without and with IV contrast	Usually Not Appropriate	○
MRA head without IV contrast	Usually Not Appropriate	○

Variant 3:**Acquired sensorineural hearing loss. Initial imaging.**

Procedure	Appropriateness Category	Relative Radiation Level
MRI head and internal auditory canal without and with IV contrast	Usually Appropriate	○
MRI head and internal auditory canal without IV contrast	Usually Appropriate	○
CT temporal bone without IV contrast	May Be Appropriate	⊕⊕⊕
CT temporal bone with IV contrast	May Be Appropriate	⊕⊕⊕
CT head with IV contrast	Usually Not Appropriate	⊕⊕⊕
CT head without and with IV contrast	Usually Not Appropriate	⊕⊕⊕
CT head without IV contrast	Usually Not Appropriate	⊕⊕⊕
CT temporal bone without and with IV contrast	Usually Not Appropriate	⊕⊕⊕
CTA head with IV contrast	Usually Not Appropriate	⊕⊕⊕
MR venography head with IV contrast	Usually Not Appropriate	○
MR venography head without IV contrast	Usually Not Appropriate	○
MRA head without and with IV contrast	Usually Not Appropriate	○
MRA head without IV contrast	Usually Not Appropriate	○

Variant 4:**Mixed conductive and sensorineural hearing loss. Initial imaging.**

Procedure	Appropriateness Category	Relative Radiation Level
CT temporal bone without IV contrast	Usually Appropriate	⊕⊕⊕
MRI head and internal auditory canal without and with IV contrast	Usually Appropriate	○
MRI head and internal auditory canal without IV contrast	Usually Appropriate	○
CT head with IV contrast	Usually Not Appropriate	⊕⊕⊕
CT head without and with IV contrast	Usually Not Appropriate	⊕⊕⊕
CT head without IV contrast	Usually Not Appropriate	⊕⊕⊕
CT temporal bone with IV contrast	Usually Not Appropriate	⊕⊕⊕
CT temporal bone without and with IV contrast	Usually Not Appropriate	⊕⊕⊕
CTA head with IV contrast	Usually Not Appropriate	⊕⊕⊕
MR venography head with IV contrast	Usually Not Appropriate	○
MR venography head without IV contrast	Usually Not Appropriate	○
MRA head without and with IV contrast	Usually Not Appropriate	○
MRA head without IV contrast	Usually Not Appropriate	○

Variant 5: Congenital hearing loss or total deafness or cochlear implant candidate. Surgical planning.

Procedure	Appropriateness Category	Relative Radiation Level
CT temporal bone without IV contrast	Usually Appropriate	☼☼☼
MRI head and internal auditory canal without and with IV contrast	Usually Appropriate	○
MRI head and internal auditory canal without IV contrast	Usually Appropriate	○
CT head with IV contrast	Usually Not Appropriate	☼☼☼
CT head without and with IV contrast	Usually Not Appropriate	☼☼☼
CT head without IV contrast	Usually Not Appropriate	☼☼☼
CT temporal bone with IV contrast	Usually Not Appropriate	☼☼☼
CT temporal bone without and with IV contrast	Usually Not Appropriate	☼☼☼
CTA head with IV contrast	Usually Not Appropriate	☼☼☼
MR venography head with IV contrast	Usually Not Appropriate	○
MR venography head without IV contrast	Usually Not Appropriate	○
MRA head without and with IV contrast	Usually Not Appropriate	○
MRA head without IV contrast	Usually Not Appropriate	○

Variant 6: Episodic vertigo with or without associated hearing loss or aural fullness (peripheral vertigo). Initial imaging.

Procedure	Appropriateness Category	Relative Radiation Level
CT temporal bone without IV contrast	Usually Appropriate	☼☼☼
MRI head and internal auditory canal without and with IV contrast	Usually Appropriate	○
MRI head and internal auditory canal without IV contrast	Usually Appropriate	○
CTA head and neck with IV contrast	May Be Appropriate	☼☼☼
MRA head and neck without and with IV contrast	May Be Appropriate	○
CT head with IV contrast	Usually Not Appropriate	☼☼☼
CT head without and with IV contrast	Usually Not Appropriate	☼☼☼
CT head without IV contrast	Usually Not Appropriate	☼☼☼
CT temporal bone with IV contrast	Usually Not Appropriate	☼☼☼
CT temporal bone without and with IV contrast	Usually Not Appropriate	☼☼☼
MR venography head with IV contrast	Usually Not Appropriate	○
MR venography head without IV contrast	Usually Not Appropriate	○
MRA head and neck without IV contrast	Usually Not Appropriate	○

Variant 7:**Persistent vertigo with or without neurological symptoms (central vertigo). Initial imaging.**

Procedure	Appropriateness Category	Relative Radiation Level
MRI head and internal auditory canal without and with IV contrast	Usually Appropriate	○
MRI head and internal auditory canal without IV contrast	Usually Appropriate	○
CT head without IV contrast	May Be Appropriate	⊕⊕⊕
CTA head and neck with IV contrast	May Be Appropriate	⊕⊕⊕
MRA head and neck without and with IV contrast	May Be Appropriate	○
MRA head and neck without IV contrast	May Be Appropriate	○
CT head with IV contrast	May Be Appropriate	⊕⊕⊕
CT head without and with IV contrast	Usually Not Appropriate	⊕⊕⊕
CT temporal bone with IV contrast	Usually Not Appropriate	⊕⊕⊕
CT temporal bone without and with IV contrast	Usually Not Appropriate	⊕⊕⊕
CT temporal bone without IV contrast	Usually Not Appropriate	⊕⊕⊕
MR venography head with IV contrast	Usually Not Appropriate	○
MR venography head without IV contrast	Usually Not Appropriate	○

HEARING LOSS AND/OR VERTIGO

Expert Panel on Neurologic Imaging: Aseem Sharma, MD^a; Claudia F. E. Kirsch, MD^b; Joseph M. Aulino, MD^c; Santanu Chakraborty, MBBS, MSc^d; Asim F. Choudhri, MD^e; Isabelle M. Germano, MD^f; A. Tuba Kendi, MD^g; H. Jeffrey Kim, MD^h; Ryan K. Lee, MD, MRMD, MBAⁱ; David S. Liebeskind, MD^j; Michael D. Luttrull, MD^k; Toshio Moritani, MD, PhD^l; Gregory J. A. Murad, MD^m; Lubdha M. Shah, MDⁿ; Robert Y. Shih, MD^o; Sophia C. Symko, MD, MS^p; Julie Bykowski, MD.^q

Summary of Literature Review

Introduction/Background

Clinical assessment and audiometric testing can determine the type of hearing loss as conductive, sensorineural, or mixed [1,2] and guide the appropriateness of subsequent imaging. Conductive hearing loss results from diseases affecting the conduction of mechanical sound wave energy to the hair cells of the organ of Corti within the cochlea. These serve as the auditory receptors, converting the mechanical energy of sound waves into electrical neural impulses that are then transmitted along the auditory pathways to the auditory cortex [1]. Sensorineural hearing loss is caused by diseases that impair the cochlear function or the transmission of electrical signal along the auditory pathway, including the cranial nerve nucleus in the brainstem through the superior olive, inferior colliculus, medial geniculate body of the thalamus, and auditory cortex in the temporal lobe.

Given the proximity of the cranial nerves and their nuclei, disorders that affect hearing may also affect vestibular function and vice versa. The vestibule and semicircular canals are the end organs responsible for balance and equilibrium. Central vestibular pathways involve extensive connections between the vestibular nuclei within the brainstem and the cerebellum, extraocular nuclei, and spinal cord. Vertigo is a sensation that you or the environment around you is moving or spinning. Although vertigo often indicates dysfunction of the vestibule or semicircular canals, patients commonly report dizziness, a less specific term that may imply disequilibrium, light-headedness, or presyncope [3-5]. Accordingly, imaging workup in these patients may require assessment for disease processes that produce symptoms reported as dizziness rather than vertigo. Appropriateness of imaging often depends upon clinical categorization of vertigo into peripheral (vestibular) and central (affecting central vestibular pathways) categories based upon factors such as onset, duration, persistence, aggravating factors, and results of clinical testing [3-7]. In some cases however, this categorization may be difficult on clinical assessment, especially in less subspecialized care [3].

Discussion of Procedures by Variant

Variant 1: Acquired conductive hearing loss in absence of clinically evident mass in the middle ear. Initial imaging.

CT Temporal Bone

Temporal bone CT is considered to be the first-line imaging modality in patients presenting with conductive hearing loss without any mass lesion seen within the middle ear cavity [1,2,5]. CT provides excellent delineation of the external auditory canal, ossicular chain, and the bony labyrinth of the inner ear structures. CT is helpful in identifying changes of otospongiosis (otosclerosis), ossicular erosion or fusion, round window occlusion, and dehiscence of the superior semicircular canal [1,2,5,8-12]. The latter can produce conductive hearing loss and Tullio phenomenon, or sound-induced vertigo, by dissipating mechanical energy through the bony dehiscence. If this condition is suspected, performing CT reconstructions along the Pöschl plane and perpendicular to the Stenver plane then the orientation of the superior semicircular canal may be helpful in identification of bony dehiscence and estimating its extent [8,9].

^aMallinckrodt Institute of Radiology, Saint Louis, Missouri. ^bPanel Chair, North Shore–Long Island Jewish Hospital, Hofstra Medical School, Hempstead, New York. ^cVanderbilt University Medical Center, Nashville, Tennessee. ^dOttawa Hospital Research Institute and the Department of Radiology, The University of Ottawa, Ottawa, Ontario, Canada. ^eLe Bonheur Children’s Hospital, University of Tennessee Health Science Center, Memphis, Tennessee. ^fMount Sinai School of Medicine, New York, New York; neurosurgical consultant. ^gMayo Clinic, Rochester, Minnesota. ^hGeorgetown University Hospital, Washington, District of Columbia; American Academy of Otolaryngology-Head and Neck Surgery. ⁱEinstein Healthcare Network, Philadelphia, Pennsylvania. ^jUniversity of California Los Angeles, Los Angeles, California; American Academy of Neurology. ^kThe Ohio State University Wexner Medical Center, Columbus, Ohio. ^lUniversity of Iowa Hospitals and Clinics, Iowa City, Iowa. ^mUniversity of Florida, Gainesville, Florida; neurosurgical consultant. ⁿUniversity of Utah, Salt Lake City, Utah. ^oWalter Reed National Military Medical Center, Bethesda, Maryland. ^pNeuroradiology consultant, Denver, Colorado. ^qSpecialty Chair, UC San Diego Health, San Diego, California.

The American College of Radiology seeks and encourages collaboration with other organizations on the development of the ACR Appropriateness Criteria through society representation on expert panels. Participation by representatives from collaborating societies on the expert panel does not necessarily imply individual or society endorsement of the final document.

Reprint requests to: publications@acr.org

Given the density of temporal bone and the rather small size of individual structures of interest, such as ossicles, details of temporal bone morphology are only evident on bone windows. Accordingly, intravenous (IV) contrast is not beneficial for evaluation of temporal bone in patients with conductive hearing loss.

CT Head

There is no evidence to support use of CT head in patients with conductive hearing loss.

CTA Head

There is no evidence to support use of CT angiography (CTA) in patients with conductive hearing loss.

MRI Head and Internal Auditory Canal

MRI of the temporal bone is insufficient in delineation of the bony details usually needed for evaluation of patients with conductive hearing loss, and there is no evidence to support its use as a first-line imaging modality in these patients.

MRA Head

There is no evidence to support use of MR angiography (MRA) for initial evaluation of patients with conductive hearing loss.

MR Venography Head

There is no evidence to support use of MR venography (MRV) for initial evaluation of patients with conductive hearing loss.

Variant 2: Acquired conductive hearing loss secondary to cholesteatoma or neoplasm with suspected intracranial or inner ear extension. Surgical planning.

CT Temporal Bone

High-spatial resolution CT of the temporal bone is helpful in defining small inflammatory or neoplastic masses within the middle ear cavity [1,2,13]. In addition, CT can help in surgical planning by demonstrating erosions of ossicles or other inner ear structures (such as perilymphatic fistulae) caused by such masses [14]. Given the surrounding dense bone, IV contrast is usually not beneficial in studying enhancement characteristics of middle ear masses. However, contrast enhancement may help delineate extraosseous soft tissue associated with invasive neoplasms.

CT Head

There is no evidence to support use of CT head for assessment of patients with conductive hearing loss and middle ear mass identified on otoscopy.

CTA Head

There is no definite evidence to support use of CTA as a first-line modality for assessment of patients with conductive hearing loss and middle ear mass identified on otoscopy. However, in patients with high clinical suspicion of middle ear paraganglioma, CTA is sometimes used initially for diagnostic confirmation and for planning further management.

MRI Head and Internal Auditory Canal

Extent of a middle ear cavity mass identified on otoscopy in a patient with conductive hearing loss is much better defined using MRI obtained without and with IV contrast [2,5,13,15]. This assessment is better done using thin sections across the temporal bone as part of a dedicated internal auditory canal (IAC) protocol rather than a routine brain MRI. Excellent soft-tissue contrast afforded by even a noncontrast MRI often complements the bony details seen on temporal bone CT for complete evaluation of such patients prior to surgical intervention.

MRA Head

MRA is usually not used as a first-line imaging modality in patients presenting with conductive hearing loss. However, it may be helpful in assessing patency of the carotid artery if initial imaging raises suspicion of vascular involvement.

MR Venography Head

Although not used as the initial imaging modality, MRV may be helpful in assessing patency of jugular vein for surgical planning in patients with documented middle ear masses.

Variant 3: Acquired sensorineural hearing loss. Initial imaging.

CT Temporal Bone

CT of the temporal bone is insensitive in detection of soft-tissue abnormalities that commonly cause sensorineural hearing loss. Small size and proximity to the dense bone of inner ear structures and IAC also precludes visualization of intralabyrinthine or intracanalicular enhancement following IV contrast administration. It may demonstrate labyrinthine ossification [16] resulting from prior infection or give an indirect clue to presence of a vestibular schwannoma in the form of bony remodeling of the IAC. In post-traumatic sensorineural hearing loss, CT can demonstrate fractures extending across the otic capsule [2,13].

CT Head

Contrast-enhanced head CT is a less-sensitive imaging modality to detect tumors, such as vestibular schwannomas [17], or assess the IAC, cerebellopontine angle cisterns, or the brainstem compared to MRI.

CTA Head

There is no evidence to support use of CTA in the initial workup of patients presenting with isolated sensorineural hearing loss.

MRI Head and Internal Auditory Canal

Imaging evaluation of patients presenting with sensorineural hearing loss involves detailed assessment of the cochlear contents, vestibulocochlear nerve, and auditory pathways. MRI is the imaging modality of choice for evaluating these soft-tissue structures [2,5,18-20]. MRI can demonstrate signal alterations induced by inflammation or hemorrhage within the cochlear contents, identify neoplasms within the cochlear labyrinth or IAC, assess the size of vestibular aqueducts, and visualize abnormalities affecting the brain parenchyma along the auditory pathways [21-25]. Although differential considerations may vary based upon sudden, fluctuating, or progressive nature of sensorineural hearing loss, MRI remains the imaging modality of choice for all these subcategories. MRI should be done using dedicated IAC protocol using thin sections across the IAC and the inner ear. These protocols include evaluation of the brainstem and thalami. Given the extreme rarity of cortical deafness, there is no strong evidence to recommend routine assessment of the entire brain parenchyma in addition to the MRI IAC protocol in patients presenting with isolated sensorineural hearing loss [26,27]. High-resolution 3-D T2-weighted images providing submillimeter assessment of fluid-filled inner ear structures and the IAC are highly sensitive for detection of diseases presenting with sensorineural hearing loss [27,28]. Visualization of inflammatory changes (eg, labyrinthitis, neuritis) as well as neoplasms, such as vestibular schwannomas, can be facilitated by administration of IV contrast [29,30]. However, there is insufficient evidence to prove incremental benefit of contrast administration beyond an MRI IAC protocol performed without IV contrast [27,28].

MRA Head

There is no evidence to support use of MRA in the initial workup of patients presenting with isolated sensorineural hearing loss.

MR Venography Head

There is no evidence to support use of MRV in the initial workup of patients presenting with isolated sensorineural hearing loss.

Variant 4: Mixed conductive and sensorineural hearing loss. Initial imaging.

CT Temporal Bone

CT scan of the temporal bones can delineate changes of otospongiosis, a common cause of mixed conductive and sensorineural hearing loss. In some patients with clinical suspicion of otospongiosis, it may suggest alternate diagnoses to explain hearing loss. [31,32]. Administration of IV contrast is usually not beneficial for assessment of temporal bone.

CT Head

Relative to MRI, CT head is much less sensitive in detecting or excluding retrocochlear pathology to account for the sensorineural component of the hearing loss [17].

CTA Head

There is no evidence to support use of CTA in the initial workup of patients presenting with mixed hearing loss.

MRI Head and Internal Auditory Canal

MRI obtained using IAC protocol can be helpful in looking for any retrocochlear pathology responsible for a sensorineural component of the hearing loss. In case IV contrast is administered, punctate enhancement can be seen within the bony otic capsule in the presence of otospongiosis [2].

MRA Head

There is no evidence to support use of MRA in the initial workup of patients presenting with mixed hearing loss.

MR Venography Head

There is no evidence to support use of MRV in the initial workup of patients presenting with mixed hearing loss.

Variant 5: Congenital hearing loss or total deafness or cochlear implant candidate. Surgical planning.

CT Temporal Bone

High-spatial resolution provided by CT of the temporal bone is valuable prior to cochlear implantation surgery in patients with profound hearing loss. It can provide preoperative delineation of underlying cochlear malformation in patients with congenital hearing loss, detect changes of otospongiosis, suggest round window occlusion, identify labyrinthitis ossificans, congenital bony fusion of the ossicles, and alert the surgeon regarding underlying otomastoiditis or variant anatomy (such as that of the facial nerve) [33,34]. It can also delineate the size of cochlear and vestibular aqueducts, alerting the surgeon for possibility of intraoperative cerebrospinal fluid gusher [24,35,36].

CT Head

High-spatial resolution of CT head is insufficient in providing anatomic details of temporal bone needed for surgical planning prior to cochlear implantation. Accordingly, there is no evidence to support routine use of CT head for this indication.

CTA Head

There is no evidence to support routine use of CTA for surgical planning prior to cochlear implantation in patients with deafness.

MRI Head and Internal Auditory Canal

MRI may provide a complementary role to temporal bone CT in preoperative assessment of patients prior to cochlear implantation. Exquisite details of inner ear structures visible on high-resolution T2-weighted images can help in detecting abnormalities, such as cochlear malformations or cochlear nerve deficiency, that directly impact surgical approach [37,38]. In addition, MRI may reveal unexpected soft-tissue abnormalities, such as vestibular schwannomas that may impact the planned surgery [39].

MRA Head

There is no evidence to support routine use of MRA for surgical planning prior to cochlear implantation in patients with deafness.

MR Venography Head

There is no evidence to support routine use of MRV for surgical planning prior to cochlear implantation in patients with deafness.

Variant 6: Episodic vertigo with or without associated hearing loss or aural fullness (peripheral vertigo). Initial imaging.

CT Temporal Bone

CT of the temporal bone provides excellent delineation of the bony labyrinth and is helpful in detecting a number of pathologies resulting in peripheral vertigo. It is highly sensitive in detecting temporal bone fractures in patients with post-traumatic vertigo, assessing for superior semicircular canal dehiscence in patients with vertigo provoked by loud noises, and diagnosing erosions in the bony labyrinth from inflammatory or iatrogenic causes [5,8,9,14].

CT Head

CT head provides insufficient details of the inner ear to be useful in patients with peripheral vertigo. Accordingly, diagnostic yield of CT head in patients presenting with vertigo is low [40].

CTA Head and Neck

There is no evidence to support use of CTA in patients presenting with peripheral causes of vertigo. In patients with episodic vertigo that cannot be confidently categorized as peripheral, CTA can be used to detect underlying vertebrobasilar insufficiency [41].

MRI Head and Internal Auditory Canal

Based on clinical assessment, peripheral vertigo in many patients is presumed to be secondary to benign processes such as benign paroxysmal positional vertigo or Meniere disease, and these patients are often managed successfully without imaging [3-5]. High-resolution T2-weighted images are capable of delineating endolymphatic sac, and delayed 3-D FLAIR images can demonstrate hydrops associated with Meniere disease following IV or intratympanic contrast administration as contrast accumulates in perilymphatic but not endolymphatic space. However, the role of such studies in management of these patients is still not clearly established [42-53]. IV contrast can be helpful in showing enhancement of vestibule or semicircular canals in patients with labyrinthitis. MRI of the brain can be used to detect rare but significant central causes of vertigo in cases where distinction between peripheral and central categories is not clinically evident [54].

MRA Head and Neck

There is no evidence to support use of MRA in patients presenting with peripheral causes of vertigo. In patients with episodic vertigo that cannot be confidently categorized as peripheral, MRA without and with IV contrast can be used to detect underlying vertebrobasilar insufficiency [55].

MR Venography Head

There is no evidence to support use of MRV in the initial workup of patients presenting with vertigo; however, in patients who may have vertigo as a symptom of pseudotumor cerebri, MRV may show narrowing of the transverse sinuses.

Variant 7: Persistent vertigo with or without neurological symptoms (central vertigo). Initial imaging.

CT Temporal Bone

CT of the temporal bone is not useful in looking for central causes of vertigo.

CT Head

Head CT without or with IV contrast may be used to look for central causes of dizziness, albeit with lesser sensitivity than MRI [40,54]. IV contrast may help in either detection or characterization of various neoplastic or inflammatory disease processes affecting the central nervous system. In patients presenting to the emergency department with acute onset of symptoms, CT may demonstrate intracranial hemorrhage as a rare central cause of dizziness [56].

CTA Head and Neck

In patients suspected of vertebrobasilar insufficiency as a cause of episodic vertigo, CTA can be used to detect vascular stenosis or occlusion [41].

MRI Head and Internal Auditory Canal

MRI is the modality of choice in evaluation of the brain in patients suspected to have central cause for vertigo. It can detect posterior fossa neoplasms, Chiari malformation, posterior fossa infarcts, and demyelinating lesions that may result in dizziness or vertigo [3,54,56-58]. Contrast administration can be helpful in detection or characterization of such lesions [58]. Compared to CT, MRI has a much higher sensitivity of detecting acute infarcts in patients with dizziness [40]. It should be noted that infarcts causing isolated vestibular symptoms are usually small, and normal initial MRI does not entirely exclude brain infarction as a cause for vertigo [59].

MRA Head and Neck

In patients suspected of vertebrobasilar insufficiency as a cause of episodic vertigo, MRA can be used to detect vascular stenosis or occlusion [55].

MR Venography Head

There is no evidence to support use of MRV in the initial workup of patients presenting with isolated vertigo.

Summary of Recommendations

- **Variant 1:** CT temporal bone without IV contrast is the first-line imaging modality in patients presenting with acquired conductive hearing loss without any mass lesion seen within the middle ear cavity.

- **Variation 2:** For presurgical planning of patients with acquired conductive hearing loss secondary to cholesteatoma or neoplasm with suspected intracranial or inner ear extension, CT temporal bone without IV contrast or MRI head and internal auditory canal without and with IV contrast is recommended. It may be necessary to do both examinations for complete presurgical evaluation.
- **Variation 3:** MRI head and internal auditory canal without IV contrast, or without and with IV contrast, is recommended for evaluating patients with acquired sensorineural hearing loss.
- **Variation 4:** Mixed conductive and sensorineural hearing loss can be evaluated with either CT temporal bone without IV contrast, or MRI head and internal auditory canal without IV contrast, or without and with IV contrast.
- **Variation 5:** Congenital hearing loss, total deafness, or cochlear implant is best assessed with either CT temporal bone without IV contrast, or MRI head and internal auditory canal without IV contrast, or without and with IV contrast.
- **Variation 6:** Episodic vertigo with or without associated hearing loss or aural fullness (peripheral vertigo) can best be assessed with either CT temporal bone without IV contrast, MRI head and internal auditory canal without IV contrast, or without and with IV contrast.
- **Variation 7:** Evaluation of persistent vertigo with or without neurological symptoms (central vertigo) can best be assessed initially with MRI head and internal auditory canal without IV contrast, or without and with IV contrast.

Summary of Evidence

Of the 60 references cited in the *ACR Appropriateness Criteria® Hearing Loss and/or Vertigo* document, 1 is categorized as a therapeutic reference of good-quality that may have design limitations. Additionally, 59 references are categorized as diagnostic references including 1 well-designed study, 11 good-quality studies, and 24 quality studies that may have design limitations. There are 23 references that may not be useful as primary evidence.

The 60 references cited in the *ACR Appropriateness Criteria® Hearing Loss and/or Vertigo* document were published from 1978 to 2017.

Although there are references that report on studies with design limitations, 13 well-designed or good-quality studies provide good evidence.

Appropriateness Category Names and Definitions

Appropriateness Category Name	Appropriateness Rating	Appropriateness Category Definition
Usually Appropriate	7, 8, or 9	The imaging procedure or treatment is indicated in the specified clinical scenarios at a favorable risk-benefit ratio for patients.
May Be Appropriate	4, 5, or 6	The imaging procedure or treatment may be indicated in the specified clinical scenarios as an alternative to imaging procedures or treatments with a more favorable risk-benefit ratio, or the risk-benefit ratio for patients is equivocal.
May Be Appropriate (Disagreement)	5	The individual ratings are too dispersed from the panel median. The different label provides transparency regarding the panel’s recommendation. “May be appropriate” is the rating category and a rating of 5 is assigned.
Usually Not Appropriate	1, 2, or 3	The imaging procedure or treatment is unlikely to be indicated in the specified clinical scenarios, or the risk-benefit ratio for patients is likely to be unfavorable.

Relative Radiation Level Information

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Patients in the pediatric age group are at inherently higher risk from exposure, because of both organ sensitivity and longer life expectancy (relevant to the long latency that appears to accompany radiation exposure). For these reasons, the RRL dose estimate ranges for pediatric examinations are lower as compared with those specified for adults (see Table below). Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® [Radiation Dose Assessment Introduction](#) document [60].

Relative Radiation Level Designations		
Relative Radiation Level*	Adult Effective Dose Estimate Range	Pediatric Effective Dose Estimate Range
O	0 mSv	0 mSv
⊕	<0.1 mSv	<0.03 mSv
⊕⊕	0.1-1 mSv	0.03-0.3 mSv
⊕⊕⊕	1-10 mSv	0.3-3 mSv
⊕⊕⊕⊕	10-30 mSv	3-10 mSv
⊕⊕⊕⊕⊕	30-100 mSv	10-30 mSv

*RRL assignments for some of the examinations cannot be made, because the actual patient doses in these procedures vary as a function of a number of factors (eg, region of the body exposed to ionizing radiation, the imaging guidance that is used). The RRLs for these examinations are designated as “Varies”.

Supporting Documents

For additional information on the Appropriateness Criteria methodology and other supporting documents go to www.acr.org/ac.

References

1. Curtin HD. Imaging of Conductive Hearing Loss With a Normal Tympanic Membrane. *AJR Am J Roentgenol* 2016;206:49-56.
2. Shah LM, Wiggins RH, 3rd. Imaging of hearing loss. *Neuroimaging Clin N Am* 2009;19:287-306.
3. Connor SE, Sriskandan N. Imaging of dizziness. *Clin Radiol* 2014;69:111-22.
4. Macleod D, McAuley D. Vertigo: clinical assessment and diagnosis. *Br J Hosp Med (Lond)* 2008;69:330-4.
5. Newman-Toker DE, Della Santina CC, Blitz AM. Vertigo and hearing loss. *Handb Clin Neurol* 2016;136:905-21.
6. Bakhit M, Heidarian A, Ehsani S, Delphi M, Latifi SM. Clinical assessment of dizzy patients: the necessity and role of diagnostic tests. *Glob J Health Sci* 2014;6:194-9.
7. Kutz JW, Jr. The dizzy patient. *Med Clin North Am* 2010;94:989-1002.
8. Belden CJ, Weg N, Minor LB, Zinreich SJ. CT evaluation of bone dehiscence of the superior semicircular canal as a cause of sound- and/or pressure-induced vertigo. *Radiology* 2003;226:337-43.
9. Branstetter BF, 4th, Harrigal C, Escott EJ, Hirsch BE. Superior semicircular canal dehiscence: oblique reformatted CT images for diagnosis. *Radiology* 2006;238:938-42.
10. Lee YH, Rivas-Rodriguez F, Song JJ, Yang KS, Mukherji SK. The prevalence of superior semicircular canal dehiscence in conductive and mixed hearing loss in the absence of other pathology using submillimetric temporal bone computed tomography. *J Comput Assist Tomogr* 2014;38:190-5.
11. Saliba I, Maniakas A, Benamira LZ, Nehme J, Benoit M, Montreuil-Jacques V. Superior canal dehiscence syndrome: clinical manifestations and radiologic correlations. *Eur Arch Otorhinolaryngol* 2014;271:2905-14.
12. Stimmer H, Hamann KF, Zeiter S, Naumann A, Rummeny EJ. Semicircular canal dehiscence in HR multislice computed tomography: distribution, frequency, and clinical relevance. *Eur Arch Otorhinolaryngol* 2012;269:475-80.

13. Eshetu T, Aygun N. Imaging of the temporal bone: a symptom-based approach. *Semin Roentgenol* 2013;48:52-64.
14. Meyer A, Bouchetemple P, Costentin B, Dehesdin D, Lerosey Y, Marie JP. Lateral semicircular canal fistula in cholesteatoma: diagnosis and management. *Eur Arch Otorhinolaryngol* 2016;273:2055-63.
15. Mohan S, Hoeffner E, Bigelow DC, Loevner LA. Applications of magnetic resonance imaging in adult temporal bone disorders. *Magn Reson Imaging Clin N Am* 2012;20:545-72.
16. Braun T, Dirr F, Berghaus A, et al. Prevalence of labyrinthine ossification in CT and MR imaging of patients with acute deafness to severe sensorineural hearing loss. *Int J Audiol* 2013;52:495-9.
17. Kulkarni BSN, Bajwa H, Chandrashekar M, et al. CT- and MRI-based gross target volume comparison in vestibular schwannomas. *Rep Pract Oncol Radiother* 2017;22:201-08.
18. Berrettini S, Seccia V, Fortunato S, et al. Analysis of the 3-dimensional fluid-attenuated inversion-recovery (3D-FLAIR) sequence in idiopathic sudden sensorineural hearing loss. *JAMA Otolaryngol Head Neck Surg* 2013;139:456-64.
19. Chau JK, Cho JJ, Fritz DK. Evidence-based practice: management of adult sensorineural hearing loss. *Otolaryngol Clin North Am* 2012;45:941-58.
20. Cueva RA. Auditory brainstem response versus magnetic resonance imaging for the evaluation of asymmetric sensorineural hearing loss. *Laryngoscope* 2004;114:1686-92.
21. Davidson HC, Harnsberger HR, Lemmerling MM, et al. MR evaluation of vestibulocochlear anomalies associated with large endolymphatic duct and sac. *AJNR Am J Neuroradiol* 1999;20:1435-41.
22. Kwan TL, Tang KW, Pak KK, Cheung JY. Screening for vestibular schwannoma by magnetic resonance imaging: analysis of 1821 patients. *Hong Kong Med J* 2004;10:38-43.
23. Mafee MF. Congenital sensorineural hearing loss and enlarged endolymphatic sac and duct: role of magnetic resonance imaging and computed tomography. *Top Magn Reson Imaging* 2000;11:10-24.
24. Valvassori GE, Clemis JD. The large vestibular aqueduct syndrome. *Laryngoscope* 1978;88:723-8.
25. Weissman JL, Curtin HD, Hirsch BE, Hirsch WL, Jr. High signal from the otic labyrinth on unenhanced magnetic resonance imaging. *AJNR Am J Neuroradiol* 1992;13:1183-7.
26. Gebarski SS, Tucci DL, Telian SA. The cochlear nuclear complex: MR location and abnormalities. *AJNR Am J Neuroradiol* 1993;14:1311-8.
27. Sharma A, Viets R, Parsons MS, Reis M, Chrisinger J, Wippold FJ, 2nd. A two-tiered approach to MRI for hearing loss: incremental cost of a comprehensive MRI over high-resolution T2-weighted imaging. *AJR Am J Roentgenol* 2014;202:136-44.
28. Daniels RL, Swallow C, Shelton C, Davidson HC, Krejci CS, Harnsberger HR. Causes of unilateral sensorineural hearing loss screened by high-resolution fast spin echo magnetic resonance imaging: review of 1,070 consecutive cases. *Am J Otol* 2000;21:173-80.
29. Held P, Fellner C, Fellner F, et al. MRI of inner ear and facial nerve pathology using 3D MP-RAGE and 3D CISS sequences. *Br J Radiol* 1997;70:558-66.
30. Zealley IA, Cooper RC, Clifford KM, et al. MRI screening for acoustic neuroma: a comparison of fast spin echo and contrast enhanced imaging in 1233 patients. *Br J Radiol* 2000;73:242-7.
31. Dudau C, Salim F, Jiang D, Connor SE. Diagnostic efficacy and therapeutic impact of computed tomography in the evaluation of clinically suspected otosclerosis. *Eur Radiol* 2017;27:1195-201.
32. Quesnel AM, Moonis G, Appel J, et al. Correlation of computed tomography with histopathology in otosclerosis. *Otol Neurotol* 2013;34:22-8.
33. Jeong SW, Kim LS. A new classification of cochleovestibular malformations and implications for predicting speech perception ability after cochlear implantation. *Audiol Neurootol* 2015;20:90-101.
34. Young JY, Ryan ME, Young NM. Preoperative imaging of sensorineural hearing loss in pediatric candidates for cochlear implantation. *Radiographics* 2014;34:E133-49.
35. El-Badry MM, Osman NM, Mohamed HM, Rafaat FM. Evaluation of the radiological criteria to diagnose large vestibular aqueduct syndrome. *Int J Pediatr Otorhinolaryngol* 2016;81:84-91.
36. Kim BG, Sim NS, Kim SH, Kim UK, Kim S, Choi JY. Enlarged cochlear aqueducts: a potential route for CSF gushers in patients with enlarged vestibular aqueducts. *Otol Neurotol* 2013;34:1660-5.
37. Glastonbury CM, Davidson HC, Harnsberger HR, Butler J, Kertesz TR, Shelton C. Imaging findings of cochlear nerve deficiency. *AJNR Am J Neuroradiol* 2002;23:635-43.
38. Parry DA, Booth T, Roland PS. Advantages of magnetic resonance imaging over computed tomography in preoperative evaluation of pediatric cochlear implant candidates. *Otol Neurotol* 2005;26:976-82.

39. Jiang ZY, Odiase E, Isaacson B, Roland PS, Kutz JW, Jr. Utility of MRIs in adult cochlear implant evaluations. *Otol Neurotol* 2014;35:1533-5.
40. Lawhn-Heath C, Buckle C, Christoforidis G, Straus C. Utility of head CT in the evaluation of vertigo/dizziness in the emergency department. *Emerg Radiol* 2013;20:45-9.
41. Pasaoglu L. Vertebrobasilar system computed tomographic angiography in central vertigo. *Medicine (Baltimore)* 2017;96:e6297.
42. Fukuoka H, Takumi Y, Tsukada K, et al. Comparison of the diagnostic value of 3 T MRI after intratympanic injection of GBCA, electrocochleography, and the glycerol test in patients with Meniere's disease. *Acta Otolaryngol* 2012;132:141-5.
43. Grieve SM, Obholzer R, Malitz N, Gibson WP, Parker GD. Imaging of endolymphatic hydrops in Meniere's disease at 1.5 T using phase-sensitive inversion recovery: (1) demonstration of feasibility and (2) overcoming the limitations of variable gadolinium absorption. *Eur J Radiol* 2012;81:331-8.
44. Hagiwara M, Roland JT, Jr., Wu X, et al. Identification of endolymphatic hydrops in Meniere's disease utilizing delayed postcontrast 3D FLAIR and fused 3D FLAIR and CISS color maps. *Otol Neurotol* 2014;35:e337-42.
45. Hornibrook J, Flook E, Greig S, et al. MRI Inner Ear Imaging and Tone Burst Electrocochleography in the Diagnosis of Meniere's Disease. *Otol Neurotol* 2015;36:1109-14.
46. Karatas A, Kocak A, Cebi IT, Salviz M. Comparison of Endolymphatic Duct Dimensions and Jugular Bulb Abnormalities Between Meniere Disease and a Normal Population. *J Craniofac Surg* 2016;27:e424-6.
47. Le CH, Truong AQ, Diaz RC. Novel techniques for the diagnosis of Meniere's disease. *Curr Opin Otolaryngol Head Neck Surg* 2013;21:492-6.
48. Liu F, Huang W, Meng X, Wang Z, Liu X, Chen Q. Comparison of noninvasive evaluation of endolymphatic hydrops in Meniere's disease and endolymphatic space in healthy volunteers using magnetic resonance imaging. *Acta Otolaryngol* 2012;132:234-40.
49. Liu Y, Jia H, Shi J, et al. Endolymphatic hydrops detected by 3-dimensional fluid-attenuated inversion recovery MRI following intratympanic injection of gadolinium in the asymptomatic contralateral ears of patients with unilateral Meniere's disease. *Med Sci Monit* 2015;21:701-7.
50. Naganawa S, Yamazaki M, Kawai H, et al. MR imaging of Meniere's disease after combined intratympanic and intravenous injection of gadolinium using HYDROPS2. *Magn Reson Med Sci* 2014;13:133-7.
51. Naganawa S, Yamazaki M, Kawai H, Bokura K, Sone M, Nakashima T. Imaging of Meniere's disease after intravenous administration of single-dose gadodiamide: utility of multiplication of MR cisternography and HYDROPS image. *Magn Reson Med Sci* 2013;12:63-8.
52. Sepahdari AR, Ishiyama G, Vorasubin N, Peng KA, Linetsky M, Ishiyama A. Delayed intravenous contrast-enhanced 3D FLAIR MRI in Meniere's disease: correlation of quantitative measures of endolymphatic hydrops with hearing. *Clin Imaging* 2015;39:26-31.
53. Wu Q, Dai C, Zhao M, Sha Y. The correlation between symptoms of definite Meniere's disease and endolymphatic hydrops visualized by magnetic resonance imaging. *Laryngoscope* 2016;126:974-9.
54. Kabra R, Robbie H, Connor SE. Diagnostic yield and impact of MRI for acute ischaemic stroke in patients presenting with dizziness and vertigo. *Clin Radiol* 2015;70:736-42.
55. Arai M, Higuchi A, Umekawa J, Mochimatsu Y, Itoh K. [The efficiency of magnetic resonance angiography (MRA) in the diagnosis and vertigo--prediction of vertebrobasilar insufficiency (VBI) and atherosclerosis]. *Nihon Jibiinkoka Gakkai Kaiho* 1999;102:925-31.
56. Doijiri R, Uno H, Miyashita K, Ihara M, Nagatsuka K. How Commonly Is Stroke Found in Patients with Isolated Vertigo or Dizziness Attack? *J Stroke Cerebrovasc Dis* 2016;25:2549-52.
57. Leker RR, Hur TB, Gomori JM, Paniri R, Eichel R, Cohen JE. Incidence of DWI-positive stroke in patients with vertigo of unclear etiology, preliminary results. *Neurol Res* 2013;35:123-6.
58. Schick B, Brors D, Koch O, Schafers M, Kahle G. Magnetic resonance imaging in patients with sudden hearing loss, tinnitus and vertigo. *Otol Neurotol* 2001;22:808-12.
59. Saber Tehrani AS, Kattah JC, Mantokoudis G, et al. Small strokes causing severe vertigo: frequency of false-negative MRIs and nonlacunar mechanisms. *Neurology* 2014;83:169-73.
60. American College of Radiology. ACR Appropriateness Criteria® Radiation Dose Assessment Introduction. Available at: <https://www.acr.org/-/media/ACR/Files/Appropriateness-Criteria/RadiationDoseAssessmentIntro.pdf>. Accessed March 30, 2018.

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.