

**American College of Radiology  
ACR Appropriateness Criteria®  
Cranial Neuropathy**

**Variant 1:                      Anosmia and abnormalities of the sense of smell. (Olfactory nerve, CN I.)**

<b>Radiologic Procedure</b>	<b>Rating</b>	<b>Comments</b>	<b>RRL*</b>
MRI head without and with IV contrast	8	This procedure is performed in conjunction with MRI of the orbit, face, and neck.	O
MRI orbit face neck without and with IV contrast	8	This procedure is performed in conjunction with MRI of the head.	O
MRI head without IV contrast	6		O
MRI orbit face neck without IV contrast	6		O
CT maxillofacial with IV contrast	6		☼☼
CT head with IV contrast	5		☼☼☼
CT head without IV contrast	5		☼☼☼
CT head without and with IV contrast	5		☼☼☼
CT maxillofacial without IV contrast	5		☼☼
CT maxillofacial without and with IV contrast	4		☼☼☼
FDG-PET/CT whole body	2		☼☼☼☼
<b><u>Rating Scale:</u> 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate</b>			<b>*Relative Radiation Level</b>

**Variant 2:****Weakness or paralysis of the mastication muscles. Sensory abnormalities of the head and neck. Trigeminal neuralgia. (Trigeminal nerve, CN V.)**

<b>Radiologic Procedure</b>	<b>Rating</b>	<b>Comments</b>	<b>RRL*</b>
MRI head without and with IV contrast	8	This procedure is performed in conjunction with MRI of the orbit, face, and neck.	O
MRI orbit face neck without and with IV contrast	8	This procedure is performed in conjunction with MRI of the head.	O
MRI head without IV contrast	6		O
MRI orbit face neck without IV contrast	6		O
MRA head without IV contrast	6		O
CTA head with IV contrast	5		⊕⊕⊕⊕
CT head with IV contrast	5		⊕⊕⊕⊕
CT maxillofacial with IV contrast	5		⊕⊕
CT maxillofacial without IV contrast	5		⊕⊕
CT head without IV contrast	4		⊕⊕⊕⊕
CT head without and with IV contrast	4		⊕⊕⊕⊕
CT neck with IV contrast	4		⊕⊕⊕⊕
CT neck without IV contrast	4	Contrast-enhanced imaging is preferred.	⊕⊕⊕⊕
CT maxillofacial without and with IV contrast	4		⊕⊕⊕⊕
CT neck without and with IV contrast	3		⊕⊕⊕⊕
FDG-PET/CT whole body	2		⊕⊕⊕⊕⊕
US neck	2		O
<b>Rating Scale:</b> 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			<b>*Relative Radiation Level</b>

**Variant 3:****Weakness or paralysis of facial expression. Hemifacial spasm. Bell palsy. (Facial nerve, CN VII.)**

<b>Radiologic Procedure</b>	<b>Rating</b>	<b>Comments</b>	<b>RRL*</b>
MRI orbit face neck without and with IV contrast	9	This procedure is performed in conjunction with MRI of the head.	O
MRI head without and with IV contrast	8	This procedure is performed in conjunction with MRI of the orbit, face, and neck.	O
MRI head without IV contrast	5		O
MRI orbit face neck without IV contrast	5		O
CT head with IV contrast	5		☼☼☼
CT head without IV contrast	5		☼☼☼
CT head without and with IV contrast	4		☼☼☼
CT neck with IV contrast	4		☼☼☼
CT neck without IV contrast	3		☼☼☼
CT neck without and with IV contrast	3		☼☼☼
FDG-PET/CT whole body	2		☼☼☼☼
US neck	2		O
<b>Rating Scale:</b> 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			<b>*Relative Radiation Level</b>

**Variant 4:****Palate weakness. Oropharyngeal pain. (Glossopharyngeal nerve, CN IX.)**

<b>Radiologic Procedure</b>	<b>Rating</b>	<b>Comments</b>	<b>RRL*</b>
MRI head without and with IV contrast	8	This procedure is performed in conjunction with MRI of the orbit, face, and neck.	O
MRI orbit face neck without and with IV contrast	8	This procedure is performed in conjunction with MRI of the head.	O
MRI head without IV contrast	6		O
MRI orbit face neck without IV contrast	6		O
CT neck with IV contrast	6		☼☼☼
CT head with IV contrast	5		☼☼☼
CT head without IV contrast	5	Contrast-enhanced imaging is preferred.	☼☼☼
CT neck without IV contrast	5	Contrast-enhanced imaging is preferred.	☼☼☼
CT head without and with IV contrast	4		☼☼☼
CT neck without and with IV contrast	4		☼☼☼
FDG-PET/CT whole body	2		☼☼☼☼
US neck	2		O
<b>Rating Scale:</b> 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			<b>*Relative Radiation Level</b>

**Variant 5:****Vocal cord paralysis. (Vagal nerve, CN X.)**

<b>Radiologic Procedure</b>	<b>Rating</b>	<b>Comments</b>	<b>RRL*</b>
MRI orbit face neck without and with IV contrast	8	This procedure is performed in conjunction with MRI of the head. CT of the neck is an alternative examination and in some instances can be complementary.	O
CT neck with IV contrast	8	MRI of the orbit, face, and neck can be an alternative examination and in some instances can be complementary.	☼☼☼
MRI head without and with IV contrast	7	This procedure is performed in conjunction with MRI of the orbit, face, and neck. CT of the neck can be useful to assess the extracranial course of CN X.	O
MRI head without IV contrast	6		O
MRI orbit face neck without IV contrast	6		O
CT chest with IV contrast	6		☼☼☼
MRI chest without and with IV contrast	5		O
CT head with IV contrast	5		☼☼☼
CT neck without IV contrast	5	Contrast-enhanced imaging is preferred.	☼☼☼
CT chest without IV contrast	5		☼☼☼
MRI chest without IV contrast	4		O
CT head without IV contrast	4		☼☼☼
CT head without and with IV contrast	4		☼☼☼
CT neck without and with IV contrast	4		☼☼☼
X-ray chest	4		☼
FDG-PET/CT whole body	4	This procedure is not a first-line examination.	☼☼☼☼
US neck	4		O
CT chest without and with IV contrast	3		☼☼☼
<b>Rating Scale:</b> 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			<b>*Relative Radiation Level</b>

**Variant 6:****Weakness or paralysis of the sternocleidomastoid and trapezius muscles. (Accessory nerve, CN XI.)**

<b>Radiologic Procedure</b>	<b>Rating</b>	<b>Comments</b>	<b>RRL*</b>
MRI head without and with IV contrast	8	This procedure is performed in conjunction with MRI of the orbit, face, and neck.	O
MRI orbit face neck without and with IV contrast	8	This procedure is performed in conjunction with MRI of the head. CT neck imaging can be complementary.	O
MRI orbit face neck without IV contrast	7	Contrast-enhanced imaging is preferred.	O
CT neck with IV contrast	7	MRI of the orbit, face, and neck can be an alternative examination and in some instances can be complementary.	☼☼☼
MRI head without IV contrast	6		O
CT head with IV contrast	6		☼☼☼
CT head without IV contrast	5		☼☼☼
CT head without and with IV contrast	5		☼☼☼
CT neck without IV contrast	5		☼☼☼
CT neck without and with IV contrast	5	The panel did not agree on a recommendation.	☼☼☼
FDG-PET/CT whole body	3		☼☼☼☼
US neck	2		O
<b>Rating Scale:</b> 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			<b>*Relative Radiation Level</b>

**Variant 7:****Weakness or paralysis of the tongue. (Hypoglossal nerve, CN XII.)**

<b>Radiologic Procedure</b>	<b>Rating</b>	<b>Comments</b>	<b>RRL*</b>
MRI head without and with IV contrast	8	This procedure is performed in conjunction with MRI of the orbit, face, and neck.	O
MRI orbit face neck without and with IV contrast	8	This procedure is performed in conjunction with MRI of the head.	O
CT neck with IV contrast	7	MRI of the orbit, face, and neck is preferred, but in some instances CT neck can be complementary.	☼☼☼
MRI head without IV contrast	6		O
MRI orbit face neck without IV contrast	6		O
CT head with IV contrast	5		☼☼☼
CT head without IV contrast	5		☼☼☼
CT head without and with IV contrast	4		☼☼☼
CT neck without IV contrast	4		☼☼☼
CT neck without and with IV contrast	4		☼☼☼
FDG-PET/CT whole body	2		☼☼☼☼
US neck	2		O
<b>Rating Scale:</b> 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			<b>*Relative Radiation Level</b>

**Variant 8:****Perineural spread of tumor. (Most commonly trigeminal nerve [CN V], facial nerve [CN VII].)**

<b>Radiologic Procedure</b>	<b>Rating</b>	<b>Comments</b>	<b>RRL*</b>
MRI head without and with IV contrast	9	This procedure is performed in conjunction with MRI of the orbit, face, and neck.	O
MRI orbit face neck without and with IV contrast	9	This procedure is performed in conjunction with MRI of the head.	O
MRI orbit face neck without IV contrast	7	Addition of contrast-enhanced imaging is preferred.	O
CT neck with IV contrast	6		☼☼☼
MRI head without IV contrast	5		O
CT head with IV contrast	5		☼☼☼
CT head without IV contrast	5		☼☼☼
CT neck without IV contrast	5		☼☼☼
CT head without and with IV contrast	4		☼☼☼
CT neck without and with IV contrast	4		☼☼☼
FDG-PET/CT whole body	4	This procedure is not a first-line examination.	☼☼☼☼
US neck	2		O
<b>Rating Scale:</b> 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			<b>*Relative Radiation Level</b>



# CRANIAL NEUROPATHY

Expert Panel on Neurologic Imaging: Bruno Policeni, MD<sup>a</sup>; Amanda S. Corey, MD<sup>b</sup>; Judah Burns, MD<sup>c</sup>; David B. Conley, MD<sup>d</sup>; R. Webster Crowley, MD<sup>e</sup>; H. Benjamin Harvey, MD, JD<sup>f</sup>; Jenny Hoang, MBBS<sup>g</sup>; Christopher H. Hunt, MD<sup>h</sup>; Bharathi D. Jagadeesan, MD<sup>i</sup>; Amy F. Juliano, MD<sup>j</sup>; Tabassum A. Kennedy, MD<sup>k</sup>; Gul Moonis, MD<sup>l</sup>; Jeffrey S. Pannell, MD<sup>m</sup>; Nandini D. Patel, MD<sup>n</sup>; Joel S. Perlmutter, MD<sup>o</sup>; Joshua M. Rosenow, MD<sup>p</sup>; Jason W. Schroeder, MD<sup>q\*</sup>; Matthew T. Whitehead, MD<sup>r</sup>; Rebecca S. Cornelius, MD.<sup>s</sup>

## Summary of Literature Review

### **Introduction/Background**

The cranial nerves arise from nuclei within the brain and brain stem and supply sensory and motor innervation to the head and neck region, whereas the spinal nerves arise from the spinal cord and supply the rest of the body. As a group, the cranial nerves have both sensory and motor components similar to those of the spinal nerves. Individually the cranial nerves may be purely sensory or purely motor or a mixture of both sensory and motor. Functions of the cranial nerves may be divided into three sensory and three motor categories. The sensory group includes visceral sensory, which supplies sensory input from the internal organs; general sensory, which supplies tactile, pain, temperature, and other sensations; and special sensory, which includes the special senses of smell, vision, taste, hearing, and balance. Of the three motor functions, somatic motor innervates muscles that develop from the body somites; branchial motor innervates muscles derived from the branchial arches; and visceral motor innervates the viscera, glands, and smooth muscle [1-3].

Cranial nerves emerge in an orderly fashion from the rostral portion of the embryologically developing neural tube, which will subsequently mature to form the brain and brain stem. Anatomically, the 12 pairs of cranial nerves are designated by numbers and are organized most rostral to most caudal in descending order. The cranial nerves include the olfactory (cranial nerve [CN] I), optic (CN II), oculomotor (CN III), trochlear (CN IV), trigeminal (CN V), abducens (CN VI), facial (CN VII), vestibulocochlear (CN VIII), glossopharyngeal (CN IX), vagus (CN X), spinal accessory (CN XI), and hypoglossal (CN XII) nerves. The olfactory (CN I) and optic (CN II) nerves are actually tracts formed from the telencephalon and diencephalon, respectively, and are not considered true nerves [1]. The optic (CN II), oculomotor (CN III), trochlear (CN IV), and abducens (CN VI) nerves are considered functionally to be part of the visual and extraocular motor system and have been discussed in the ACR Appropriateness Criteria<sup>®</sup> “[Orbits, Vision and Visual Loss](#)” [4]. Also, the vestibulocochlear nerve (CN VIII) has been reviewed in the ACR Appropriateness Criteria<sup>®</sup> “[Hearing Loss and/or Vertigo](#)” [5]. Therefore, this discussion will focus on CN I, CN V, CN VII, CN IX, CN X, CN XI, and CN XII.

In approaching cranial neuropathy, several concepts should be emphasized:

1. Because of the complex anatomic structures within the brain and brain stem and because the cranial nerves may take long, circuitous routes to their destinations, a detailed knowledge of cranial nerve anatomy is essential for proper clinical localization of potential lesions and for appropriate application of specific imaging protocols.
2. Because some individual nerve fibers, such as the autonomic nerves, may travel with several different cranial nerves from their nuclei of origin to their ultimate destinations, loss of a specific function may indicate involvement of potentially more than one cranial nerve.

---

<sup>a</sup>Principal Author and Panel Vice-chair, University of Iowa Hospitals and Clinics, Iowa City, Iowa. <sup>b</sup>Panel Chair, Emory University, Atlanta, Georgia. <sup>c</sup>Montefiore Medical Center, Bronx, New York. <sup>d</sup>Northwestern University Feinberg School of Medicine, Chicago, Illinois; American Academy of Otolaryngology-Head and Neck Surgery. <sup>e</sup>Rush University Medical Center, Chicago, Illinois; neurosurgical consultant. <sup>f</sup>Massachusetts General Hospital, Boston, Massachusetts. <sup>g</sup>Duke University Medical Center, Durham, North Carolina. <sup>h</sup>Mayo Clinic, Rochester, Minnesota. <sup>i</sup>University of Minnesota, Minneapolis, Minnesota. <sup>j</sup>Massachusetts Eye and Ear Infirmary, Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts. <sup>k</sup>University of Wisconsin Hospital and Clinic, Madison, Wisconsin. <sup>l</sup>Columbia University Medical Center, New York, New York. <sup>m</sup>University of California San Diego Medical Center, San Diego, California. <sup>n</sup>Fairfax Radiology Consultants PC, Fairfax, Virginia. <sup>o</sup>Washington University School of Medicine, Saint Louis, Missouri; American Academy of Neurology. <sup>p</sup>Northwestern University Feinberg School of Medicine, Chicago, Illinois; neurosurgical consultant. <sup>q</sup>Walter Reed National Military Medical Center, Bethesda, Maryland. <sup>r</sup>Children’s National Medical Center, Washington, District of Columbia. <sup>s</sup>Specialty Chair, University of Cincinnati Medical Center, Cincinnati, Ohio.

The American College of Radiology seeks and encourages collaboration with other organizations on the development of the ACR Appropriateness Criteria through society representation on expert panels. Participation by representatives from collaborating societies on the expert panel does not necessarily imply individual or society endorsement of the final document.

\*The views expressed in this manuscript are those of the author and do not reflect the official policy of the Department of Army/Navy/Air Force, Department of Defense, or United States Government.

Reprint requests to: [publications@acr.org](mailto:publications@acr.org)

3. Because of the close proximity of many cranial nerve nuclei and of many exiting sites of the nerves themselves, some mass lesions may involve multiple cranial nerves.

### **Special Imaging Considerations**

In the evaluation of cranial neuropathy complete evaluation of the nerves from their brain stem nuclei to their “end organs” must be performed. The pathology can be located in the nucleus, cisternae, and skull base segments of the cranial nerves. For CN VII, the lesion can also be located in the parotid. For CN IX, CN X, CN XI, and CN XII, the lesion can also be located in the neck. Evaluation of the upper chest is necessary for complete evaluation of the CN X (recurrent laryngeal) course. This can be accomplished by extending the neck scan into the mid thorax (aortic pulmonary window) or dedicated chest computed tomography (CT). Patients presenting with otalgia may require evaluation of CN V, VII, IX, and X and upper cervical nerves C2 and C3 because any of these nerves may be the source for the otalgia [6]. The use of intravenous contrast is imperative for the evaluation of cranial neuropathy with magnetic resonance imaging (MRI). Neck CT also requires the use of contrast when evaluating pathology affecting the neck. Dual-phase CT before and after administration of contrast is rarely necessary. This should be avoided because of the extra radiation exposure and minimal added benefits.

The primary plane of study for head and neck evaluation of cranial neuropathy is usually the axial plane. Additional orthogonal planes are required depending upon the course of the various nerves. Coronal and sagittal reconstructions are typically performed on CT. To obtain high-resolution orthogonal reconstructions, the axial plane is acquired with thin sections, typically <1 mm. On MRI, orthogonal reconstructions are typically performed on the postcontrast T1-weighted images; however, they can also be obtained on T2-weighted images. Thin-section MRI are required to evaluate the cisternal segment and should be performed.

High-field strength magnets (1.5-3.0T) are preferred to low-field-strength units because of achievable signal to noise ratios, gradient strength, and spatial resolution [7]. A phased-array head coil suffices for most examinations; specialized surface coils may supplement examinations of peripherally located nerves.

Fundamental techniques include T1-weighted, T2-weighted, and enhanced T1-weighted imaging sequences. The unenhanced T1-weighted sequence remains an excellent baseline technique for anatomic evaluation because of the natural contrast provided by neck and skull base fat. Specialized versions of sequences may be available on scanners depending on manufacturer options. For example, various 3-D and heavily T2-weighted sequences—such as constructive interference in steady state, 3-D-balanced fast field echo, 3-D-driven equilibrium radio frequency reset pulse, 3-D fast spin echo, fast imaging using steady-state acquisition, and 3-D fast spin-echo extended echo-train acquisition—may provide excellent spatial resolution of the cisternal segments of some of the cranial nerves, but they must be used judiciously because of potentially misleading artifacts [7-17]. Enhanced fat-suppression T1-weighted techniques may emphasize abnormally enhancing lesions and nerves but may potentially mask subtle pathology if the suppression is nonuniform. Additional sequences, such as diffusion-weighted imaging, may be added to evaluate specific pathologies, such as infarctions, or specific lesions, such as epidermoids, that may affect cranial nerve function. Slice thickness should be calculated for optimal spatial resolution without introducing partial-volume effect. Because cranial nerve examinations tend to be lengthy, strategies such as parallel imaging may improve patient compliance and image quality [7].

### **Discussion of Procedures by Variant**

#### **Variant 1: Anosmia and abnormalities of the sense of smell. (Olfactory nerve, CN I.)**

Abnormalities of the special sense of smell are mediated by the olfactory nerve (CN I) and can be grouped into clinical categories. Quantitative disturbances imply diminished or enhanced sense of smell (anosmia, hyposmia, or hyperosmia). Qualitative disturbances involve distortions of the sense of smell (dysosmia). Discrimination disturbances involve an inability to differentiate among various smells. Hallucinations or delusions in the sense of smell may also occur. The latter may be caused by temporal lobe dysfunction (see the ACR Appropriateness Criteria® [“Seizures and Epilepsy”](#) [18]) or by degenerative or psychiatric disease. Taste, mediated by the facial (CN VII) and glossopharyngeal (CN IX) nerves, may also be affected by pathology involving the olfactory nerve (CN I).

Most patients with olfactory symptoms do not require imaging. Chronic tobacco use, upper respiratory infections, and inflammatory conditions most commonly affect the sense of smell [19]. More serious conditions affecting the olfactory nerve include trauma (the olfactory nerve is the nerve most commonly disrupted by trauma); cribriform plate tumors such as invasive squamous cell carcinomas of the paranasal sinuses, meningiomas, and

esthesioneuroblastomas; inflammatory lesions such as sarcoidosis and granulomatosis with polyangiitis (formerly known as Wegener granulomatosis); and congenital conditions such as cephaloceles and Kallmann syndrome [8,20-22]. Recent investigations have focused on olfactory bulb volume as an indicator of olfactory dysfunction and even a marker for such disorders as early Parkinson disease and depression [20,23-26].

### **MRI and CT**

MRI is the mainstay for examining the olfactory apparatus, although CT remains useful when evaluating fractures, paranasal sinus inflammatory disease, and bony anatomy [20,27,28]. Imaging protocols should cover the major anatomic divisions of the olfactory nerve and pathway, including the olfactory epithelium, which is located in the upper nasal cavity; the olfactory neurons and bulbs, located in the cribriform plate and inferior frontal lobes; and the olfactory pathways, which travel in portions of the temporal and frontal lobes [7].

### **FDG-PET/CT and Other Imaging Modalities**

Efforts using functional MRI, single-photon emission computed tomography, and fluorine-18-2-fluoro-2-deoxy-D-glucose positron emission tomography (FDG-PET) in studying olfactory dysfunction remain largely investigative and are not generally used in routine evaluations [29-33].

### **Variant 2: Weakness or paralysis of the mastication muscles. Sensory abnormalities of the head and neck. Trigeminal neuralgia. (Trigeminal nerve, CN V.)**

The trigeminal nerve (CN V) provides general sensation to large portions of the head and neck and branchial motor innervation to the muscles of mastication [3]. It is the largest cranial nerve and is divided into three main divisions, known as the ophthalmic (V1), maxillary (V2), and mandibular (V3) branches [8]. Symptoms largely depend on the involved segment and division [34]. Abnormalities of the nerve may manifest as sensory disturbances, such as trigeminal neuralgia (*tic douloureux*), facial numbness, or motor abnormalities such as weakness when chewing food.

The trigeminal nerve (CN V) is the nerve of the first branchial arch and may be involved in congenital conditions such as Goldenhar-Gorlin syndrome [34]. Intra-axial and extra-axial processes may affect the brain stem trigeminal nuclei and nerve root entry and exit zones. Conditions localized to the brain stem portion of the trigeminal nerve (CN V) include vascular lesions (such as compressing vascular loops, aneurysms, vertebrobasilar dolichoectasia, and infarctions), inflammatory and infectious conditions (such as meningitis, encephalitis, sarcoidosis, and multiple sclerosis), and tumors (such as gliomas, lymphomas, metastases, and meningiomas) [35-38]. The cisternal portion of the nerve may be especially vulnerable to compression from adjacent vascular loops, causing trigeminal neuralgia [38,39]. Tumors, vascular lesions, and inflammatory processes may also affect the branches of the nerve as they traverse the Meckel cave, the pterygopalatine fossa, the orbit, the skull base, and the masticator space [34,40].

### **MRI and CT**

MRI is the preferred modality for investigating the trigeminal nerve (CN V) [7,34]. CT is very useful for evaluating the skull base and neural foramina. Three-dimensional and heavily T2-weighted MRI sequences and MR angiography and CT angiography are helpful noninvasive methods for reviewing the anatomy of potentially compressing vascular loops [41-56]. Patients may benefit from MRI studies performed in a high-strength magnet (3 T), given the higher anatomic resolution [57,58]. With the growing popularity of radiosurgery, such as Gamma Knife procedures, and radiofrequency thermocoagulation in the treatment of trigeminal neuralgia, both CT and MRI have become indispensable planning and follow-up tools, although imaging may not reliably predict outcome [35,59-73]. Because of the complex branching patterns of the nerve, multiple imaging planes are essential [34].

### **US**

US is not routinely used in the initial evaluation of the trigeminal nerve.

### **FDG-PET/CT**

FDG-PET/CT is not routinely used in the initial evaluation of the trigeminal nerve.

### **Advanced Imaging Modalities**

Advanced imaging applications, such as fractional anisotropy derived from diffusion tensor imaging and virtual endoscopy, are promising future directions in investigating trigeminal neuralgia [74-76].

### **Variant 3: Weakness or paralysis of facial expression. Hemifacial spasm. Bell palsy. (Facial nerve, CN VII.)**

The facial nerve (CN VII) is one of the most complex cranial nerves and contains branchial motor (innervation to the muscles of facial expression), visceral motor (parasympathetic innervation to most of the glands of the head), general sensory (surface innervations to a small portion of the external ear and tympanic membrane), and special sensory (taste to the anterior two-thirds of the tongue) functions [3]. It is the one of the most commonly paralyzed nerves in the body, and most of the clinical attention it receives focuses on its role in facial expression [8]. Tinnitus, conductive and sensorineural hearing loss, and hemifacial spasm may also signal a lesion involving the facial nerve [15].

The intracranial course of the facial nerve includes pontine, cisternal, and intratemporal segments [7]. Within the pons, the facial nuclei can be affected by intra-axial conditions such as infarction, vascular malformations, tumors, and multiple sclerosis [77]. As the nerve exits the brain stem and courses through the temporal bone, it may be affected by facial and vestibular schwannomas, meningiomas, vascular lesions, inflammation, cholesteatomas, paragangliomas, trauma, and intrinsic bone tumors [15,78]. The extracranial segment of the facial nerve courses through the parotid gland and may be affected by parotid tumors and inflammation and conditions of the neighboring anatomic spaces and skull base such as carcinomas, sarcomas, trauma, and inflammatory disease.

#### **MRI**

MRI is the mainstay of evaluating both intracranial and extracranial portions of the facial nerve [15,79-83]. Facial paralysis in the form of Bell palsy is one of the most common syndromes confronting the otolaryngologist. In general, Bell palsy patients need not be imaged unless the symptoms are atypical or persist for >2 months [15]. When imaging is considered, MRI is the method of choice [15,82,84]. Enhancement may be seen in the canicular, labyrinthine, geniculate, tympanic, and mastoid portions of the nerve in neuritis, although geniculate, tympanic, and mastoid portions may enhance normally [15,85-89]. MRI may also be useful in establishing prognosis [88,90-94]; however, there is one current study with a small cohort of patients that shows no association between the degree of enhancement and the clinical severity of facial nerve palsy in the early stage, stating that predicting the prognosis is difficult [95].

#### **CT**

CT provides useful information regarding temporal bone fractures and trauma, presurgical osseous anatomy, nerve involvement with inflammatory middle ear disease, foraminal expansion, patterns of bone erosion, and intrinsic bone tumor matrices [83,84,96-98]. In patients with risk for contrast allergy and contrast-induced nephropathy, noncontrast CT may be sufficient if patients cannot undergo MRI. A dedicated temporal bone CT with thin sections should be obtained instead of a head CT to evaluate the course of CN VII.

#### **US**

US is not routinely used in the initial evaluation of the facial nerve.

#### **FDG-PET/CT**

FDG-PET/CT is not routinely used in the initial evaluation of the facial nerve.

### **Variant 4: Palate weakness. Oropharyngeal pain. (Glossopharyngeal nerve, CN IX.)**

The glossopharyngeal nerve (CN IX) arises in the medulla and is responsible for branchial motor innervation to the stylopharyngeus muscle, which elevates the palate, and visceral motor parasympathetic innervation to the parotid gland [3]. Visceral sensory innervation to the carotid sinus plays a role in regulating circulation and general and special sensory functions that supply sensation and taste to the posterior tongue. The nerve exits the jugular foramen in close proximity to the vagus (CN X) and the spinal accessory (CN XI) nerves [3,7,77,99]. Therefore, isolated syndromes involving the glossopharyngeal nerve are rare. Intra-axial lesions include gliomas, lymphomas, metastases, vascular malformations, infarctions, and inflammatory abnormalities. Multiple sclerosis may also affect the medulla adjacent to the cranial nerve nuclei. Leptomeningeal metastases, granulomatous disease, and even tortuous or aneurysmal dilatation of vessels may affect the nerve as it enters the cistern. Lesions in the region of the posterior skull base and jugular foramen, such as metastases, schwannomas, paragangliomas, and meningiomas, usually also involve the other lower cranial nerves [99]. Tonsillar pain syndromes, palate weakness, and loss of gag reflex accompanied by loss of taste and sensation in the posterior pharynx may signal a glossopharyngeal nerve lesion [99].

## **MRI and CT**

As with the other cranial nerves, MRI of CN IX is the preferred modality for investigating possible lesions such as masses or vascular compression, with CT providing information on the bony integrity of the foramina [7,100-103]. Imaging protocols should focus on the posterior skull base and upper neck.

## **US**

US is not routinely used in the initial evaluation of the glossopharyngeal nerve.

## **FDG-PET/CT**

FDG-PET/CT is not routinely used in the initial evaluation of the glossopharyngeal nerve.

## **Variant 5: Vocal cord paralysis. (Vagal nerve, CN X.)**

The vagus nerve (CN X) supplies visceral sensation to the pharynx, larynx, and viscera and general sensation to the ear. Branchial motor branches innervate muscles of the pharynx and larynx, whereas visceral motor branches play a predominant role in parasympathetic supply to the thorax and abdomen [3,7,99]. The vagus nerve boasts the longest course in the body of any cranial nerve and is therefore vulnerable to a wide range of pathologies occurring throughout its trajectory from the posterior fossa and skull base to the neck, thorax, and abdomen [8]. Intracranial processes such as meningiomas, schwannomas, metastases, granulomatous disease, ischemia, vascular conditions, and infection may affect the vagal nuclei and the nerve as it exits the medulla. Paragangliomas, schwannomas, and metastases involving the skull base may affect the nerve and the neighboring glossopharyngeal nerve (CN IX) by infiltration of fibers or by compression. Within the neck, trauma may also affect the vagus nerve, in addition to masses, vascular lesions, thyroid conditions, infection, or inflammation [104]. Viral neuropathy may be one of the most common causes of idiopathic vagal palsies [104].

One of the most troubling symptoms of vagus dysfunction is vocal cord paralysis. Because lesions anywhere in the long course of the nerve may potentially cause paralysis, the imaging protocol must visualize the full extent of the nerve from the skull base to the mid chest [99].

## **MRI and CT**

With its rapid scanning time and availability, CT provides an excellent means of examining the lower course of the nerve [105]. Moreover, thoracic causes of paralysis, such as lung cancer, tuberculosis, and thoracic aortic aneurysm, are common [106]. Although chest radiographs may detect many of these causes, chest CT is more sensitive, especially for lesions concealed in the aortopulmonary window [106,107]. This can also be accomplished by extending the neck CT scanning to the mid thorax. For imaging of the upper course of the nerve including the skull base, MRI is preferred [103,108,109]. For the mid neck and larynx, CT and MRI complement one another [110,111]. For example, CT may differentiate traumatic arytenoid dislocation from neurogenic paralysis [112]. Rapid multislice CT scanning, including functional 3-D applications, also allows the patient to perform phonation and breathing maneuvers during imaging to augment diagnosis [113-117]. In patients with risk for contrast allergy and contrast-induced nephropathy, noncontrast CT may be sufficient if patients cannot undergo MRI.

## **US**

US may also have a role in imaging of the neck [104,105]. It may be useful for assessing lesions such as tumors or lymphadenopathy that have caused CN X neuropathy. It can be used in cases of neck lesions as a problem-solving technique. It can also be used as a technique to guide biopsies of lesions in the neck.

## **FDG-PET/CT**

FDG-PET/CT imaging used for evaluating head and neck malignancy may yield false-positive findings in the larynx for patients with vocal cord paralysis or unrecognized physiological asymmetry [118,119]. It may be useful as a problem-solving technique after initial cross-sectional imaging in patients with a known primary malignancy. FDG-PET/CT may also be superior to cross-sectional imaging for both localization and determination of response to therapy [120-124].

## **Radiography**

Chest radiographs can be used as a screening tool if chest CT or chest MRI is unavailable or contraindicated. It can reveal lesions in the lung apex or mediastinum that may cause CN X deficits.

**Variant 6: Weakness or paralysis of the sternocleidomastoid and trapezius muscles. (Accessory nerve, CN XI.)**

The spinal accessory nerve (CN XI) supplies the sternocleidomastoid muscle and the upper portion of the trapezius muscle as its sole branchial motor function [3,7,99]. Palsy is clinically manifested by weakness and atrophy of these muscles and may be accompanied by evidence of involvement of the glossopharyngeal (CN IX) and vagus (CN X) nerves in combined syndromes [99]. Loss of volume and fatty infiltration of the sternocleidomastoid and trapezius muscles may be noted on imaging.

**MRI and CT**

CT and MRI are complementary in diagnosing conditions such as posterior fossa and skull base infarctions, vascular lesions, Chiari malformations, paragangliomas, schwannomas, meningiomas, and metastases or in recognizing nerve involvement from prior neck surgeries [99,103,104]. In patients with risk for contrast allergy and contrast-induced nephropathy, noncontrast CT may be sufficient if patients cannot undergo MRI. Protocol with thin-section MRI should be performed to evaluate the cranial nerves.

**US**

US is not routinely used in the initial evaluation of the accessory nerve.

**FDG-PET/CT**

FDG-PET/CT is not routinely used in the initial evaluation of the accessory nerve.

**Variant 7: Weakness or paralysis of the tongue. (Hypoglossal nerve, CN XII.)**

The hypoglossal nerve (CN XII) supplies somatic motor innervation to the intrinsic and extrinsic muscles of the tongue, except the palatoglossus muscle [3,7,99]. Palsy of this nerve is recognized by dysarthria and deviation of the tongue to the affected side on protrusion. Atrophy and fatty infiltration of the tongue may be noted on imaging. Lesions of the posterior fossa, skull base, upper neck, and floor of the mouth may affect the hypoglossal nerve. They include infarctions, meningiomas, schwannomas, paragangliomas, carcinomas, metastases, subarachnoid hemorrhage, Chiari malformations, basilar invagination, and fractures [99].

**MRI and CT**

As with the other lower cranial nerves, MRI is the preferred modality for CN XII, and CT provides complementary information on the integrity of the bony structures and foramina [125]. Evaluation of the entire course of the nerve is required, which includes evaluation of the nucleus in the brain stem medulla and the nerve in the cisternal segment and high carotid space. This is preferably obtained with a neck MRI that covers the entire nerve pathway. In patients with risk for contrast allergy and contrast-induced nephropathy, noncontrast CT may be sufficient if patients cannot undergo MRI.

**US**

US is not routinely used in the initial evaluation of the hypoglossal nerve.

**FDG-PET/CT**

FDG-PET/CT is not routinely used in the initial evaluation of the hypoglossal nerve.

**Variant 8: Perineural spread of tumor. (Most commonly trigeminal nerve [CN V], facial nerve [CN VII].)**

Because of the complex anatomy of the head and neck and the close proximity of several cranial nerves, many clinical presentations of cranial neuropathy involve multiple nerves. As in syndromes of combined neuropathy of the upper cranial nerves, such as those related to vision and the extraocular muscles (which are covered in other Appropriateness Criteria), syndromes involving the lower cranial nerves are also grouped primarily by the proximity of the involved cranial nerves. For example, Gradenigo syndrome involves CNs V and VI as they travel in the vicinity of the petrous apex, whereas Vernet syndrome involves CNs IX, X, and XI as they travel within the jugular foramen. Collet-Sicard syndrome involves CNs IX, X, XI, and XII related to lesions just below the skull base or large lesions affecting both the jugular foramen and the hypoglossal canal. Imaging protocols should be tailored to evaluate the suspected region of anatomy when the syndrome is identified by the clinician.

A difficult problem for the surgeon is the perineural spread of head and neck malignancy. The trigeminal (CN V) and facial (CN VII) are the most common nerves involved; however, any cranial nerve traveling in the vicinity of a malignancy may become involved [126]. Perineural spread of tumor along the facial nerve may evade even the most meticulous imaging [127]. Subtle clues such as nerve enhancement, nerve enlargement, foraminal expansion, or muscle volume loss may indicate cranial nerve involvement with tumor [128]. For example,

asymmetry of facial musculature may be useful in detecting perineural tumor spread along the facial nerve or predicting return of function after nerve grafting [91,128].

### **MRI and CT**

MRI has emerged as the preferred imaging method for evaluating the perineural spread of tumor, although CT may be very useful for visualizing the neural foramina [126,129-134]. In patients with risk for contrast allergy and contrast-induced nephropathy, noncontrast CT may be sufficient if patients cannot undergo MRI. Protocol with thin-section MRI should be performed to evaluate the cranial nerves.

### **FDG-PET/CT**

FDG-PET/CT imaging may also be helpful [133,135]. It may be useful as a problem-solving technique after initial cross-sectional imaging in patients with a known primary malignancy. FDG-PET/CT may also be superior to cross-sectional imaging for both localization and determination of response to therapy [120-124].

### **US**

US is not routinely used in the initial evaluation of the perineural spread of tumor.

### **Summary of Recommendations**

- Pathology affecting the olfactory nerve is best evaluated with contrast-enhanced MRI. The protocol should be tailored to the anterior cranial fossa. CT may be useful in the evaluation of sinus inflammatory disease and trauma.
- Contrast-enhanced MRI is the primary technique to evaluate trigeminal neuralgia. MR angiography can be used to evaluate vascular compression.
- Facial nerve neuropathy is best evaluated with a dedicated contrast-enhanced MRI of the orbit, face, and neck, tailored to the temporal bone and parotid area.
- In the evaluation of cranial nerve IX symptoms, a focused contrast-enhanced MRI tailored to the posterior fossa is the study of choice.
- Cranial nerve X paralysis is well evaluated with either contrast-enhanced MRI or contrast-enhanced neck CT. Pathology in the posterior fossa will be better demonstrated with MRI. The complete evaluation of the nerve requires imaging the upper chest to the level of the anteroposterior window.
- In the evaluation of cranial nerve XI symptoms, a focused contrast-enhanced MRI tailored to the posterior fossa is the study of choice. Contrast-enhanced CT of the neck is complementary to the skull base imaging.
- Evaluation of lesions affecting cranial nerve XII is best done with a contrast-enhanced MRI tailored to the posterior fossa. The evaluation of the neck can also be done with contrast-enhanced neck CT.
- Perineural tumor spread most commonly affects CNs V and VII. Evaluation is best done with contrast-enhanced MRI tailored to the skull base.

### **Summary of Evidence**

Of the 136 references cited in the *ACR Appropriateness Criteria® Cranial Neuropathy* document, 2 are categorized as therapeutic references, including 1 good-quality study. Additionally, 134 references are categorized as diagnostic references, including 3 well-designed studies, 19 good-quality studies, and 46 quality studies that may have design limitations. There are 67 references that may not be useful as primary evidence.

The 136 references cited in the *ACR Appropriateness Criteria® Cranial Neuropathy* document were published from 1998 to 2017.

Although there are references that report on studies with design limitations, 23 well-designed or good-quality studies provide good evidence.

### **Relative Radiation Level Information**

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Patients in the pediatric age group are at inherently higher risk from exposure, both because of organ sensitivity and longer life expectancy (relevant to the

long latency that appears to accompany radiation exposure). For these reasons, the RRL dose estimate ranges for pediatric examinations are lower as compared to those specified for adults (see Table below). Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® [Radiation Dose Assessment Introduction](#) document [136].

Relative Radiation Level Designations		
Relative Radiation Level*	Adult Effective Dose Estimate Range	Pediatric Effective Dose Estimate Range
○	0 mSv	0 mSv
⊕	<0.1 mSv	<0.03 mSv
⊕⊕	0.1-1 mSv	0.03-0.3 mSv
⊕⊕⊕	1-10 mSv	0.3-3 mSv
⊕⊕⊕⊕	10-30 mSv	3-10 mSv
⊕⊕⊕⊕⊕	30-100 mSv	10-30 mSv

\*RRL assignments for some of the examinations cannot be made, because the actual patient doses in these procedures vary as a function of a number of factors (eg, region of the body exposed to ionizing radiation, the imaging guidance that is used). The RRLs for these examinations are designated as “Varies”.

### Supporting Documents

For additional information on the Appropriateness Criteria methodology and other supporting documents go to [www.acr.org/ac](http://www.acr.org/ac).

### References

- Laine FJ, Smoker WR. Anatomy of the cranial nerves. *Neuroimaging Clin N Am*. 1998;8(1):69-100.
- Rubin M, Safdieh JE, Netter FH. Cranial nerves I-XII. *Netter's concise neuroanatomy*. Philadelphia: Saunders Elsevier; 2007:215-263.
- Wilson-Pauwels L, Akesson EJ, Stewart PA. Introduction. In: *Cranial nerves: anatomy and clinical comments*. Toronto, Ontario, Canada: B.C. Decker; 1988:vii-xiii.
- American College of Radiology. ACR Appropriateness Criteria®: Orbits, Vision and Visual Loss. Available at: <https://acsearch.acr.org/docs/69486/Narrative/>. Accessed March 1, 2017.
- American College of Radiology. ACR Appropriateness Criteria®: Hearing Loss and/or Vertigo. Available at: <https://acsearch.acr.org/docs/69488/Narrative/>. Accessed March 1, 2017.
- Chen RC, Khorsandi AS, Shatzkes DR, Holliday RA. The radiology of referred otalgia. *AJNR Am J Neuroradiol*. 2009;30(10):1817-1823.
- Casselmann J, Mermuys K, Delanote J, Ghekiere J, Coenegrachts K. MRI of the cranial nerves--more than meets the eye: technical considerations and advanced anatomy. *Neuroimaging Clin N Am*. 2008;18(2):197-231, preceding x.
- Borges A, Casselman J. Imaging the cranial nerves: Part I: methodology, infectious and inflammatory, traumatic and congenital lesions. *Eur Radiol*. 2007;17(8):2112-2125.
- Casselmann JW, Kuhweide R, Deimling M, Ampe W, Dehaene I, Meeus L. Constructive interference in steady state-3DFT MR imaging of the inner ear and cerebellopontine angle. *AJNR Am J Neuroradiol*. 1993;14(1):47-57.
- Ciftci E, Anik Y, Arslan A, Akansel G, Sarisoy T, Demirci A. Driven equilibrium (drive) MR imaging of the cranial nerves V-VIII: comparison with the T2-weighted 3D TSE sequence. *Eur J Radiol*. 2004;51(3):234-240.
- Jung NY, Moon WJ, Lee MH, Chung EC. Magnetic resonance cisternography: comparison between 3-dimensional driven equilibrium with sensitivity encoding and 3-dimensional balanced fast-field echo sequences with sensitivity encoding. *J Comput Assist Tomogr*. 2007;31(4):588-591.
- Lane JI, Ward H, Witte RJ, Bernstein MA, Driscoll CL. 3-T imaging of the cochlear nerve and labyrinth in cochlear-implant candidates: 3D fast recovery fast spin-echo versus 3D constructive interference in the steady state techniques. *AJNR Am J Neuroradiol*. 2004;25(4):618-622.



13. Tsuchiya K, Aoki C, Hachiya J. Evaluation of MR cisternography of the cerebellopontine angle using a balanced fast-field-echo sequence: preliminary findings. *Eur Radiol.* 2004;14(2):239-242.
14. Tsuchiya K, Yamakami N, Hachiya J, Kassai Y. MR cisternography using a three-dimensional half-fourier single-shot fast spin-echo sequence. *Eur Radiol.* 1998;8(3):424-426.
15. Veillon F, Taboada LR, Eid MA, et al. Pathology of the facial nerve. *Neuroimaging Clin N Am.* 2008;18(2):309-320, x.
16. Linn J, Peters F, Moriggl B, Naidich TP, Bruckmann H, Yousry I. The jugular foramen: imaging strategy and detailed anatomy at 3T. *AJNR Am J Neuroradiol.* 2009;30(1):34-41.
17. Erdogan N, Altay C, Akay E, et al. MRI assessment of internal acoustic canal variations using 3D-FIESTA sequences. *Eur Arch Otorhinolaryngol.* 2013;270(2):469-475.
18. American College of Radiology. ACR Appropriateness Criteria®: Seizures and Epilepsy. Available at: <https://acsearch.acr.org/docs/69479/Narrative/>. Accessed March 1, 2017.
19. Castillo M. Imaging of the upper cranial nerves I, III-VIII, and the cavernous sinuses. *Magn Reson Imaging Clin N Am.* 2002;10(3):415-431, v.
20. Abolmaali N, Gudziol V, Hummel T. Pathology of the olfactory nerve. *Neuroimaging Clin N Am.* 2008;18(2):233-242, preceding x.
21. Madan R, Sawlani V, Gupta S, Phadke RV. MRI findings in Kallmann syndrome. *Neurol India.* 2004;52(4):501-503.
22. Koenigkam-Santos M, Santos AC, Versiani BR, Diniz PR, Junior JE, de Castro M. Quantitative magnetic resonance imaging evaluation of the olfactory system in Kallmann syndrome: correlation with a clinical smell test. *Neuroendocrinology.* 2011;94(3):209-217.
23. Berendse HW, Booij J, Francot CM, et al. Subclinical dopaminergic dysfunction in asymptomatic Parkinson's disease patients' relatives with a decreased sense of smell. *Ann Neurol.* 2001;50(1):34-41.
24. Mueller A, Rodewald A, Reden J, Gerber J, von Kummer R, Hummel T. Reduced olfactory bulb volume in post-traumatic and post-infectious olfactory dysfunction. *Neuroreport.* 2005;16(5):475-478.
25. Rombaux P, Mouraux A, Bertrand B, Nicolas G, Duprez T, Hummel T. Olfactory function and olfactory bulb volume in patients with postinfectious olfactory loss. *Laryngoscope.* 2006;116(3):436-439.
26. Negoias S, Croy I, Gerber J, et al. Reduced olfactory bulb volume and olfactory sensitivity in patients with acute major depression. *Neuroscience.* 2010;169(1):415-421.
27. Haruna S, Otori N, Moriyama H, Nakanishi M. Olfactory dysfunction in sinusitis with infiltration of numerous activated eosinophils. *Auris Nasus Larynx.* 2006;33(1):23-30.
28. Trotier D, Bensimon JL, Herman P, Tran Ba Huy P, Doving KB, Eloit C. Inflammatory obstruction of the olfactory clefts and olfactory loss in humans: a new syndrome? *Chem Senses.* 2007;32(3):285-292.
29. Eftekhari M, Assadi M, Kazemi M, et al. Brain perfusion single photon emission computed tomography findings in patients with posttraumatic anosmia and comparison with radiological imaging. *Am J Rhinol.* 2006;20(6):577-581.
30. Mainland JD, Johnson BN, Khan R, Ivry RB, Sobel N. Olfactory impairments in patients with unilateral cerebellar lesions are selective to inputs from the contralesional nostril. *J Neurosci.* 2005;25(27):6362-6371.
31. Mann NM, Vento JA. A study comparing SPECT and MRI in patients with anosmia after traumatic brain injury. *Clin Nucl Med.* 2006;31(8):458-462.
32. Westermann B, Wattendorf E, Schwerdtfeger U, et al. Functional imaging of the cerebral olfactory system in patients with Parkinson's disease. *J Neurol Neurosurg Psychiatry.* 2008;79(1):19-24.
33. Wang J, Eslinger PJ, Doty RL, et al. Olfactory deficit detected by fMRI in early Alzheimer's disease. *Brain Res.* 2010;1357:184-194.
34. Becker M, Kohler R, Vargas MI, Viallon M, Delavelle J. Pathology of the trigeminal nerve. *Neuroimaging Clin N Am.* 2008;18(2):283-307, x.
35. Broggi G, Ferroli P, Franzini A, et al. Operative findings and outcomes of microvascular decompression for trigeminal neuralgia in 35 patients affected by multiple sclerosis. *Neurosurgery.* 2004;55(4):830-838; discussion 838-839.
36. da Silva CJ, da Rocha AJ, Mendes MF, Maia AC, Jr., Braga FT, Tilbery CP. Trigeminal involvement in multiple sclerosis: magnetic resonance imaging findings with clinical correlation in a series of patients. *Mult Scler.* 2005;11(3):282-285.
37. Ma X, Sun X, Yao J, et al. Clinical analysis of trigeminal neuralgia caused by vertebrobasilar dolichoectasia. *Neurosurg Rev.* 2013;36(4):573-577; discussion 577-578.

38. Maarbjerg S, Wolfram F, Gozalov A, Olesen J, Bendtsen L. Significance of neurovascular contact in classical trigeminal neuralgia. *Brain*. 2015;138(Pt 2):311-319.
39. Kress B, Rasche D, Fiebach J, Tronnier V, Sartor K, Stippich C. [MR volumetry of the trigeminal nerve in patients with unilateral facial pain]. *Rofo*. 2004;176(5):719-723.
40. Kato K, Tomura N, Takahashi S, Watarai J. Motor denervation of tumors of the head and neck: changes in MR appearance. *Magn Reson Med Sci*. 2002;1(3):157-164.
41. Anderson VC, Berryhill PC, Sandquist MA, Ciaverella DP, Nesbit GM, Burchiel KJ. High-resolution three-dimensional magnetic resonance angiography and three-dimensional spoiled gradient-recalled imaging in the evaluation of neurovascular compression in patients with trigeminal neuralgia: a double-blind pilot study. *Neurosurgery*. 2006;58(4):666-673; discussion 666-673.
42. Chang JW, Chang JH, Park YG, Chung SS. Microvascular decompression in trigeminal neuralgia: a correlation of three-dimensional time-of-flight magnetic resonance angiography and surgical findings. *Stereotact Funct Neurosurg*. 2000;74(3-4):167-174.
43. Gorgulho A, De Salles AA, McArthur D, et al. Brainstem and trigeminal nerve changes after radiosurgery for trigeminal pain. *Surg Neurol*. 2006;66(2):127-135; discussion 135.
44. Kress B, Schindler M, Rasche D, et al. MRI volumetry for the preoperative diagnosis of trigeminal neuralgia. *Eur Radiol*. 2005;15(7):1344-1348.
45. Kuncz A, Voros E, Barzo P, Tajti J, Milassin P, Bodosi M. [The role of MR angiography in predicting operative results of microvascular decompression in patients with trigeminal neuralgia]. *Orv Hetil*. 2005;146(51):2595-2602.
46. Kuncz A, Voros E, Barzo P, et al. Comparison of clinical symptoms and magnetic resonance angiographic (MRA) results in patients with trigeminal neuralgia and persistent idiopathic facial pain. Medium-term outcome after microvascular decompression of cases with positive MRA findings. *Cephalalgia*. 2006;26(3):266-276.
47. Miller J, Acar F, Hamilton B, Burchiel K. Preoperative visualization of neurovascular anatomy in trigeminal neuralgia. *J Neurosurg*. 2008;108(3):477-482.
48. Ogiwara M, Shimizu T. Surface rendered three-dimensional MR imaging for the evaluation of trigeminal neuralgia and hemifacial spasm. *J Clin Neurosci*. 2004;11(8):840-844.
49. Satoh T, Omi M, Ohsako C, Nabeshima M, Onoda K, Date I. [Novel 3D MR angiographic findings of neurovascular compressive sites in patients with trigeminal neuralgia]. *No Shinkei Geka*. 2007;35(3):259-265.
50. Satoh T, Onoda K, Date I. Preoperative simulation for microvascular decompression in patients with idiopathic trigeminal neuralgia: visualization with three-dimensional magnetic resonance cisternogram and angiogram fusion imaging. *Neurosurgery*. 2007;60(1):104-113; discussion 113-104.
51. Tanaka T, Morimoto Y, Shiiba S, et al. Utility of magnetic resonance cisternography using three-dimensional fast asymmetric spin-echo sequences with multiplanar reconstruction: the evaluation of sites of neurovascular compression of the trigeminal nerve. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2005;100(2):215-225.
52. Voros E, Palko A, Horvath K, Barzo P, Kardos L, Kuncz A. Three-dimensional time-of-flight MR angiography in trigeminal neuralgia on a 0.5-T system. *Eur Radiol*. 2001;11(4):642-647.
53. Yamakami I, Kobayashi E, Hirai S, Yamaura A. Preoperative assessment of trigeminal neuralgia and hemifacial spasm using constructive interference in steady state-three-dimensional Fourier transformation magnetic resonance imaging. *Neurol Med Chir (Tokyo)*. 2000;40(11):545-555; discussion 555-546.
54. Borges A, Casselman J. Imaging the trigeminal nerve. *Eur J Radiol*. 2010;74(2):323-340.
55. Shimizu M, Imai H, Kagoshima K, Umezawa E, Shimizu T, Yoshimoto Y. Detection of compression vessels in trigeminal neuralgia by surface-rendering three-dimensional reconstruction of 1.5- and 3.0-T magnetic resonance imaging. *World Neurosurg*. 2013;80(3-4):378-385.
56. Zeng Q, Zhou Q, Liu Z, Li C, Ni S, Xue F. Preoperative detection of the neurovascular relationship in trigeminal neuralgia using three-dimensional fast imaging employing steady-state acquisition (FIESTA) and magnetic resonance angiography (MRA). *J Clin Neurosci*. 2013;20(1):107-111.
57. Garcia M, Naraghi R, Zumbunn T, Rosch J, Hastreiter P, Dorfler A. High-resolution 3D-constructive interference in steady-state MR imaging and 3D time-of-flight MR angiography in neurovascular compression: a comparison between 3T and 1.5T. *AJNR Am J Neuroradiol*. 2012;33(7):1251-1256.
58. Leal PR, Hermier M, Souza MA, Cristino-Filho G, Froment JC, Sindou M. Visualization of vascular compression of the trigeminal nerve with high-resolution 3T MRI: a prospective study comparing

- preoperative imaging analysis to surgical findings in 40 consecutive patients who underwent microvascular decompression for trigeminal neuralgia. *Neurosurgery*. 2011;69(1):15-25; discussion 26.
59. Alberico RA, Fenstermaker RA, Lobel J. Focal enhancement of cranial nerve V after radiosurgery with the Leksell gamma knife: experience in 15 patients with medically refractory trigeminal neuralgia. *AJNR Am J Neuroradiol*. 2001;22(10):1944-1948.
  60. Erbay SH, Bhadelia RA, O'Callaghan M, et al. Nerve atrophy in severe trigeminal neuralgia: noninvasive confirmation at MR imaging--initial experience. *Radiology*. 2006;238(2):689-692.
  61. Erbay SH, Bhadelia RA, Riesenburger R, et al. Association between neurovascular contact on MRI and response to gamma knife radiosurgery in trigeminal neuralgia. *Neuroradiology*. 2006;48(1):26-30.
  62. Flickinger JC, Pollock BE, Kondziolka D, et al. Does increased nerve length within the treatment volume improve trigeminal neuralgia radiosurgery? A prospective double-blind, randomized study. *Int J Radiat Oncol Biol Phys*. 2001;51(2):449-454.
  63. Lang E, Naraghi R, Tanrikulu L, et al. Neurovascular relationship at the trigeminal root entry zone in persistent idiopathic facial pain: findings from MRI 3D visualisation. *J Neurol Neurosurg Psychiatry*. 2005;76(11):1506-1509.
  64. Lim M, Cotrutz C, Romanelli P, et al. Stereotactic radiosurgery using CT cisternography and non-isocentric planning for the treatment of trigeminal neuralgia. *Comput Aided Surg*. 2006;11(1):11-20.
  65. Liu M, Wu CY, Liu YG, Wang HW, Meng FG. Three-dimensional computed tomography-guided radiofrequency trigeminal rhizotomy for treatment of idiopathic trigeminal neuralgia. *Chin Med Sci J*. 2005;20(3):206-209.
  66. Lorenzoni JG, Massager N, David P, et al. Neurovascular compression anatomy and pain outcome in patients with classic trigeminal neuralgia treated by radiosurgery. *Neurosurgery*. 2008;62(2):368-375; discussion 375-366.
  67. Massager N, Abeloos L, Devriendt D, Op de Beeck M, Levivier M. Clinical evaluation of targeting accuracy of gamma knife radiosurgery in trigeminal neuralgia. *Int J Radiat Oncol Biol Phys*. 2007;69(5):1514-1520.
  68. Peker S, Bayrakli F, Kilic T, Pamir MN. Gamma-knife radiosurgery in the treatment of trigeminal schwannomas. *Acta Neurochir (Wien)*. 2007;149(11):1133-1137; discussion 1137.
  69. Sekimoto K, Koizuka S, Saito S, Goto F. Thermogangliolysis of the Gasserian ganglion under computed tomography fluoroscopy. *J Anesth*. 2005;19(2):177-179.
  70. Shaya M, Jawahar A, Caldito G, Sin A, Willis BK, Nanda A. Gamma knife radiosurgery for trigeminal neuralgia: a study of predictors of success, efficacy, safety, and outcome at LSUHSC. *Surg Neurol*. 2004;61(6):529-534; discussion 534-525.
  71. Zhang WJ, Wang Y, Chen MJ. [Research of the puncture depth in treatment of trigeminal neuralgia (TN) by radiofrequency thermocoagulation(RFT) under CT location]. *Shanghai Kou Qiang Yi Xue*. 2003;12(2):94-95.
  72. Park KJ, Kano H, Berkowitz O, et al. Computed tomography-guided gamma knife stereotactic radiosurgery for trigeminal neuralgia. *Acta Neurochir (Wien)*. 2011;153(8):1601-1609.
  73. Sheehan JP, Ray DK, Monteith S, et al. Gamma Knife radiosurgery for trigeminal neuralgia: the impact of magnetic resonance imaging-detected vascular impingement of the affected nerve. *J Neurosurg*. 2010;113(1):53-58.
  74. Herweh C, Kress B, Rasche D, et al. Loss of anisotropy in trigeminal neuralgia revealed by diffusion tensor imaging. *Neurology*. 2007;68(10):776-778.
  75. Oishi M, Fukuda M, Takao T, Ishida G, Sato M, Fujii Y. [The utility of presurgical simulation of microvascular decompression by MR virtual endoscopy]. *No Shinkei Geka*. 2007;35(11):1087-1095.
  76. Lutz J, Linn J, Mehrkens JH, et al. Trigeminal neuralgia due to neurovascular compression: high-spatial-resolution diffusion-tensor imaging reveals microstructural neural changes. *Radiology*. 2011;258(2):524-530.
  77. Laine FJ, Underhill T. Imaging of the lower cranial nerves. *Neuroimaging Clin N Am*. 2004;14(4):595-609.
  78. Toulgoat F, Sarrazin JL, Benoudiba F, et al. Facial nerve: from anatomy to pathology. *Diagn Interv Imaging*. 2013;94(10):1033-1042.
  79. Jemec B, Grobbelaar AO, Harrison DH. The abnormal nucleus as a cause of congenital facial palsy. *Arch Dis Child*. 2000;83(3):256-258.

80. Kinoshita T, Ishii K, Okitsu T, Ogawa T, Okudera T. High-intensity facial nerve lesions on T2-weighted images in chronic persistent facial nerve palsy. *Neuroradiology*. 2001;43(5):388-392.
81. Park SU, Kim HJ, Cho YK, et al. The usefulness of MR imaging of the temporal bone in the evaluation of patients with facial and audiovestibular dysfunction. *Korean J Radiol*. 2002;3(1):16-23.
82. Shinohara Y, Kinoshita T, Sugihara S, et al. [MR imaging for evaluation of severe facial nerve damage in patients with facial nerve palsy]. *Nihon Igaku Hoshasen Gakkai Zasshi*. 2005;65(4):353-358.
83. Raghavan P, Mukherjee S, Phillips CD. Imaging of the facial nerve. *Neuroimaging Clin N Am*. 2009;19(3):407-425.
84. Bodenez C, Darrouzet V, Rouanet-Larriviere M, et al. [Facial paralysis after temporal bone trauma]. *Ann Otolaryngol Chir Cervicofac*. 2006;123(1):9-16.
85. Kim IS, Shin SH, Kim J, Lee WS, Lee HK. Correlation between MRI and operative findings in Bell's palsy and Ramsay Hunt syndrome. *Yonsei Med J*. 2007;48(6):963-968.
86. Kinoshita T, Ishii K, Okitsu T, Okudera T, Ogawa T. Facial nerve palsy: evaluation by contrast-enhanced MR imaging. *Clin Radiol*. 2001;56(11):926-932.
87. Seok JI, Lee DK, Kim KJ. The usefulness of clinical findings in localising lesions in Bell's palsy: comparison with MRI. *J Neurol Neurosurg Psychiatry*. 2008;79(4):418-420.
88. Yetiser S, Kazkayas M, Altinok D, Karadeniz Y. Magnetic resonance imaging of the intratemporal facial nerve in idiopathic peripheral facial palsy. *Clin Imaging*. 2003;27(2):77-81.
89. Hong HS, Yi BH, Cha JG, et al. Enhancement pattern of the normal facial nerve at 3.0 T temporal MRI. *Br J Radiol*. 2010;83(986):118-121.
90. Belveze P, Guichard C, Gabrillargues J, Mom T, Gilain L. [Magnetic Resonance Imaging in Facial Bell's Palsy]. *Ann Otolaryngol Chir Cervicofac*. 2002;119(2):81-88.
91. Kaylie DM, Wax MK, Weissman JL. Preoperative facial muscle imaging predicts final facial function after facial nerve grafting. *AJNR Am J Neuroradiol*. 2003;24(3):326-330.
92. Kress B, Griesbeck F, Stippich C, Bahren W, Sartor K. Bell palsy: quantitative analysis of MR imaging data as a method of predicting outcome. *Radiology*. 2004;230(2):504-509.
93. Kress BP, Griesbeck F, Efinger K, et al. [The prognostic value of quantified MRI at an early stage of Bell's palsy]. *Rofo*. 2002;174(4):426-432.
94. Yla-Kotola TM, Kauhanen MS, Koskinen SK, Asko-Seljavaara SL. Magnetic resonance imaging of microneurovascular free muscle flaps in facial reanimation. *Br J Plast Surg*. 2005;58(1):22-27.
95. Jun BC, Chang KH, Lee SJ, Park YS. Clinical feasibility of temporal bone magnetic resonance imaging as a prognostic tool in idiopathic acute facial palsy. *J Laryngol Otol*. 2012;126(9):893-896.
96. Ulug T, Arif Ulubil S. Management of facial paralysis in temporal bone fractures: a prospective study analyzing 11 operated fractures. *Am J Otolaryngol*. 2005;26(4):230-238.
97. Yu Z, Han D, Dai H, Zhao S, Zheng Y. Diagnosis of the pathological exposure of the mastoid portion of the facial nerve by CT scanning. *Acta Otolaryngol*. 2007;127(3):323-327.
98. Vianna M, Adams M, Schachern P, Lazarini PR, Paparella MM, Cureoglu S. Differences in the diameter of facial nerve and facial canal in bell's palsy--a 3-dimensional temporal bone study. *Otol Neurotol*. 2014;35(3):514-518.
99. Policeni BA, Smoker WR. Pathologic conditions of the lower cranial nerves IX, X, XI, and XII. *Neuroimaging Clin N Am*. 2008;18(2):347-368, xi.
100. Hiwatashi A, Matsushima T, Yoshiura T, et al. MRI of glossopharyngeal neuralgia caused by neurovascular compression. *AJR Am J Roentgenol*. 2008;191(2):578-581.
101. Tanrikulu L, Hastreiter P, Troescher-Weber R, Buchfelder M, Naraghi R. Intraoperative three-dimensional visualization in microvascular decompression. *J Neurosurg*. 2007;107(6):1137-1143.
102. Gaul C, Hastreiter P, Duncker A, Naraghi R. Diagnosis and neurosurgical treatment of glossopharyngeal neuralgia: clinical findings and 3-D visualization of neurovascular compression in 19 consecutive patients. *J Headache Pain*. 2011;12(5):527-534.
103. Linn J, Moriggl B, Schwarz F, et al. Cisternal segments of the glossopharyngeal, vagus, and accessory nerves: detailed magnetic resonance imaging-demonstrated anatomy and neurovascular relationships. *J Neurosurg*. 2009;110(5):1026-1041.
104. Larson TC, 3rd, Aulino JM, Laine FJ. Imaging of the glossopharyngeal, vagus, and accessory nerves. *Semin Ultrasound CT MR*. 2002;23(3):238-255.
105. Robinson S, Pitkaranta A. Radiology findings in adult patients with vocal fold paralysis. *Clin Radiol*. 2006;61(10):863-867.

106. El Badawey MR, Punekar S, Zammit-Maempel I. Prospective study to assess vocal cord palsy investigations. *Otolaryngol Head Neck Surg.* 2008;138(6):788-790.
107. Bando H, Nishio T, Bamba H, Uno T, Hisa Y. Vocal fold paralysis as a sign of chest diseases: a 15-year retrospective study. *World J Surg.* 2006;30(3):293-298.
108. Infante-Velazquez EJ, Gomez-Fernandez L, Perez del Campo YH, Diaz-Perez MJ, Vergara-Consuegra O. [Brainstem lesions: clinicoradiological electrophysiological correlation when chronic]. *Rev Neurol.* 2002;34(4):317-321.
109. Saglitz SA, Gaab MR. Investigations using magnetic resonance imaging: is neurovascular compression present in patients with essential hypertension? *J Neurosurg.* 2002;96(6):1006-1012.
110. Chin SC, Edelstein S, Chen CY, Som PM. Using CT to localize side and level of vocal cord paralysis. *AJR Am J Roentgenol.* 2003;180(4):1165-1170.
111. Kumar VA, Lewin JS, Ginsberg LE. CT assessment of vocal cord medialization. *AJNR Am J Neuroradiol.* 2006;27(8):1643-1646.
112. Rubin AD, Hawkshaw MJ, Moyer CA, Dean CM, Sataloff RT. Arytenoid cartilage dislocation: a 20-year experience. *J Voice.* 2005;19(4):687-701.
113. Jun BC, Kim HT, Kim HS, Cho SH. Clinical feasibility of the new technique of functional 3D laryngeal CT. *Acta Otolaryngol.* 2005;125(7):774-778.
114. Kim BS, Ahn KJ, Park YH, Hahn ST. Usefulness of laryngeal phonation CT in the diagnosis of vocal cord paralysis. *AJR Am J Roentgenol.* 2008;190(5):1376-1379.
115. Oyamada Y, Yumoto E, Nakano K, Goto H. Asymmetry of the vocal folds in patients with vocal fold immobility. *Arch Otolaryngol Head Neck Surg.* 2005;131(5):399-406.
116. Yumoto E, Nakano K, Oyamada Y. Relationship between 3D behavior of the unilaterally paralyzed larynx and aerodynamic vocal function. *Acta Otolaryngol.* 2003;123(2):274-278.
117. Yumoto E, Oyamada Y, Nakano K, Nakayama Y, Yamashita Y. Three-dimensional characteristics of the larynx with immobile vocal fold. *Arch Otolaryngol Head Neck Surg.* 2004;130(8):967-974.
118. Heller MT, Meltzer CC, Fukui MB, et al. Superphysiologic FDG Uptake in the Non-Paralyzed Vocal Cord. Resolution of a False-Positive PET Result with Combined PET-CT Imaging. *Clin Positron Imaging.* 2000;3(5):207-211.
119. Kamel EM, Goerres GW, Burger C, von Schulthess GK, Steinert HC. Recurrent laryngeal nerve palsy in patients with lung cancer: detection with PET-CT image fusion -- report of six cases. *Radiology.* 2002;224(1):153-156.
120. Bronstein Y, Tummala S, Rohren E. F-18 FDG PET/CT for detection of malignant involvement of peripheral nerves: case series and literature review. *Clin Nucl Med.* 2011;36(2):96-100.
121. Fukui MB, Blodgett TM, Snyderman CH, et al. Combined PET-CT in the head and neck: part 2. Diagnostic uses and pitfalls of oncologic imaging. *Radiographics.* 2005;25(4):913-930.
122. Kim JH, Jang JH, Koh SB. A case of neurolymphomatosis involving cranial nerves: MRI and fusion PET-CT findings. *J Neurooncol.* 2006;80(2):209-210.
123. Komissarova M, Wong KK, Piert M, Mukherji SK, Fig LM. Spectrum of 18F-FDG PET/CT findings in oncology-related recurrent laryngeal nerve palsy. *AJR Am J Roentgenol.* 2009;192(1):288-294.
124. Matsue K, Hayama BY, Iwama K, et al. High frequency of neurolymphomatosis as a relapse disease of intravascular large B-cell lymphoma. *Cancer.* 2011;117(19):4512-4521.
125. Yousry I, Moriggl B, Schmid UD, et al. Detailed anatomy of the intracranial segment of the hypoglossal nerve: neurovascular relationships and landmarks on magnetic resonance imaging sequences. *J Neurosurg.* 2002;96(6):1113-1122.
126. Gandhi D, Gujar S, Mukherji SK. Magnetic resonance imaging of perineural spread of head and neck malignancies. *Top Magn Reson Imaging.* 2004;15(2):79-85.
127. Jungehuelsing M, Sittel C, Fischbach R, Wagner M, Stennert E. Limitations of magnetic resonance imaging in the evaluation of perineural tumor spread causing facial nerve paralysis. *Arch Otolaryngol Head Neck Surg.* 2000;126(4):506-510.
128. Smoker WR, Reede DL. Denervation atrophy of motor cranial nerves. *Neuroimaging Clin N Am.* 2008;18(2):387-411, xi.
129. Blandino A, Gaeta M, Minutoli F, Pandolfo I. CT and MR findings in neoplastic perineural spread along the vidian nerve. *Eur Radiol.* 2000;10(3):521-526.
130. Bowyer JD, Sullivan TJ, Whitehead KJ, Kelly LE, Allison RW. The management of perineural spread of squamous cell carcinoma to the ocular adnexae. *Ophthal Plast Reconstr Surg.* 2003;19(4):275-281.

131. Chang PC, Fischbein NJ, McCalmont TH, et al. Perineural spread of malignant melanoma of the head and neck: clinical and imaging features. *AJNR Am J Neuroradiol*. 2004;25(1):5-11.
132. Galloway TJ, Morris CG, Mancuso AA, Amdur RJ, Mendenhall WM. Impact of radiographic findings on prognosis for skin carcinoma with clinical perineural invasion. *Cancer*. 2005;103(6):1254-1257.
133. Lee KJ, Abemayor E, Sayre J, Bhuta S, Kirsch C. Determination of perineural invasion preoperatively on radiographic images. *Otolaryngol Head Neck Surg*. 2008;139(2):275-280.
134. Maroldi R, Farina D, Borghesi A, Marconi A, Gatti E. Perineural tumor spread. *Neuroimaging Clin N Am*. 2008;18(2):413-429, xi.
135. Conrad GR, Sinha P, Holzhauer M. Perineural spread of skin carcinoma to the base of the skull: detection with FDG PET and CT fusion. *Clin Nucl Med*. 2004;29(11):717-719.
136. American College of Radiology. ACR Appropriateness Criteria® Radiation Dose Assessment Introduction. Available at: <http://www.acr.org/~media/ACR/Documents/AppCriteria/RadiationDoseAssessmentIntro.pdf>. Accessed March 1, 2017.

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.