American College of Radiology
ACR Appropriateness Criteria®

Clinically Suspected Adnexal Mass, No Acute Symptoms

Variant 1: Adult female. Clinically suspected adnexal mass, no acute symptoms. Premenopausal or postmenopausal. Initial imaging.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
US duplex Doppler pelvis	Usually appropriate	Strong	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	1	1	4	1	7

References	Study Quality
19 (9618853)	3
23 (18504770)	3
21 (9646799)	2
20 (10739694)	3
25 (7839975)	3
24 (11176291)	4
22 (31687921)	4

US pelvis transvaginal	Usually appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	0	0	2	2	10
US pelvis transabdominal and US pelvis transvaginal	Usually appropriate	Strong	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	0	0	2	1	11

References	Study Quality
39 (11169340)	4
35 (14529661)	3
32 (19244063)	3
23 (18504770)	3
27 (10586480)	3

26 (10201081)	3
37 (30149950)	4
36 (25320247)	3
43 (28727500)	3
44 (31770022)	3
34 (30303910)	3
22 (31687921)	4
28 (27776067)	4
30 (32736853)	4
29 (34474226)	4
41 (31549945)	4
45 (32809166)	2
31 (12634641)	4
33 (20505067)	4
42 (23358210)	3
38 (24937676)	3
40 (36520422)	3

US pelvis transabdominal	Usually appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	7	7	0	0	0	0	0	4	5	4	1
MRI pelvis without and with IV contrast	May be appropriate	Limited	O 0 mSv	O 0 mSv [ped]	6	6	0	0	1	3	3	6	0	1	0

References	Study Quality
15 (31482379)	3
16 (35040672)	4

MRI pelvis without IV contrast	May be appropriate	Limited	O 0 mSv	O 0 mSv [ped]	5	5	1	0	3	2	4	3	0	1	0

References	Study Quality
17 (7824738)	2
18 (33725187)	3

CT pelvis with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	3	5	2	1	2	1	0	0	0
CT pelvis without IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	6	5	3	0	0	0	0	0	0
CT pelvis without and with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	4	5	3	0	2	0	0	0	0
FDG-PET/CT skull base to mid-thigh	Usually not appropriate	Expert Consensus	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	1	1	13	0	1	0	0	0	0	0	0

Variant 2: Adult female. Adnexal mass, likely benign, no acute symptoms. Premenopausal. Follow-up imaging.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
US pelvis transvaginal	Usually appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	0	0	3	2	9
US pelvis transabdominal and US pelvis transvaginal	Usually appropriate	Strong	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	0	1	1	1	11

References	Study Quality
51 (27282774)	3
24 (11176291)	4
48 (30419104)	3
34 (30303910)	3
22 (31687921)	4
50 (30737137)	2
53 (29112653)	3
55 (29702449)	Good
41 (31549945)	4
45 (32809166)	2
33 (20505067)	4

		46 (20096820)		4												
		47 (21762355)		3												
		49 (23583217)		4												
		52 (21351179)		3												
		54 (35315722)		3												
US duplex Doppler pelvis	Usually appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	8	8	0	0	0	0	1	1	4	3	5
US pelvis transabdominal	Usually appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	7	7	0	0	0	0	0	4	6	3	1
MRI pelvis without IV contrast	May be appropriate	Limited	O 0 mSv	O 0 mSv [ped]	4	4	0	0	1	8	4	1	0	0	0
		References		Study Quality												
		17 (7824738)		2												
		18 (33725187)		3												
MRI pelvis without and with IV contrast	May be appropriate	Limited	O 0 mSv	O 0 mSv [ped]	4	4	1	0	0	8	5	1	1	0	0
		References		Study Quality												
		15 (31482379)		3												
		16 (35040672)		4												
CT pelvis with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	3	8	2	0	0	0	1	0	0
CT pelvis without IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	6	7	1	0	0	0	0	0	0
CT pelvis without and with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	5	6	3	0	0	0	0	0	0
FDG-PET/CT skull base to mid-thigh	Usually not appropriate	Expert Consensus	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	1	1	11	2	1	0	0	0	0	0	0

Variant 3: Adult female. Adnexal mass, likely benign, no acute symptoms. Postmenopausal. Follow-up Imaging.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
US pelvis transvaginal	Usually appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	0	0	3	2	9
US pelvis transabdominal and US pelvis transvaginal	Usually appropriate	Strong	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	0	0	1	2	11

References	Study Quality
58 (12962948)	4
56 (23001924)	3
59 (20503240)	3
48 (30419104)	3
34 (30303910)	3
55 (29702449)	Good
57 (31396641)	3
41 (31549945)	4
46 (20096820)	4
47 (21762355)	3

US duplex Doppler pelvis	Usually appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	8	8	0	0	0	0	2	0	5	1	6
US pelvis transabdominal	Usually appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	7	7	0	0	0	1	0	2	7	3	1
MRI pelvis without and with IV contrast	May be appropriate	Limited	O 0 mSv	O 0 mSv [ped]	6	6	0	0	0	1	4	8	0	1	0

References	Study Quality
15 (31482379)	3
16 (35040672)	4

MRI pelvis without IV contrast	May be appropriate	Limited	O 0 mSv	O 0 mSv [ped]	5	5	0	0	1	6	4	3	0	0	0

References	Study Quality

					17 (7824738)	2										
					18 (33725187)	3										
CT pelvis with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	3	7	1	1	0	2	0	0	0
CT pelvis without IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	6	7	1	0	0	0	0	0	0
CT pelvis without and with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	6	5	2	0	1	0	0	0	0
FDG-PET/CT skull base to mid-thigh	Usually not appropriate	Expert Consensus	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	1	1	10	2	2	0	0	0	0	0	0

Variant 4: Adult female. Adnexal mass, indeterminate on initial pelvic US, no acute symptoms. Premenopausal or postmenopausal. Next imaging study for characterization.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
MRI pelvis without and with IV contrast	Usually appropriate	Strong	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	0	1	1	1	11

References	Study Quality
69 (24183731)	4
73 (22095438)	2
77 (22105841)	3
81 (19214523)	3
79 (21193596)	2
84 (29467113)	3
68 (27770228)	4
85 (27583370)	3
43 (28727500)	3
72 (31977064)	3
82 (33484725)	4

		75 (27367786)	3												
		71 (33929901)	4												
		83 (32715577)	2												
		16 (35040672)	4												
		70 (23468574)	2												
		74 (20419493)	3												
		76 (22814827)	3												
		78 (18581400)	3												
		80 (18972356)	3												
MRI pelvis without IV contrast	May be appropriate	Strong	O 0 mSv	O 0 mSv [ped]	6	6	0	0	0	1	4	10	1	0	0
		References	Study Quality												
		17 (7824738)	2												
		72 (31977064)	3												
		86 (30177450)	4												
		18 (33725187)	3												
		16 (35040672)	4												
		70 (23468574)	2												
CT pelvis with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	4	4	3	0	2	1	0	0	0
CT pelvis without IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	7	3	3	0	1	0	0	0	0
CT pelvis without and with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	4	5	1	0	3	1	0	0	0
FDG-PET/CT skull base to mid-thigh	Usually not appropriate	Limited	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	1	1	9	1	4	0	0	0	0	0	0
		References	Study Quality												
		65 (24463640)	3												
		62 (34112736)	4												

63 (15599527)	4
64 (29399968)	3
66 (18423470)	3
67 (20093590)	4

Variant 5: Adult female. Adnexal mass, indeterminate on initial pelvic US, no acute symptoms. Premenopausal. Follow-up imaging.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
US pelvis transvaginal	Usually appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	9	9	0	0	1	0	0	1	3	2	7
US pelvis transabdominal and US pelvis transvaginal	Usually appropriate	Limited	O 0 mSv	O 0 mSv [ped]	9	9	0	0	1	0	0	0	2	2	9

References	Study Quality
60 (29480762)	3
44 (31770022)	3
72 (31977064)	3
22 (31687921)	4
16 (35040672)	4

MRI pelvis without and with IV contrast	Usually appropriate	Limited	O 0 mSv	O 0 mSv [ped]	8	8	0	0	0	1	1	1	4	5	2

References	Study Quality
16 (35040672)	4

US duplex Doppler pelvis	Usually appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	8	8	0	0	1	0	1	1	3	2	6
MRI pelvis without IV contrast	May be appropriate	Strong	O 0 mSv	O 0 mSv [ped]	6	6	0	0	1	4	1	6	2	0	0

References	Study Quality
17 (7824738)	2
72 (31977064)	3

References	Study Quality
86 (30177450)	4
18 (33725187)	3
16 (35040672)	4
70 (23468574)	2

US pelvis transabdominal	May be appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	6	6	1	0	0	0	2	5	5	1	0
CT pelvis with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	6	3	2	2	1	0	0	0	0
CT pelvis without IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	7	5	2	0	0	0	0	0	0
CT pelvis without and with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	6	3	2	1	2	0	0	0	0
FDG-PET/CT skull base to mid-thigh	Usually not appropriate	Expert Consensus	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	1	1	12	2	0	0	0	0	0	0	0

Variant 6: Adult female. Adnexal mass, indeterminate on initial pelvic US, no acute symptoms. Postmenopausal. Follow-up imaging.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
US duplex Doppler pelvis	Usually appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	2	1	1	3	7
US pelvis transvaginal	Usually appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	0	2	1	4	7
US pelvis transabdominal and US pelvis transvaginal	Usually appropriate	Limited	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	0	1	1	3	9

References	Study Quality
90 (27876070)	3
60 (29480762)	3

		88 (25068555)		3												
		44 (31770022)		3												
		72 (31977064)		3												
		89 (33992598)		3												
		91 (28118296)		4												
		16 (35040672)		4												
MRI pelvis without and with IV contrast	Usually appropriate	Limited	O 0 mSv	O 0 mSv [ped]	8	8	0	0	0	0	1	0	6	3	4
		References		Study Quality												
		16 (35040672)		4												
		87 (25273949)		3												
US pelvis transabdominal	Usually appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	7	7	0	0	0	1	1	5	5	0	2
MRI pelvis without IV contrast	May be appropriate	Strong	O 0 mSv	O 0 mSv [ped]	6	6	0	0	0	0	3	11	2	0	0
		References		Study Quality												
		17 (7824738)		2												
		72 (31977064)		3												
		86 (30177450)		4												
		18 (33725187)		3												
		16 (35040672)		4												
		70 (23468574)		2												
CT pelvis with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	4	4	3	1	1	0	1	0	0
CT pelvis without IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	7	4	3	0	0	0	0	0	0
CT pelvis without and with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	2	2	5	3	2	0	3	0	1	0	0

FDG-PET/CT skull base to mid-thigh	Usually not appropriate	Expert Consensus	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	1	1	10	2	1	0	0	1	0	0	0

Variant 7: Adult female. Adnexal mass, suspicious for malignancy on pelvic US, no acute symptoms. Premenopausal or postmenopausal. Next imaging study for characterization.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
MRI pelvis without and with IV contrast	Usually appropriate	Strong	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	0	1	2	0	13

References	Study Quality
61 (21156740)	3
73 (22095438)	2
77 (22105841)	3
23 (18504770)	3
79 (21193596)	2
60 (29480762)	3
68 (27770228)	4
36 (25320247)	3
44 (31770022)	3
72 (31977064)	3
22 (31687921)	4
75 (27367786)	3
16 (35040672)	4
33 (20505067)	4
54 (35315722)	3
70 (23468574)	2
74 (20419493)	3
76 (22814827)	3
78 (18581400)	3

		94 (35230183)		3												
MRI pelvis without IV contrast	May be appropriate	Strong	O 0 mSv	O 0 mSv [ped]	6	6	0	0	1	2	2	6	4	1	0
		References		Study Quality												
		17 (7824738)		2												
		72 (31977064)		3												
		86 (30177450)		4												
		18 (33725187)		3												
		16 (35040672)		4												
		70 (23468574)		2												
CT pelvis without and with IV contrast	May be appropriate (Disagreement)	Expert Opinion	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	5	5	5	1	7	0	0	1	0	0	2
CT pelvis with IV contrast	Usually not appropriate	Limited	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	3	3	4	4	6	0	0	0	1	1	0
		References		Study Quality												
		92 (29724422)		4												
FDG-PET/CT skull base to mid-thigh	Usually not appropriate	Limited	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	3	3	7	0	5	0	3	1	0	0	0
		References		Study Quality												
		93 (23988417)		2												
CT pelvis without IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	1	1	10	2	2	0	1	0	1	0	0

Variant 8: Female. Clinically suspected adnexal mass, no acute symptoms. Pregnant. Initial imaging.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
US pelvis transvaginal	Usually appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	0	1	2	4	7

US pelvis transabdominal and US pelvis transvaginal	Usually appropriate	Strong	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	0	0	2	2	10

References	Study Quality
108 (26482730)	3
109 (24496773)	4
100 (9315194)	4
110 (27484484)	4
103 (33515746)	Good
28 (27776067)	4
101 (32347749)	4
102 (10214838)	3
104 (21194826)	4
105 (16310839)	4
106 (15863550)	3
107 (10411786)	4
111 (11127014)	4
112 (33671023)	3
113 (34902893)	4

US duplex Doppler pelvis	Usually appropriate	Strong	O 0 mSv	O 0 mSv [ped]	8	8	0	0	0	0	1	0	6	3	4

References	Study Quality
19 (9618853)	3
23 (18504770)	3
21 (9646799)	2
20 (10739694)	3
25 (7839975)	3
24 (11176291)	4
22 (31687921)	4

US pelvis transabdominal	Usually appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	7	7	0	0	0	0	1	2	5	5	1

MRI pelvis without IV contrast	May be appropriate	Limited	O 0 mSv	O 0 mSv [ped]	5	5	0	0	1	6	5	3	1	0	0

References	Study Quality
17 (7824738)	2
95 (-3188538)	4
18 (33725187)	3
15 (31482379)	3
16 (35040672)	4

MRI pelvis without and with IV contrast	Usually not appropriate	Limited	O 0 mSv	O 0 mSv [ped]	3	3	5	1	5	1	1	0	1	0	0

References	Study Quality
97 (27599330)	3
99 (28668416)	2
95 (-3188538)	4
96 (22003244)	3
98 (-3195845)	4

CT pelvis with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	1	1	11	3	0	0	0	0	0	0	0
CT pelvis without IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	1	1	11	3	0	0	0	0	0	0	0
CT pelvis without and with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	1	1	12	2	0	0	0	0	0	0	0
FDG-PET/CT skull base to mid-thigh	Usually not appropriate	Expert Consensus	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	1	1	14	0	0	0	0	0	0	0	0

Variant 9: Female . Adnexal mass, indeterminate on initial pelvic US, no acute symptoms. Pregnant. Next imaging study for characterization.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
MRI pelvis without IV contrast	Usually appropriate	Strong	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	0	0	4	2	8

References	Study Quality
61 (21156740)	3
108 (26482730)	3
110 (27484484)	4
60 (29480762)	3
95 (-3188538)	4
72 (31977064)	3
86 (30177450)	4
18 (33725187)	3
16 (35040672)	4
70 (23468574)	2
114 (27130065)	3

MRI pelvis without and with IV contrast	Usually not appropriate	Limited	O 0 mSv	O 0 mSv [ped]	3	3	1	0	8	5	1	0	1	0	0

References	Study Quality
97 (27599330)	3
99 (28668416)	2
95 (-3188538)	4
96 (22003244)	3
98 (-3195845)	4

CT pelvis with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	1	1	10	3	0	0	0	1	0	0	0
CT pelvis without IV contrast	Usually not appropriate	Expert Consensus	☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	1	1	10	3	0	1	0	0	0	0	0
CT pelvis without and with IV contrast	Usually not appropriate	Expert Consensus	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	1	1	11	2	0	0	0	1	0	0	0

FDG-PET/CT skull base to mid-thigh	Usually not appropriate	Expert Consensus	☢☢☢☢ 10-30 mSv	☢☢☢☢ 3-10 mSv [ped]	1	1	13	0	1	0	0	0	0	0	0

Appendix Key

A more complete discussion of the items presented below can be found by accessing the supporting documents at the designated hyperlinks.

Appropriateness Category: The panel's recommendation for a procedure based on the assessment of the risks and benefits of performing the procedure for the specified clinical scenario.

SOE: Strength of Evidence. The assessment of the amount and quality of evidence found in the peer reviewed medical literature for an appropriateness recommendation.

- **References:** The citation number and PMID for the reference(s) associated with the recommendation.
- **Study Quality**: The assessment of the quality of an individual reference based on the number of study quality elements described in the reference.

RRL: Relative Radiation Level. A population based assessment of the amount of radiation a typical patient may be exposed to during the specified procedure.

Rating: The final rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

Median: The median rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

Final tabulations: A histogram showing the number of panel members who rated the procedure as noted in the column heading (ie, 1, 2, 3, etc.).

Additional supporting documents about the AC methodology and processes can be found at **www.acr.org/ac.**