American College of Radiology ACR Appropriateness Criteria®

Malignant or Aggressive Primary Musculoskeletal Tumor-Staging And Surveillance

Variant 1: Malignant or aggressive primary musculoskeletal tumor. Initial staging. Evaluation for pulmonary metastasis.

D .	Appropriateness	COF	4.1.4. DDT	D 1 DD1		3.5.31			F	inal '	Tabu	latio	ns		
Procedure	Category	SOE	Adults RRL	Peds RRL	Rating	Median	1	2	3	4	5	6	7	8	9
CT chest without IV contrast	Usually appropriate	Strong	��� 1-10 mSv	���� 3- 10 mSv [ped]	9	9	0	0	0	0	0	1	1	2	13
		References		Study	Quality										
		9 (26472138)			2										
		11 (16410133)			2										
		12 (21513852)			4		_								
FDG-PET/CT whole body	May be appropriate	Strong	���� 10-30 mSv	0	5	5	0	0	0	2	10	4	0	0	0
		References		Study	Quality										
		16 (22745027)			2										
		6 (30953476)		C	Good										
		13 (19352275)			3										
		15 (22134535)			3										
		14 (23251096)			3										
CT chest with IV contrast	Usually not appropriate	Expert Consensus	��� 1-10 mSv	❤️❤️❤️ 3- 10 mSv [ped]	1	1	10	0	6	0	0	0	0	0	0
CT chest without and with IV contrast	Usually not appropriate	Expert Consensus	��� 1-10 mSv	���� 3- 10 mSv [ped]	1	1	10	1	4	0	0	1	0	0	0

Radiography chest	Usually approp		Expert Consensus	⊕ <0.1 mSv		1	1	14	0	1	0	0	1	0	0	0
FDG-PET/MRI whole body	Usually approp	4	Limited	≎≎≎ 1-10 mSv	���� 3- 10 mSv [ped]	1	1	10	1	3	1	1	1	0	0	0
			References		Study	Quality										
			17 (27951459)			3										

Variant 2: Malignant or aggressive primary musculoskeletal tumor. Initial staging. Evaluation for extrapulmonary metastasis.

D 1	Appropri	ateness	COF	4 1 14 DD		D I DDI	D (1	37.11			F	inal	Tabu	latio	ns		
Procedure	Categ	ory	SOE	Adults RR	L	Peds RRL	Rating	Median	1	2	3	4	5	6	7	8	9
FDG-PET/CT whole body	Usua approp		Strong	⊕⊕⊕⊕ 10-3 mSv	30	���� 3- 10 mSv [ped]	8	8	0	0	0	0	2	0	4	4	7
			References			Study	Quality										
			37 (18048826)				3										
			34 (16951929)				2										
			36 (21716019)				2										
			33 (11418435)				2										
			22 (25539251)				3										
			32 (14693872)				2										
			30 (31601386)				2										
			35 (31016339)				2										
			6 (30953476)			G	lood										
			31 (29189489)				4										
			15 (22134535)				3										
MRI whole body without IV contrast	May approp (Disagree	riate	Expert Opinion	O 0 mSv		O 0 mSv [ped]	5	5	1	0	1	1	10	0	0	2	1
			References			Study	Quality										
			34 (16951929)				2										

			33 (11418435)			2										
			32 (14693872)			2										
			29 (30915559)			4										
			42 (31447287)			4										
			35 (31016339)			2										
			44 (26831007)			3										
			40 (29558901)			4										
			41 (29275455)			4										
			43 (20369068)			4										
Bone scan whole body with SPECT or SPECT/CT area of interest	May approp		Limited	��� 1-10 mSv	≎≎≎≎ 3- 10 mSv [ped]	5	5	0	0	2	1	7	6	0	0	0
	•		References		Study	Quality		•			•		•			
			30 (31601386)			2										
Fluoride PET/CT whole body	May approp (Disagre	riate	Expert Opinion	���� 10-30 mSv	���� 3- 10 mSv [ped]	5	5	3	0	4	2	4	3	0	0	0
			References		Study	Quality										
			39 (28798033)			2										
FDG-PET/MRI whole body	May approp		Limited	≎	���� 3- 10 mSv [ped]	5	5	0	0	1	3	11	0	1	0	0
			References		Study	Quality										
			38 (24309383)			3										
			17 (27951459)			3										
Bone scan whole body	May approp		Strong	≎ ≎≎ 1-10 mSv	���� 3- 10 mSv [ped]	4	4	0	0	2	7	5	1	0	0	1
			References		Study	Quality										
			22 (25539251)			3										
			28 (20224457)			2										
			23 (25533313)			2										
			29 (30915559)			4										
			24 (23995264)			3										
		·	·	·	·	·	· · · · · · · · · · · · · · · · · · ·									

25 (9779715)	3
26 (12750613)	3
27 (19542943)	4

MRI whole body without and with IV contrast	Usually not appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	2	2	8	1	5	2	0	0	0	0	0
CT area of interest with IV contrast	Usually not appropriate	Expert Consensus	Varies	Varies	1	1	11	0	4	1	0	0	0	0	0
CT area of interest without IV contrast	Usually not appropriate	Expert Consensus	Varies	Varies	1	1	11	0	5	0	0	0	0	0	0
CT area of interest without and with IV contrast	Usually not appropriate	Expert Consensus	Varies	Varies	1	1	11	0	5	0	0	0	0	0	0
US area of interest	Usually not appropriate	Expert Consensus	O 0 mSv	O 0 mSv [ped]	1	1	14	1	0	0	1	0	0	0	0
Radiography area of interest	Usually not appropriate	Expert Consensus	Varies	Varies	1	1	14	1	0	0	0	0	0	0	1

Variant 3: Malignant or aggressive primary musculoskeletal tumor with no suspected or known recurrence. Surveillance for pulmonary metastasis.

D 1	Appropriateness	COF	A L I/ DDI	D I DDI	D 4	3.7.19			F	inal '	Гаbu	latio	ns		
Procedure	Category	SOE	Adults RRL	Peds RRL	Rating	Median	1	2	3	4	5	6	7	8	9
CT chest without IV contrast	Usually appropriate	Strong	��� 1-10 mSv	���� 3- 10 mSv [ped]	9	9	0	0	0	1	0	0	1	2	13

References	Study Quality
56 (21537873)	3
55 (17438474)	3
9 (26472138)	2
45 (31377411)	4
54 (24249538)	2
12 (21513852)	4

			53 (29437071)			2										
FDG-PET/CT whole body	May approp	be riate	Limited	≎≎≎≎ 10-3 mSv	0	5	5	0	0	0	1	13	1	1	0	0
			References		Study	y Quality										
			16 (22745027)			2										
			15 (22134535)			3										
CT chest with IV contrast	Usually approp		Expert Consensus	≎≎≎ 1-10 mSv	���� 3- 10 mSv [ped]	1	1	11	0	2	2	0	1	0	0	0
CT chest without and with IV contrast	Usually approp		Expert Consensus	≎≎≎ 1-10 mSv	���� 3- 10 mSv [ped]	1	1	12	1	2	1	0	0	0	0	0
Radiography chest	Usually approp		Limited	≎ <0.1 mS	v <0.03 mSv [ped]	1	1	10	3	3	0	0	0	0	0	0
			References		Study	y Quality										
			57 (10674442)			3										
FDG-PET/MRI whole body	Usually approp		Expert Consensus	��� 1-10 mSv	���� 3- 10 mSv [ped]	1	1	10	0	4	2	0	1	0	0	0

Variant 4: Malignant or aggressive primary bone tumor. Surveillance for local recurrence.

D 1	Appropria	teness	COF	414 DD		D I DDI	D 41	3.6.11			F	inal '	Tabu	latio	ns		
Procedure	Catego		SOE	Adults RR	L	Peds RRL	Rating	Median	1	2	3	4	5	6	7	8	9
MRI area of interest without and with IV contrast	Usuall appropri		Limited	O 0 mSv	,	O 0 mSv [ped]	9	9	0	0	0	0	0	0	1	1	15
			References			Study	Quality										
			67 (23246334)				3										
			68 (9285993)				4										
Radiography area of interest	Usuall appropri	.*	Expert Consensus	Varies		Varies	9	9	1	0	0	0	0	2	1	1	12

MRI area of interest without IV contrast	Usually appropria	Expert Consensus	O 0 mSv	/	O 0 mSv [ped]	7	7	1	0	1	0	2	3	6	3	1
CT area of interest with IV contrast	May be appropria	Expert Consensus	Varies		Varies	5	5	0	0	1	1	10	3	0	1	0
FDG-PET/CT whole body	May be appropria	Limited	≎≎≎≎ 10- mSv	-30	���� 3- 10 mSv [ped]	5	5	0	0	0	0	9	2	4	1	0
		References			Study	Quality										
		64 (23558689)				3										
		65 (29261626)				3										
		66 (25431093)				2										
Bone scan whole body with SPECT or SPECT/CT area of interest	Usually n appropria	Expert Consensus	431093) ert ��� 1-10		���� 3- 10 mSv [ped]	2	2	8	3	2	2	1	0	1	0	0
Fluoride PET/CT whole body	Usually n appropria	Expert Consensus			���� 3- 10 mSv [ped]	2	2	8	2	3	1	1	1	1	0	0
CT area of interest without IV contrast	Usually n appropria	Expert Consensus	Varies		Varies	1	1	9	0	2	2	3	0	0	0	0
CT area of interest without and with IV contrast	Usually n appropria	Expert Consensus	Varies		Varies	1	1	9	1	2	2	1	1	0	0	0
Bone scan whole body	Usually n appropria	Limited	��� 1-1 mSv	0	���� 3- 10 mSv [ped]	1	1	10	2	4	0	0	0	0	0	0
		References			Study	Quality										
		22 (25539251)				3										
		 23 (25533313)		<u> </u>		2										
US area of interest	Usually n appropria	Expert Consensus	O 0 mSv	/	O 0 mSv [ped]	1	1	10	2	1	2	1	1	0	0	0
FDG-PET/MRI whole body	Usually n appropria	Expert Consensus	��� 1-1 mSv	0	���� 3- 10 mSv [ped]	1	1	11	3	2	0	0	0	0	0	0

Variant 5: Malignant or aggressive primary soft tissue tumor. Surveillance for local recurrence.

	Appropriatenes	S GOT		p 1 ppr	- ·	3.5.11			F	inal	Tabu	latio	ns		
Procedure	Category	SOE	Adults RRL	Peds RRL	Rating	Median	1	2	3	4	5	6	7	8	9
MRI area of interest without and with IV contrast	Usually appropriate	Limited	O 0 mSv	O 0 mSv [ped]	9	9	0	0	0	0	0	0	0	1	16
		References		Study	y Quality		•		•		•		•		
		82 (27060189)			3										
		77 (30352764))		3										
		83 (19744809))		3										
		67 (23246334))		3										
		68 (9285993)			4										
MRI area of interest without IV contrast	Usually appropriate	Limited	O 0 mSv	O 0 mSv [ped]	7	7	0	0	0	1	2	0	11	2	0
		References		Study	y Quality										
		82 (27060189))		3										
		77 (30352764))		3										
		83 (19744809))		3										
		67 (23246334))		3										
		68 (9285993)			4						1				
CT area of interest with IV contrast	May be appropriate	Expert Consensus	Varies	Varies	6	6	0	0	0	0	7	4	4	1	1
FDG-PET/CT whole body	May be appropriate	Strong	���� 10-3 mSv	0 \$9993- 10 mSv [ped]	5	5	0	0	0	1	9	4	2	0	0
		References		Study	y Quality										
		79 (23233156))		2										
		80 (27488833))		2										

US area of interest	May be appropriate (Disagreement)		Expert Opinion	O 0 mSv	O 0 mSv [ped]	5	5	2	0	1	3	7	3	0	0	0
			References		Study Quality											
			84 (25794664)		3											
FDG-PET/MRI whole body	May be appropriate		Moderate	��� 1-10 mSv	���� 3- 10 mSv [ped]	5	5	1	0	1	5	7	1	1	0	0
	References				Study Quality											
				81 (28567495)												
CT area of interest without and with IV contrast	Usually not appropriate		Expert Consensus	Varies	Varies	2	2	8	3	2	2	1	0	0	0	0
CT area of interest without IV contrast	Usually not appropriate		Expert Consensus	Varies	Varies	1	1	11	1	1	2	1	0	0	0	0
Radiography area of interest	Usually not appropriate		Expert Consensus	Varies	Varies	1	1	10	1	0	3	1	1	0	0	0

Appendix Key

A more complete discussion of the items presented below can be found by accessing the supporting documents at the designated hyperlinks.

Appropriateness Category: The panel's recommendation for a procedure based on the assessment of the risks and benefits of performing the procedure for the specified clinical scenario.

SOE: Strength of Evidence. The assessment of the amount and quality of evidence found in the peer reviewed medical literature for an appropriateness recommendation.

- **References:** The citation number and PMID for the reference(s) associated with the recommendation.
- Study Quality: The assessment of the quality of an individual reference based on the number of study quality elements described in the reference.

RRL: Relative Radiation Level. A population based assessment of the amount of radiation a typical patient may be exposed to during the specified procedure.

Rating: The final rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

Median: The median rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

Final tabulations: A histogram showing the number of panel members who rated the procedure as noted in the column heading (ie, 1, 2, 3, etc.).

Additional supporting documents about the AC methodology and processes can be found at www.acr.org/ac.