

CT complete spine with IV contrast	Usually not appropriate	Expert Consensus	☹☹☹☹ 10-30 mSv	☹☹☹☹ 3-10 mSv [ped]	1	1	14	2	1	0	0	0	1	0	0
------------------------------------	-------------------------	------------------	----------------	---------------------	---	---	----	---	---	---	---	---	---	---	---

Variant 2: Child. Acute ataxia, history of recent trauma. Initial imaging.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations									
							1	2	3	4	5	6	7	8	9	
CT head without IV contrast	Usually appropriate	Limited	☹☹☹ 1-10 mSv	☹☹☹ 0.3-3 mSv [ped]	9	9	0	0	0	0	1	0	1	2	14	
		References	Study Quality													
		7 (27789117)	4													
MRI head without IV contrast	Usually appropriate	Limited	○ 0 mSv	○ 0 mSv [ped]	7	7	0	0	1	0	4	4	3	2	4	
		References	Study Quality													
		4 (22595899)	4													
		26 (25535060)	2													
		33 (21292802)	4													
		32 (19863638)	4													
CTA head and neck with IV contrast	May be appropriate	Expert Consensus	☹☹☹ 1-10 mSv	☹☹☹☹ 3-10 mSv [ped]	6	6	0	0	0	1	6	4	8	0	0	
CTA head with IV contrast	May be appropriate	Expert Consensus	☹☹☹ 1-10 mSv	☹☹☹☹ 3-10 mSv [ped]	5	5	0	0	0	0	15	3	1	0	0	
CTA neck with IV contrast	May be appropriate	Expert Consensus	☹☹☹ 1-10 mSv	☹☹☹ 0.3-3 mSv [ped]	5	5	0	0	0	0	15	3	1	0	0	
MRA head without IV contrast	May be appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	5	5	0	0	0	0	18	1	0	0	0	
MRA head and neck without IV contrast	May be appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	5	5	0	0	0	0	17	2	0	0	0	

MRA neck without IV contrast	May be appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	5	5	0	0	0	0	18	1	0	0	0
MRA neck with IV contrast	May be appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	5	5	0	0	0	5	13	1	0	0	0
MRI head without and with IV contrast	Usually not appropriate	Limited	0 0 mSv	0 0 mSv [ped]	3	3	4	4	4	4	2	0	1	0	0
		References	Study Quality												
		4 (22595899)	4												
		26 (25535060)	2												
		33 (21292802)	4												
		32 (19863638)	4												
CT head with IV contrast	Usually not appropriate	Limited	☹☹☹ 1-10 mSv	☹☹☹ 0.3-3 mSv [ped]	2	2	7	4	3	0	3	1	0	0	0
		References	Study Quality												
		7 (27789117)	4												
CT head without and with IV contrast	Usually not appropriate	Limited	☹☹☹ 1-10 mSv	☹☹☹☹ 3-10 mSv [ped]	2	2	9	2	3	0	3	1	0	0	0
		References	Study Quality												
		7 (27789117)	4												
MRI complete spine without IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	2	2	7	4	8	0	0	0	0	0	0
MRI complete spine without and with IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	2	2	9	0	4	0	3	1	1	0	0
MRV head without IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	2	2	9	10	0	0	0	0	0	0	0
CTV head with IV contrast	Usually not appropriate	Expert Consensus	☹☹☹ 1-10 mSv	☹☹☹☹ 3-10 mSv [ped]	2	2	9	3	1	2	1	2	0	0	0
MRV head and neck without IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	2	2	7	12	0	0	0	0	0	0	0

MR spectroscopy head without IV contrast	May be appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	5	5	1	0	1	1	14	2	0	0	0	
CTA neck with IV contrast	Usually not appropriate	Expert Consensus	☼☼☼ 1-10 mSv	☼☼☼ 0.3-3 mSv [ped]	3	3	8	1	2	2	3	1	1	0	0	
CTA head and neck with IV contrast	Usually not appropriate	Expert Consensus	☼☼☼ 1-10 mSv	☼☼☼☼ 3-10 mSv [ped]	3	3	7	1	4	2	2	1	1	0	0	
CT head without IV contrast	Usually not appropriate	Expert Consensus	☼☼☼ 1-10 mSv	☼☼☼ 0.3-3 mSv [ped]	2	2	6	11	1	0	0	0	1	0	0	
MRA head without IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	2	2	7	9	3	0	0	0	0	0	0	
MRA head and neck without IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	2	2	6	5	8	0	0	0	0	0	0	
MRA neck without IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	2	2	7	4	3	4	1	0	0	0	0	
MRA neck with IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	2	2	7	8	3	0	1	0	0	0	0	
CTA head with IV contrast	Usually not appropriate	Expert Consensus	☼☼☼ 1-10 mSv	☼☼☼☼ 3-10 mSv [ped]	1	1	10	2	2	1	1	1	1	1	0	0
CT head with IV contrast	Usually not appropriate	Expert Consensus	☼☼☼ 1-10 mSv	☼☼☼ 0.3-3 mSv [ped]	1	1	14	1	1	1	0	0	1	0	0	
CT head without and with IV contrast	Usually not appropriate	Expert Consensus	☼☼☼ 1-10 mSv	☼☼☼☼ 3-10 mSv [ped]	1	1	14	0	1	0	3	0	0	0	0	
MRI complete spine without IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	1	1	14	4	0	0	1	0	0	0	0	
MRI complete spine without and with IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	1	1	12	1	1	1	2	1	0	0	0	
MRV head without IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	1	1	11	1	2	0	2	2	0	0	0	

CTV head with IV contrast	Usually not appropriate	Expert Consensus	☹☹☹ 1-10 mSv	☹☹☹☹ 3-10 mSv [ped]	1	1	12	2	1	1	1	1	0	0	0
MRV head without and with IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	18	1	0	0	0	0	0	0	0
MRV head and neck without IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	10	3	1	0	2	1	1	0	0
MRV head with IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	17	2	0	0	0	0	0	0	0
CT complete spine without and with IV contrast	Usually not appropriate	Expert Consensus	☹☹☹☹ 10-30 mSv	☹☹☹☹ 3-10 mSv [ped]	1	1	17	0	1	0	0	0	0	0	0
CT complete spine without IV contrast	Usually not appropriate	Expert Consensus	☹☹☹☹ 10-30 mSv	☹☹☹☹ 3-10 mSv [ped]	1	1	16	0	2	0	0	0	0	0	0
CT complete spine with IV contrast	Usually not appropriate	Expert Consensus	☹☹☹☹ 10-30 mSv	☹☹☹☹ 3-10 mSv [ped]	1	1	15	1	2	0	0	0	0	0	0

Variant 4: Child. Chronic progressive ataxia. Initial imaging.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
MRI head without IV contrast	Usually appropriate	Strong	○ 0 mSv	○ 0 mSv [ped]	8	8	0	0	0	0	1	0	3	6	8

References	Study Quality
41 (21701895)	3
46 (19631275)	2
45 (22764178)	4
4 (22595899)	4
39 (22816526)	4
42 (29922574)	3

1 (28257643)	4
44 (27126632)	4
40 (20378176)	4
47 (30086112)	4
43 (27423800)	4
11 (27142713)	2
28 (26444039)	4
5 (23622410)	4
25 (22764177)	4

MRI head without and with IV contrast	Usually appropriate	Strong	0 0 mSv	0 0 mSv [ped]	8	8	0	0	0	0	2	1	3	5	7
---------------------------------------	---------------------	--------	---------	---------------	---	---	---	---	---	---	---	---	---	---	---

References	Study Quality
41 (21701895)	3
46 (19631275)	2
45 (22764178)	4
4 (22595899)	4
39 (22816526)	4
42 (29922574)	3
1 (28257643)	4
44 (27126632)	4
40 (20378176)	4
47 (30086112)	4
43 (27423800)	4
11 (27142713)	2
28 (26444039)	4
5 (23622410)	4
25 (22764177)	4

CT head without IV contrast	May be appropriate	Limited	☼☼☼ 1-10 mSv	☼☼☼ 0.3-3 mSv [ped]	5	5	1	0	0	2	13	2	1	0	0
-----------------------------	--------------------	---------	--------------	---------------------	---	---	---	---	---	---	----	---	---	---	---

References	Study Quality
14 (26940752)	4

			28 (26444039)		4													
MRI complete spine without IV contrast	May be appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	5	5	1	0	0	1	13	3	1	0	0			
MRI complete spine without and with IV contrast	May be appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	5	5	1	0	0	0	15	3	0	0	0			
CT head with IV contrast	Usually not appropriate	Limited	⊗⊗⊗ 1-10 mSv	⊗⊗⊗ 0.3-3 mSv [ped]	3	3	6	3	2	2	3	2	0	0	0			
			References		Study Quality													
			14 (26940752)		4													
			28 (26444039)		4													
CT head without and with IV contrast	Usually not appropriate	Limited	⊗⊗⊗ 1-10 mSv	⊗⊗⊗⊗ 3-10 mSv [ped]	2	2	8	2	1	1	4	2	0	0	0			
			References		Study Quality													
			14 (26940752)		4													
			28 (26444039)		4													
MRA head without IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	2	2	8	3	1	2	2	2	0	0	0			
MRA head and neck without IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	2	2	8	4	1	1	2	2	0	0	0			
MRA neck without IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	2	2	9	3	1	1	2	2	0	0	0			
MR spectroscopy head without IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	2	2	7	3	2	0	7	0	0	0	0			
CTA head with IV contrast	Usually not appropriate	Expert Consensus	⊗⊗⊗ 1-10 mSv	⊗⊗⊗⊗ 3-10 mSv [ped]	1	1	11	1	1	2	2	1	0	0	0			
CTA neck with IV contrast	Usually not appropriate	Expert Consensus	⊗⊗⊗ 1-10 mSv	⊗⊗⊗ 0.3-3 mSv [ped]	1	1	11	1	1	2	2	1	0	0	0			

CTA head and neck with IV contrast	Usually not appropriate	Expert Consensus	⊗⊗⊗ 1-10 mSv	⊗⊗⊗⊗ 3-10 mSv [ped]	1	1	11	1	1	2	2	1	0	0	0
MRV head without IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	11	3	0	0	2	2	0	0	0
CTV head with IV contrast	Usually not appropriate	Expert Consensus	⊗⊗⊗ 1-10 mSv	⊗⊗⊗⊗ 3-10 mSv [ped]	1	1	13	1	0	2	2	0	0	0	0
MRV head without and with IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	13	2	0	0	1	2	0	0	0
MRV head and neck without IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	10	3	0	1	3	1	0	0	0
MRA neck with IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	11	3	1	0	2	1	0	0	0
MRV head with IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	13	2	0	0	1	2	0	0	0
CT complete spine without and with IV contrast	Usually not appropriate	Expert Consensus	⊗⊗⊗⊗ 10-30 mSv	⊗⊗⊗⊗ 3-10 mSv [ped]	1	1	17	0	1	0	0	0	0	0	0
CT complete spine without IV contrast	Usually not appropriate	Expert Consensus	⊗⊗⊗⊗ 10-30 mSv	⊗⊗⊗⊗ 3-10 mSv [ped]	1	1	15	0	2	0	1	0	0	0	0
CT complete spine with IV contrast	Usually not appropriate	Expert Consensus	⊗⊗⊗⊗ 10-30 mSv	⊗⊗⊗⊗ 3-10 mSv [ped]	1	1	15	1	2	0	0	0	0	0	0

Variant 5: Child. Chronic nonprogressive ataxia. Initial imaging.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
MRI head without IV contrast	Usually appropriate	Limited	○ 0 mSv	○ 0 mSv [ped]	9	9	0	0	0	1	0	0	1	3	13
			References	Study Quality											

4 (22595899)	4
52 (27785688)	3
43 (27423800)	4
6 (26413479)	3
32 (19863638)	4
50 (26132530)	4

CT head without IV contrast	May be appropriate	Limited	☼☼☼ 1-10 mSv	☼☼☼ 0.3-3 mSv [ped]	6	6	0	0	0	2	7	9	1	0	0
-----------------------------	--------------------	---------	--------------	---------------------	---	---	---	---	---	---	---	---	---	---	---

References	Study Quality
14 (26940752)	4
28 (26444039)	4

MRI complete spine without IV contrast	May be appropriate	Limited	○ 0 mSv	○ 0 mSv [ped]	5	5	1	0	0	0	16	2	0	0	0
--	--------------------	---------	---------	---------------	---	---	---	---	---	---	----	---	---	---	---

References	Study Quality
44 (27126632)	4

CT head with IV contrast	Usually not appropriate	Limited	☼☼☼ 1-10 mSv	☼☼☼ 0.3-3 mSv [ped]	3	3	7	2	2	3	4	0	0	0	0
--------------------------	-------------------------	---------	--------------	---------------------	---	---	---	---	---	---	---	---	---	---	---

References	Study Quality
14 (26940752)	4
28 (26444039)	4

MRI head without and with IV contrast	Usually not appropriate	Limited	○ 0 mSv	○ 0 mSv [ped]	2	2	5	5	2	0	6	0	1	0	0
---------------------------------------	-------------------------	---------	---------	---------------	---	---	---	---	---	---	---	---	---	---	---

References	Study Quality
4 (22595899)	4
52 (27785688)	3
43 (27423800)	4
6 (26413479)	3
32 (19863638)	4
50 (26132530)	4

MRI complete spine without and with IV contrast	Usually not appropriate	Limited	0 0 mSv	0 0 mSv [ped]	2	2	5	5	3	5	1	0	0	0	0
		References		Study Quality											
		44 (27126632)		4											
CTA head with IV contrast	Usually not appropriate	Expert Consensus	☼☼☼ 1-10 mSv	☼☼☼☼ 3-10 mSv [ped]	1	1	11	3	1	1	1	1	0	0	0
CTA neck with IV contrast	Usually not appropriate	Expert Consensus	☼☼☼ 1-10 mSv	☼☼☼ 0.3-3 mSv [ped]	1	1	11	2	2	1	1	1	0	0	0
CT head without and with IV contrast	Usually not appropriate	Limited	☼☼☼ 1-10 mSv	☼☼☼☼ 3-10 mSv [ped]	1	1	10	2	2	2	2	0	0	0	0
		References		Study Quality											
		14 (26940752)		4											
		28 (26444039)		4											
CTA head and neck with IV contrast	Usually not appropriate	Expert Consensus	☼☼☼ 1-10 mSv	☼☼☼☼ 3-10 mSv [ped]	1	1	11	3	1	0	2	1	0	0	0
MRA head without IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	1	1	10	3	1	1	2	1	0	0	0
MRA head and neck without IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	1	1	10	3	1	0	2	2	0	0	0
MRA neck without IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	1	1	10	3	1	1	2	1	0	0	0
MR spectroscopy head without IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	1	1	10	2	3	2	0	0	0	1	0
MRV head without IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	1	1	11	3	0	1	2	1	0	0	0
CTV head with IV contrast	Usually not appropriate	Expert Consensus	☼☼☼ 1-10 mSv	☼☼☼☼ 3-10 mSv [ped]	1	1	13	1	1	0	2	1	0	0	0

MRV head without and with IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	13	2	0	0	1	2	0	0	0
MRV head and neck without IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	11	3	0	0	3	1	0	0	0
MRA neck with IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	11	3	1	0	2	1	0	0	0
MRV head with IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	12	3	0	0	1	2	0	0	0
CT complete spine without and with IV contrast	Usually not appropriate	Expert Consensus	⊗⊗⊗⊗ 10-30 mSv	⊗⊗⊗⊗ 3-10 mSv [ped]	1	1	17	1	0	0	0	0	0	0	0
CT complete spine without IV contrast	Usually not appropriate	Expert Consensus	⊗⊗⊗⊗ 10-30 mSv	⊗⊗⊗⊗ 3-10 mSv [ped]	1	1	15	1	1	0	1	0	0	0	0
CT complete spine with IV contrast	Usually not appropriate	Expert Consensus	⊗⊗⊗⊗ 10-30 mSv	⊗⊗⊗⊗ 3-10 mSv [ped]	1	1	16	2	0	0	0	0	0	0	0

Appendix Key

A more complete discussion of the items presented below can be found by accessing the supporting documents at the designated hyperlinks.

Appropriateness Category: The panel's recommendation for a procedure based on the assessment of the risks and benefits of performing the procedure for the specified clinical scenario.

SOE: Strength of Evidence. The assessment of the amount and quality of evidence found in the peer reviewed medical literature for an appropriateness recommendation.

- **References:** The citation number and PMID for the reference(s) associated with the recommendation.
- **Study Quality:** The assessment of the quality of an individual reference based on the number of study quality elements described in the reference.

RRL: Relative Radiation Level. A population based assessment of the amount of radiation a typical patient may be exposed to during the specified procedure.

Rating: The final rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

Median: The median rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

Final tabulations: A histogram showing the number of panel members who rated the procedure as noted in the column heading (ie, 1, 2, 3, etc.).

Additional supporting documents about the AC methodology and processes can be found at www.acr.org/ac.