



28 (25241315)	4
31 (10887229)	M
27 (12239057)	3
32 (16864668)	1
25 (9052573)	4
29 (25079487)	3

CTA abdomen and pelvis with bilateral lower extremity runoff without and with IV contrast	Usually appropriate	Strong	☼☼☼☼☼ 30-100 mSv		8	8	0	0	0	0	1	1	4	9	1
---	---------------------	--------	------------------	--	---	---	---	---	---	---	---	---	---	---	---

References	Study Quality
21 (21345629)	2
23 (27399244)	4
15 (21658691)	2
22 (16966479)	4
18 (16580545)	2
10 (12591682)	2
19 (17848679)	M
20 (12601190)	1
13 (20609871)	4
17 (15128997)	2
16 (18094974)	2
14 (20233780)	4
24 (17709077)	4

Arteriography lower extremity	Usually appropriate	Limited	☼☼ 0.1-1mSv	☼☼☼ 0.3-3 mSv [ped]	7	7	0	0	1	0	2	1	9	4	0
-------------------------------	---------------------	---------	-------------	---------------------	---	---	---	---	---	---	---	---	---	---	---

References	Study Quality
10 (12591682)	2
2 (16545667)	4
12 (6359269)	4
11 (18201915)	3

US duplex Doppler lower extremity	Usually appropriate	Strong	0 0 mSv	0 0 mSv [ped]	7	7	1	0	0	0	1	2	8	0	2
-----------------------------------	---------------------	--------	---------	---------------	---	---	---	---	---	---	---	---	---	---	---

References	Study Quality
21 (21345629)	2
40 (8796687)	M
31 (10887229)	M
37 (2931880)	3
38 (2196267)	2
39 (8445746)	4
41 (8164280)	2
32 (16864668)	1
42 (30342218)	2
43 (32240007)	4

CTA lower extremity with IV contrast	May be appropriate	Strong	☼☼☼ 1-10 mSv	☼☼☼☼ 3-10 mSv [ped]	6	6	0	0	0	0	3	7	2	5	0
--------------------------------------	--------------------	--------	--------------	---------------------	---	---	---	---	---	---	---	---	---	---	---

References	Study Quality
21 (21345629)	2
23 (27399244)	4
15 (21658691)	2
22 (16966479)	4
18 (16580545)	2
10 (12591682)	2
19 (17848679)	M
20 (12601190)	1
13 (20609871)	4
17 (15128997)	2
16 (18094974)	2

MRA lower extremity without IV contrast	May be appropriate	Moderate	0 0 mSv	0 0 mSv [ped]	6	6	0	0	0	1	7	8	1	0	0
---	--------------------	----------	---------	---------------	---	---	---	---	---	---	---	---	---	---	---

References	Study Quality
30 (22109304)	2

36 (21098208)	4
33 (22173999)	4
35 (21274982)	4
29 (25079487)	3

MRA lower extremity without and with IV contrast	May be appropriate	Strong	0 0 mSv	0 0 mSv [ped]	6	6	0	0	0	0	4	8	4	1	0
--	--------------------	--------	---------	---------------	---	---	---	---	---	---	---	---	---	---	---

References	Study Quality
26 (23801421)	M
30 (22109304)	2
28 (25241315)	4
31 (10887229)	M
27 (12239057)	3
36 (21098208)	4
32 (16864668)	1
25 (9052573)	4
33 (22173999)	4
35 (21274982)	4
29 (25079487)	3

MRA abdomen and pelvis with bilateral lower extremity runoff without IV contrast	May be appropriate	Moderate	0 0 mSv	0 0 mSv [ped]	6	6	0	1	0	3	2	5	3	0	0
--	--------------------	----------	---------	---------------	---	---	---	---	---	---	---	---	---	---	---

References	Study Quality
30 (22109304)	2
36 (21098208)	4
33 (22173999)	4
35 (21274982)	4
29 (25079487)	3

CTA lower extremity without and with IV contrast	May be appropriate	Strong	☹☹☹ 1-10 mSv		6	6	0	0	0	0	1	10	2	2	1
--	--------------------	--------	--------------	--	---	---	---	---	---	---	---	----	---	---	---

References	Study Quality
21 (21345629)	2
23 (27399244)	4

15 (21658691)	2
22 (16966479)	4
18 (16580545)	2
10 (12591682)	2
19 (17848679)	M
20 (12601190)	1
13 (20609871)	4
17 (15128997)	2
16 (18094974)	2
24 (17709077)	4

## **Appendix Key**

A more complete discussion of the items presented below can be found by accessing the supporting documents at the designated hyperlinks.

**Appropriateness Category:** The panel's recommendation for a procedure based on the assessment of the risks and benefits of performing the procedure for the specified clinical scenario.

**SOE:** Strength of Evidence. The assessment of the amount and quality of evidence found in the peer reviewed medical literature for an appropriateness recommendation.

- **References:** The citation number and PMID for the reference(s) associated with the recommendation.
- **Study Quality:** The assessment of the quality of an individual reference based on the number of study quality elements described in the reference.

**RRL:** Relative Radiation Level. A population based assessment of the amount of radiation a typical patient may be exposed to during the specified procedure.

**Rating:** The final rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

**Median:** The median rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

**Final tabulations:** A histogram showing the number of panel members who rated the procedure as noted in the column heading (ie, 1, 2, 3, etc.).

Additional supporting documents about the AC methodology and processes can be found at [www.acr.org/ac](http://www.acr.org/ac).