American College of Radiology ACR Appropriateness Criteria®

Pretreatment Planning of Invasive Cancer of the Cervix

Variant 1: FIGO stage 1b1, tumor size <4 cm.

	Appropriateness	G07		D 1 DD1		3.5.31			F	inal '	Tabu	latio	ns		
Procedure	Category	SOE	Adults RRL	Peds RRL	Rating	Median	1	2	3	4	5	6	7	8	9
FDG-PET/CT whole body	Usually appropriate		���� 10-30 mSv		8	n/a	0	0	0	0	0	0	0	0	0
MRI pelvis without and with IV contrast	Usually appropriate		O 0 mSv	O 0 mSv [ped]	8	n/a	0	0	0	0	0	0	0	0	0
MRI pelvis without IV contrast	May be appropriate		O 0 mSv	O 0 mSv [ped]	6	n/a	0	0	0	0	0	0	0	0	0
CT abdomen and pelvis with IV contrast	May be appropriate		≎≎≎ 1-10 mSv	���� 3- 10 mSv [ped]	5	n/a	0	0	0	0	0	0	0	0	0
Radiography chest	May be appropriate		≎ <0.1 mSv		4	n/a	0	0	0	0	0	0	0	0	0
CT abdomen and pelvis without IV contrast	Usually not appropriate		≎≎≎ 1-10 mSv	���� 3- 10 mSv [ped]	2	n/a	0	0	0	0	0	0	0	0	0
US abdomen	Usually not appropriate		O 0 mSv	O 0 mSv [ped]	2	n/a	0	0	0	0	0	0	0	0	0
US pelvis transabdominal	Usually not appropriate		O 0 mSv	O 0 mSv [ped]	2	n/a	0	0	0	0	0	0	0	0	0
US pelvis transvaginal	Usually not appropriate		O 0 mSv	O 0 mSv [ped]	2	n/a	0	0	0	0	0	0	0	0	0

CT abdomen and pelvis without and with IV contrast	Usually not appropriate	୫୫୫୫ 10-30 mSv	≎≎≎≎≎ 10-30 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0
Bone scan whole body	Usually not appropriate	��� 1-10 mSv	���� 3- 10 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0
Fluoroscopy contrast enema	Usually not appropriate	≎≎≎ 1-10 mSv	���� 3- 10 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0
Radiography intravenous urography	Usually not appropriate	≎≎≎ 1-10 mSv	��� 0.3- 3 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0

Variant 2: FIGO stage Ib2, tumor size >4 cm.

Duo oo daano	Appropriateness	SOE	Adults RRL	Peds RRL	Dating	Median			F	inal T	Гаbu	lation	ıs		
Procedure	Category				Rating		1	2	3	4	5	6	7	8	9
FDG-PET/CT whole body	Usually appropriate		���� 10-30 mSv		9	n/a	0	0	0	0	0	0	0	0	0
MRI pelvis without and with IV contrast	Usually appropriate		O 0 mSv	O 0 mSv [ped]	9	n/a	0	0	0	0	0	0	0	0	0
MRI pelvis without IV contrast	May be appropriate		O 0 mSv	O 0 mSv [ped]	6	n/a	0	0	0	0	0	0	0	0	0
CT abdomen and pelvis with IV contrast	May be appropriate		≎≎≎ 1-10 mSv	���� 3- 10 mSv [ped]	5	n/a	0	0	0	0	0	0	0	0	0
Radiography chest	May be appropriate		⊕ <0.1 mSv		5	n/a	0	0	0	0	0	0	0	0	0
CT abdomen and pelvis without IV contrast	Usually not appropriate		≎≎≎ 1-10 mSv	���� 3- 10 mSv [ped]	2	n/a	0	0	0	0	0	0	0	0	0

US abdomen	Usually not appropriate	O 0 mSv	O 0 mSv [ped]	2	n/a	0	0	0	0	0	0	0	0	0
US pelvis transabdominal	Usually not appropriate	O 0 mSv	O 0 mSv [ped]	2	n/a	0	0	0	0	0	0	0	0	0
US pelvis transvaginal	Usually not appropriate	O 0 mSv	O 0 mSv [ped]	2	n/a	0	0	0	0	0	0	0	0	0
CT abdomen and pelvis without and with IV contrast	Usually not appropriate	୫୫୫୫ 10-30 mSv	����� 10-30 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0
Bone scan whole body	Usually not appropriate	��� 1-10 mSv	���� 3- 10 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0
Fluoroscopy contrast enema	Usually not appropriate	≎≎≎ 1-10 mSv	���� 3- 10 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0
Radiography intravenous urography	Usually not appropriate	≎≎≎ 1-10 mSv	��� 0.3- 3 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0

Variant 3: FIGO stage >Ib.

Procedure	Appropriateness	COF	A J14- DDI	RRL Peds RRL	Rating	Median	. Final Tabulations												
	Category	SOE	Adults RRL				1	2	3	4	5	6	7	8	9				
FDG-PET/CT whole body	Usually appropriate		���⊕ 10-30 mSv		9	n/a	0	0	0	0	0	0	0	0	0				
MRI pelvis without and with IV contrast	Usually appropriate		O 0 mSv	O 0 mSv [ped]	9	n/a	0	0	0	0	0	0	0	0	0				
CT abdomen and pelvis with IV contrast	Usually appropriate		≎≎≎ 1-10 mSv	���� 3- 10 mSv [ped]	7	n/a	0	0	0	0	0	0	0	0	0				
CT chest with IV contrast	May be appropriate		≎≎≎ 1-10 mSv	���� 3- 10 mSv [ped]	6	n/a	0	0	0	0	0	0	0	0	0				

MRI pelvis without IV contrast	May be appropriate	O 0 mSv	O 0 mSv [ped]	6	n/a	0	0	0	0	0	0	0	0	0
CT chest without IV contrast	May be appropriate	≎≎≎ 1-10 mSv	���� 3- 10 mSv [ped]	4	n/a	0	0	0	0	0	0	0	0	0
CT abdomen and pelvis without IV contrast	Usually not appropriate	≎≎≎ 1-10 mSv	���� 3- 10 mSv [ped]	2	n/a	0	0	0	0	0	0	0	0	0
Bone scan whole body	Usually not appropriate	��� 1-10 mSv	���� 3- 10 mSv [ped]	2	n/a	0	0	0	0	0	0	0	0	0
US abdomen	Usually not appropriate	O 0 mSv	O 0 mSv [ped]	2	n/a	0	0	0	0	0	0	0	0	0
US pelvis transabdominal	Usually not appropriate	O 0 mSv	O 0 mSv [ped]	2	n/a	0	0	0	0	0	0	0	0	0
US pelvis transvaginal	Usually not appropriate	O 0 mSv	O 0 mSv [ped]	2	n/a	0	0	0	0	0	0	0	0	0
Radiography chest	Usually not appropriate	≎ <0.1 mSv		2	n/a	0	0	0	0	0	0	0	0	0
CT abdomen and pelvis without and with IV contrast	Usually not appropriate	୫୫୫୫ 10-30 mSv	≎≎≎≎≎ 10-30 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0
CT chest without and with IV contrast	Usually not appropriate	��� 1-10 mSv	���� 3- 10 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0
Fluoroscopy contrast enema	Usually not appropriate	��� 1-10 mSv	���� 3- 10 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0
Radiography intravenous urography	Usually not appropriate	��� 1-10 mSv	��� 0.3- 3 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0

Appendix Key

A more complete discussion of the items presented below can be found by accessing the supporting documents at the designated hyperlinks.

Appropriateness Category: The panel's recommendation for a procedure based on the assessment of the risks and benefits of performing the procedure for the specified clinical scenario.

SOE: Strength of Evidence. The assessment of the amount and quality of evidence found in the peer reviewed medical literature for an appropriateness recommendation.

- **References:** The citation number and PMID for the reference(s) associated with the recommendation.
- Study Quality: The assessment of the quality of an individual reference based on the number of study quality elements described in the reference.

RRL: Relative Radiation Level. A population based assessment of the amount of radiation a typical patient may be exposed to during the specified procedure.

Rating: The final rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

Median: The median rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

Final tabulations: A histogram showing the number of panel members who rated the procedure as noted in the column heading (ie, 1, 2, 3, etc.).

Additional supporting documents about the AC methodology and processes can be found at www.acr.org/ac.