

**American College of Radiology
ACR Appropriateness Criteria®**

Sinonasal Disease

Variant 1: Acute (<4 weeks) uncomplicated rhinosinusitis.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
CT paranasal sinuses without IV contrast	May be appropriate		☼☼ 0.1-1mSv	☼☼☼ 0.3-3 mSv [ped]	4	n/a	0	0	0	0	0	0	0	0	0
CT cone beam paranasal sinuses without IV contrast	May be appropriate		☼☼ 0.1-1mSv		4	n/a	0	0	0	0	0	0	0	0	0
CT paranasal sinuses with IV contrast	Usually not appropriate		☼☼ 0.1-1mSv	☼☼☼ 0.3-3 mSv [ped]	2	n/a	0	0	0	0	0	0	0	0	0
Radiography paranasal sinuses	Usually not appropriate		☼ <0.1 mSv	☼ <0.03 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0
CT paranasal sinuses without and with IV contrast	Usually not appropriate		☼☼☼ 1-10 mSv	☼☼☼☼ 3-10 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0
MRI maxillofacial without IV contrast	Usually not appropriate		○ 0 mSv	○ 0 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0
MRI maxillofacial without and with IV contrast	Usually not appropriate		○ 0 mSv	○ 0 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0

Variant 2: Possible surgical candidate. Recurrent acute rhinosinusitis, chronic rhinosinusitis, sinonasal polyposis, or noninvasive fungal sinusitis.

CT paranasal sinuses with IV contrast	Usually appropriate		☼☼ 0.1-1mSv	☼☼☼ 0.3-3 mSv [ped]	7	n/a	0	0	0	0	0	0	0	0	0
MRI maxillofacial without IV contrast	May be appropriate		○ 0 mSv	○ 0 mSv [ped]	6	n/a	0	0	0	0	0	0	0	0	0
Arteriography craniofacial	May be appropriate		☼☼☼ 1-10 mSv		4	n/a	0	0	0	0	0	0	0	0	0
CTA head with IV contrast	May be appropriate		☼☼☼ 1-10 mSv	☼☼☼☼ 3-10 mSv [ped]	4	n/a	0	0	0	0	0	0	0	0	0
MRA head without IV contrast	May be appropriate		○ 0 mSv	○ 0 mSv [ped]	4	n/a	0	0	0	0	0	0	0	0	0
CTA neck with IV contrast	May be appropriate		☼☼☼ 1-10 mSv	☼☼☼ 0.3-3 mSv [ped]	4	n/a	0	0	0	0	0	0	0	0	0
FDG-PET/CT skull base to mid-thigh	May be appropriate		☼☼☼☼ 10-30 mSv	☼☼☼☼☼ 10-30 mSv [ped]	4	n/a	0	0	0	0	0	0	0	0	0
MRA head with IV contrast	May be appropriate		○ 0 mSv	○ 0 mSv [ped]	4	n/a	0	0	0	0	0	0	0	0	0
MRA neck with IV contrast	May be appropriate		○ 0 mSv	○ 0 mSv [ped]	4	n/a	0	0	0	0	0	0	0	0	0
MRA neck without and with IV contrast	Usually not appropriate		○ 0 mSv	○ 0 mSv [ped]	3	n/a	0	0	0	0	0	0	0	0	0
CT paranasal sinuses without and with IV contrast	Usually not appropriate		☼☼☼ 1-10 mSv	☼☼☼☼ 3-10 mSv [ped]	2	n/a	0	0	0	0	0	0	0	0	0
CT cone beam paranasal sinuses without IV contrast	Usually not appropriate		☼☼ 0.1-1mSv		2	n/a	0	0	0	0	0	0	0	0	0

Variant 5: Suspected invasive fungal sinusitis.

Appendix Key

A more complete discussion of the items presented below can be found by accessing the supporting documents at the designated hyperlinks.

Appropriateness Category: The panel's recommendation for a procedure based on the assessment of the risks and benefits of performing the procedure for the specified clinical scenario.

SOE: Strength of Evidence. The assessment of the amount and quality of evidence found in the peer reviewed medical literature for an appropriateness recommendation.

- **References:** The citation number and PMID for the reference(s) associated with the recommendation.
- **Study Quality:** The assessment of the quality of an individual reference based on the number of study quality elements described in the reference.

RRL: Relative Radiation Level. A population based assessment of the amount of radiation a typical patient may be exposed to during the specified procedure.

Rating: The final rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

Median: The median rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

Final tabulations: A histogram showing the number of panel members who rated the procedure as noted in the column heading (ie, 1, 2, 3, etc.).

Additional supporting documents about the AC methodology and processes can be found at www.acr.org/ac.