



US abdomen	May be appropriate	Limited	0 0 mSv	0 0 mSv [ped]	6	6	0	0	0	0	0	0	0	0	0	0	
		References	Study Quality														
		44 (27203288)	3														
		43 (25771443)	3														
		42 (25475515)	3														
		40 (24768483)	3														
		41 (22360013)	3														
MRI abdomen and pelvis without and with IV contrast	May be appropriate	Limited	0 0 mSv	0 0 mSv [ped]	5	5	0	0	0	0	0	0	0	0	0	0	
		References	Study Quality														
		51 (22033948)	2														
US pelvis	May be appropriate	Limited	0 0 mSv	0 0 mSv [ped]	5	5	0	0	0	0	0	0	0	0	0	0	
		References	Study Quality														
		47 (25499245)	3														
		48 (22421290)	3														
		46 (22273324)	3														
MRI abdomen and pelvis without IV contrast	May be appropriate	Strong	0 0 mSv	0 0 mSv [ped]	4	4	0	0	0	0	0	0	0	0	0	0	
		References	Study Quality														
		55 (26901006)	Good														
		49 (26807893)	2														
		50 (26526902)	3														
		54 (24272981)	2														
		52 (24168926)	2														
		53 (23481162)	2														
CT abdomen and pelvis without and with IV contrast	Usually not appropriate	Strong	☼☼☼☼ 10-30 mSv	☼☼☼☼☼ 10-30 mSv [ped]	3	3	3	1	7	2	3	1	0	0	0		
		References	Study Quality														





Radiography abdomen	Usually not appropriate	Limited	☹☹ 0.1-1mSv	☹☹☹ 0.03-0.3 mSv [ped]	3	3	2	6	4	2	2	0	0	1	0
		References		Study Quality											
		56 (22574093)		3											
Fluoroscopy contrast enema	Usually not appropriate	Expert Consensus	☹☹☹ 1-10 mSv	☹☹☹☹ 3-10 mSv [ped]	2	2	0	0	0	0	0	0	0	0	0
WBC scan abdomen and pelvis	Usually not appropriate	Limited	☹☹☹☹ 10-30 mSv		2	2	0	0	0	0	0	0	0	0	0
		References		Study Quality											
		57 (1456603)		3											

**Variant 3: Pregnant woman. Right lower quadrant pain, fever, leukocytosis. Suspected appendicitis. Initial imaging.**

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
US abdomen	Usually appropriate	Limited	○ 0 mSv	○ 0 mSv [ped]	8	8	0	0	0	0	0	0	0	0	0
		References		Study Quality											
		63 (27554180)		3											
		62 (24151869)		3											
		64 (22370694)		3											
MRI abdomen and pelvis without IV contrast	Usually appropriate	Limited	○ 0 mSv	○ 0 mSv [ped]	7	7	0	0	0	0	0	0	0	0	0
		References		Study Quality											
		67 (26338256)		3											
		65 (26215327)		3											
		66 (25291255)		3											
		61 (25148766)		3											
		68 (23360736)		3											

US pelvis	May be appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	6	6	0	0	0	0	0	0	0	0	0
CT abdomen and pelvis with IV contrast	May be appropriate	Limited	☹☹☹ 1-10 mSv	☹☹☹☹ 3-10 mSv [ped]	5	5	0	0	0	0	0	0	0	0	0
		References	Study Quality												
		61 (25148766)	3												
		58 (25419588)	3												
CT abdomen and pelvis without IV contrast	May be appropriate	Limited	☹☹☹ 1-10 mSv	☹☹☹☹ 3-10 mSv [ped]	4	4	0	0	0	0	0	0	0	0	0
		References	Study Quality												
		61 (25148766)	3												
		58 (25419588)	3												
CT abdomen and pelvis without and with IV contrast	Usually not appropriate	Limited	☹☹☹☹ 10-30 mSv	☹☹☹☹☹ 10-30 mSv [ped]	3	3	0	0	0	0	0	0	0	0	0
		References	Study Quality												
		61 (25148766)	3												
		58 (25419588)	3												
Fluoroscopy contrast enema	Usually not appropriate	Expert Consensus	☹☹☹ 1-10 mSv	☹☹☹☹ 3-10 mSv [ped]	2	2	0	0	0	0	0	0	0	0	0
MRI abdomen and pelvis without and with IV contrast	Usually not appropriate	Limited	0 0 mSv	0 0 mSv [ped]	2	2	0	0	0	0	0	0	0	0	0
		References	Study Quality												
		67 (26338256)	3												
		65 (26215327)	3												
		66 (25291255)	3												
		61 (25148766)	3												
		68 (23360736)	3												



## **Appendix Key**

A more complete discussion of the items presented below can be found by accessing the supporting documents at the designated hyperlinks.

**Appropriateness Category:** The panel's recommendation for a procedure based on the assessment of the risks and benefits of performing the procedure for the specified clinical scenario.

**SOE:** Strength of Evidence. The assessment of the amount and quality of evidence found in the peer reviewed medical literature for an appropriateness recommendation.

- **References:** The citation number and PMID for the reference(s) associated with the recommendation.
- **Study Quality:** The assessment of the quality of an individual reference based on the number of study quality elements described in the reference.

**RRL:** Relative Radiation Level. A population based assessment of the amount of radiation a typical patient may be exposed to during the specified procedure.

**Rating:** The final rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

**Median:** The median rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

**Final tabulations:** A histogram showing the number of panel members who rated the procedure as noted in the column heading (ie, 1, 2, 3, etc.).

Additional supporting documents about the AC methodology and processes can be found at [www.acr.org/ac](http://www.acr.org/ac).