American College of Radiology
ACR Appropriateness Criteria®

Chronic Chest Pain—High Probability of Coronary Artery Disease

Variant 1: Chronic chest pain; high probability of coronary artery disease.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
Radiography chest	Usually appropriate		☢ <0.1 mSv	☢ <0.03 mSv [ped]	9	n/a	0	0	0	0	0	0	0	0	0
Tc-99m SPECT MPI rest and stress	Usually appropriate		☢☢☢☢ 10-30 mSv		9	n/a	0	0	0	0	0	0	0	0	0
MRI heart with function and vasodilator stress perfusion without and with IV contrast	Usually appropriate		O 0 mSv	O 0 mSv [ped]	9	n/a	0	0	0	0	0	0	0	0	0
Arteriography coronary	Usually appropriate		☢☢☢ 1-10 mSv		9	n/a	0	0	0	0	0	0	0	0	0
CTA coronary arteries with IV contrast	Usually appropriate		☢☢☢ 1-10 mSv		8	n/a	0	0	0	0	0	0	0	0	0
US echocardiography transthoracic stress	Usually appropriate		O 0 mSv	O 0 mSv [ped]	8	n/a	0	0	0	0	0	0	0	0	0
Rb-82 PET heart stress	Usually appropriate		☢☢☢ 1-10 mSv		8	n/a	0	0	0	0	0	0	0	0	0
MRI heart function and morphology without and with IV contrast	Usually appropriate		O 0 mSv	O 0 mSv [ped]	7	n/a	0	0	0	0	0	0	0	0	0
MRI heart with function and inotropic stress without IV contrast	Usually appropriate		O 0 mSv	O 0 mSv [ped]	7	n/a	0	0	0	0	0	0	0	0	0

MRI heart with function and inotropic stress without and with IV contrast	Usually appropriate		O 0 mSv	O 0 mSv [ped]	7	n/a	0	0	0	0	0	0	0	0	0
MRA coronary arteries without and with IV contrast	May be appropriate		O 0 mSv	O 0 mSv [ped]	5	n/a	0	0	0	0	0	0	0	0	0
MRI heart function and morphology without IV contrast	May be appropriate		O 0 mSv	O 0 mSv [ped]	4	n/a	0	0	0	0	0	0	0	0	0
US echocardiography transthoracic resting	May be appropriate		O 0 mSv	O 0 mSv [ped]	4	n/a	0	0	0	0	0	0	0	0	0
CT chest with IV contrast	May be appropriate		☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	4	n/a	0	0	0	0	0	0	0	0	0
MRA coronary arteries without IV contrast	May be appropriate		O 0 mSv	O 0 mSv [ped]	4	n/a	0	0	0	0	0	0	0	0	0
CT chest without IV contrast	May be appropriate		☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	4	n/a	0	0	0	0	0	0	0	0	0
CT chest without and with IV contrast	May be appropriate		☢☢☢ 1-10 mSv	☢☢☢☢ 3-10 mSv [ped]	4	n/a	0	0	0	0	0	0	0	0	0
CT coronary calcium	Usually not appropriate		☢☢☢ 1-10 mSv		3	n/a	0	0	0	0	0	0	0	0	0
Tc-99m ventriculography	Usually not appropriate		☢☢☢ 1-10 mSv		3	n/a	0	0	0	0	0	0	0	0	0
US abdomen	Usually not appropriate		O 0 mSv	O 0 mSv [ped]	3	n/a	0	0	0	0	0	0	0	0	0

Appendix Key

A more complete discussion of the items presented below can be found by accessing the supporting documents at the designated hyperlinks.

Appropriateness Category: The panel's recommendation for a procedure based on the assessment of the risks and benefits of performing the procedure for the specified clinical scenario.

SOE: Strength of Evidence. The assessment of the amount and quality of evidence found in the peer reviewed medical literature for an appropriateness recommendation.

- **References:** The citation number and PMID for the reference(s) associated with the recommendation.
- **Study Quality**: The assessment of the quality of an individual reference based on the number of study quality elements described in the reference.

RRL: Relative Radiation Level. A population based assessment of the amount of radiation a typical patient may be exposed to during the specified procedure.

Rating: The final rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

Median: The median rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

Final tabulations: A histogram showing the number of panel members who rated the procedure as noted in the column heading (ie, 1, 2, 3, etc.).

Additional supporting documents about the AC methodology and processes can be found at **www.acr.org/ac.**