

CT knee without IV contrast	Usually not appropriate	Expert Consensus	⊕ <0.1 mSv	⊕⊕ 0.03-0.3 mSv [ped]	1	1	10	4	1	1	1	0	0	0	0
CT knee without and with IV contrast	Usually not appropriate	Expert Consensus	⊕ <0.1 mSv	⊕⊕ 0.03-0.3 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0
MRA knee without IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	16	1	0	0	0	0	0	0	0
MRA knee without and with IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0
MR arthrography knee	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0
MRI knee without IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	13	0	1	1	1	0	0	1	0
MRI knee without and with IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0
US knee	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	11	4	0	0	1	1	0	0	0
Bone scan with SPECT or SPECT/CT knee	Usually not appropriate	Expert Consensus	⊕⊕⊕ 1-10 mSv	⊕⊕⊕⊕ 3-10 mSv [ped]	1	1	16	1	0	0	0	0	0	0	0

Variant 3: Adult or skeletally mature child. Fall or acute twisting trauma to the knee. No fracture seen on radiographs. Suspect occult fracture or internal derangement. Next study.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
MRI knee without IV contrast	Usually appropriate	Strong	○ 0 mSv	○ 0 mSv [ped]	9	9	0	0	0	0	0	0	0	3	14
			References	Study Quality											
			23 (16118171)	1											
			26 (16861540)	3											

21 (23677359)	2
22 (17394470)	4
24 (18795300)	4
20 (23773879)	4
30 (21380741)	2
35 (17483932)	2
36 (12324825)	3
19 (-3093387)	4
25 (29752650)	4
34 (28680889)	3
32 (28432395)	4
33 (28424850)	4
29 (27775896)	3
31 (28293698)	4
27 (27097792)	4
28 (27177438)	3

CT knee without IV contrast	May be appropriate	Strong	☼ <0.1 mSv	☼☼ 0.03-0.3 mSv [ped]	6	6	0	0	0	0	3	6	2	1	5
-----------------------------	--------------------	--------	------------	-----------------------	---	---	---	---	---	---	---	---	---	---	---

References	Study Quality
14 (26025120)	3
17 (20551186)	1
12 (16392612)	3
13 (17136560)	3
16 (23020888)	2
15 (28214478)	4
18 (25816992)	2

CT knee with IV contrast	Usually not appropriate	Strong	☼ <0.1 mSv	☼☼ 0.03-0.3 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0
--------------------------	-------------------------	--------	------------	-----------------------	---	---	----	---	---	---	---	---	---	---	---

References	Study Quality
14 (26025120)	3

17 (20551186)	1
12 (16392612)	3
13 (17136560)	3
16 (23020888)	2
15 (28214478)	4
18 (25816992)	2

CT knee without and with IV contrast	Usually not appropriate	Strong	☼ <0.1 mSv	☼☼☼ 0.03-0.3 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0
--------------------------------------	-------------------------	--------	------------	------------------------	---	---	----	---	---	---	---	---	---	---	---

References	Study Quality
14 (26025120)	3
17 (20551186)	1
12 (16392612)	3
13 (17136560)	3
16 (23020888)	2
15 (28214478)	4
18 (25816992)	2

MRA knee without IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0
MRA knee without and with IV contrast	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0
MR arthrography knee	Usually not appropriate	Expert Consensus	○ 0 mSv	○ 0 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0
MRI knee without and with IV contrast	Usually not appropriate	Strong	○ 0 mSv	○ 0 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0

References	Study Quality
23 (16118171)	1
26 (16861540)	3
21 (23677359)	2
22 (17394470)	4
24 (18795300)	4

20 (23773879)	4
30 (21380741)	2
35 (17483932)	2
36 (12324825)	3
19 (-3093387)	4
25 (29752650)	4
34 (28680889)	3
32 (28432395)	4
33 (28424850)	4
29 (27775896)	3
31 (28293698)	4
27 (27097792)	4
28 (27177438)	3

US knee	Usually not appropriate	Strong	0 0 mSv	0 0 mSv [ped]	1	1	13	2	1	0	0	1	0	0	0
---------	-------------------------	--------	---------	---------------	---	---	----	---	---	---	---	---	---	---	---

References	Study Quality
40 (16786321)	2
43 (17678659)	3
44 (21982460)	3
41 (28374066)	4
47 (29466145)	Good
49 (27145775)	4
48 (28343806)	4
46 (25916746)	Good
42 (25911713)	3
45 (23412237)	2

Bone scan with SPECT or SPECT/CT knee	Usually not appropriate	Limited	☼☼☼ 1-10 mSv	☼☼☼☼ 3-10 mSv [ped]	1	1	14	2	0	0	0	0	0	0	1
---------------------------------------	-------------------------	---------	--------------	---------------------	---	---	----	---	---	---	---	---	---	---	---

References	Study Quality
38 (16894322)	2
37 (12218448)	3

39 (25558700)	3
---------------	---

Variant 4: Skeletally immature child. Fall or acute twisting trauma to the knee. No fracture seen on radiographs. Suspect occult fracture or internal derangement. Next study.

Procedure	Appropriateness Category	SOE	Adults RRL	Peds RRL	Rating	Median	Final Tabulations								
							1	2	3	4	5	6	7	8	9
MRI knee without IV contrast	Usually appropriate	Strong	○ 0 mSv	○ 0 mSv [ped]	9	9	1	0	0	0	0	0	0	1	15
		References		Study Quality											
		26 (16861540)		3											
		21 (23677359)		2											
		34 (28680889)		3											
		32 (28432395)		4											
		33 (28424850)		4											
		29 (27775896)		3											
		31 (28293698)		4											
		27 (27097792)		4											
		28 (27177438)		3											
CT knee without IV contrast	May be appropriate	Strong	⊕ <0.1 mSv	⊕⊕ 0.03-0.3 mSv [ped]	5	5	1	0	0	2	12	1	1	0	0
		References		Study Quality											
		14 (26025120)		3											
		17 (20551186)		1											
		26 (16861540)		3											
		21 (23677359)		2											
		12 (16392612)		3											
		50 (22965381)		4											
		16 (23020888)		2											
		51 (28190582)		3											
		34 (28680889)		3											

References	Study Quality
14 (26025120)	3
17 (20551186)	1
26 (16861540)	3
21 (23677359)	2
12 (16392612)	3
50 (22965381)	4
16 (23020888)	2
51 (28190582)	3
34 (28680889)	3
32 (28432395)	4
33 (28424850)	4
29 (27775896)	3
31 (28293698)	4
15 (28214478)	4
27 (27097792)	4
28 (27177438)	3
18 (25816992)	2

MRA knee without IV contrast	Usually not appropriate	Expert Consensus	0 mSv	0 mSv [ped]	1	1	16	0	0	0	0	0	0	0	0	1
MRA knee without and with IV contrast	Usually not appropriate	Expert Consensus	0 mSv	0 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0	0
MRI knee without and with IV contrast	Usually not appropriate	Strong	0 mSv	0 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0	0

References	Study Quality
26 (16861540)	3
21 (23677359)	2
34 (28680889)	3
32 (28432395)	4
33 (28424850)	4
29 (27775896)	3

26 (16861540)	3
21 (23677359)	2
34 (28680889)	3
32 (28432395)	4
33 (28424850)	4
29 (27775896)	3
31 (28293698)	4
27 (27097792)	4
28 (27177438)	3

CT knee without IV contrast	May be appropriate	Strong	☼ <0.1 mSv	☼☼ 0.03-0.3 mSv [ped]	5	8	0	0	0	1	0	2	5	1	8
-----------------------------	--------------------	--------	------------	-----------------------	---	---	---	---	---	---	---	---	---	---	---

References	Study Quality
14 (26025120)	3
17 (20551186)	1
26 (16861540)	3
21 (23677359)	2
12 (16392612)	3
50 (22965381)	4
16 (23020888)	2
51 (28190582)	3
34 (28680889)	3
32 (28432395)	4
33 (28424850)	4
29 (27775896)	3
31 (28293698)	4
15 (28214478)	4
27 (27097792)	4
28 (27177438)	3
18 (25816992)	2

CT knee with IV contrast	Usually not appropriate	Strong	☼ <0.1 mSv	☼☼ 0.03-0.3 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0
--------------------------	-------------------------	--------	------------	-----------------------	---	---	----	---	---	---	---	---	---	---	---

References	Study Quality
14 (26025120)	3
17 (20551186)	1
26 (16861540)	3
21 (23677359)	2
12 (16392612)	3
50 (22965381)	4
16 (23020888)	2
51 (28190582)	3
34 (28680889)	3
32 (28432395)	4
33 (28424850)	4
29 (27775896)	3
31 (28293698)	4
15 (28214478)	4
27 (27097792)	4
28 (27177438)	3
18 (25816992)	2

CT knee without and with IV contrast	Usually not appropriate	Strong	☼ <0.1 mSv	☼☼ 0.03-0.3 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0
--------------------------------------	-------------------------	--------	------------	-----------------------	---	---	----	---	---	---	---	---	---	---	---

References	Study Quality
14 (26025120)	3
17 (20551186)	1
26 (16861540)	3
21 (23677359)	2
12 (16392612)	3
50 (22965381)	4
16 (23020888)	2

51 (28190582)	3
34 (28680889)	3
32 (28432395)	4
33 (28424850)	4
29 (27775896)	3
31 (28293698)	4
15 (28214478)	4
27 (27097792)	4
28 (27177438)	3
18 (25816992)	2

MRA knee without IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0
MRA knee without and with IV contrast	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0
MRI knee without and with IV contrast	Usually not appropriate	Strong	0 0 mSv	0 0 mSv [ped]	1	1	16	0	0	0	0	1	0	0	0

References	Study Quality
26 (16861540)	3
21 (23677359)	2
34 (28680889)	3
32 (28432395)	4
33 (28424850)	4
29 (27775896)	3
31 (28293698)	4
27 (27097792)	4
28 (27177438)	3

Bone scan with SPECT or SPECT/CT lower extremity	Usually not appropriate	Limited	☹☹☹ 1-10 mSv	☹☹☹☹ 3-10 mSv [ped]	1	n/a	0	0	0	0	0	0	0	0	0
--	-------------------------	---------	--------------	---------------------	---	-----	---	---	---	---	---	---	---	---	---

References	Study Quality
38 (16894322)	2

58 (21156317)	4
59 (17869471)	4
60 (16498092)	2

CT knee with IV contrast	May be appropriate (Disagreement)	Expert Opinion	☼ <0.1 mSv	☼☼ 0.03-0.3 mSv [ped]	5	5	9	0	2	1	3	0	1	1	0
--------------------------	-----------------------------------	----------------	------------	-----------------------	---	---	---	---	---	---	---	---	---	---	---

References	Study Quality
12 (16392612)	3

CT knee without IV contrast	May be appropriate	Limited	☼ <0.1 mSv	☼☼ 0.03-0.3 mSv [ped]	5	5	3	3	1	1	7	2	0	0	0
-----------------------------	--------------------	---------	------------	-----------------------	---	---	---	---	---	---	---	---	---	---	---

References	Study Quality
12 (16392612)	3

Arteriography lower extremity	May be appropriate	Limited	☼☼ 0.1-1mSv	☼☼☼ 0.3-3 mSv [ped]	5	5	0	1	2	1	11	1	1	0	0
-------------------------------	--------------------	---------	-------------	---------------------	---	---	---	---	---	---	----	---	---	---	---

References	Study Quality
58 (21156317)	4
56 (19932064)	4
57 (19074408)	4

MRA knee without and with IV contrast	May be appropriate	Limited	○ 0 mSv	○ 0 mSv [ped]	5	5	1	1	1	2	10	2	0	0	0
---------------------------------------	--------------------	---------	---------	---------------	---	---	---	---	---	---	----	---	---	---	---

References	Study Quality
61 (11972076)	4
62 (21446625)	3

MRI knee without IV contrast	May be appropriate	Strong	○ 0 mSv	○ 0 mSv [ped]	5	5	2	0	1	1	12	1	0	0	0
------------------------------	--------------------	--------	---------	---------------	---	---	---	---	---	---	----	---	---	---	---

References	Study Quality
26 (16861540)	3
21 (23677359)	2
62 (21446625)	3
32 (28432395)	4
29 (27775896)	3

			31 (28293698)		4													
			27 (27097792)		4													
			28 (27177438)		3													
MRA knee without IV contrast	Usually not appropriate	Limited	0 0 mSv	0 0 mSv [ped]	3	3	7	0	3	1	4	1	0	0	1			
			References		Study Quality													
			61 (11972076)		4													
			62 (21446625)		3													
CT knee without and with IV contrast	Usually not appropriate	Limited	☼ <0.1 mSv	☼☼ 0.03-0.3 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0	0	0	0
			References		Study Quality													
			12 (16392612)		3													
MR arthrography knee	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	1	1	17	0	0	0	0	0	0	0	0	0	0	0
MRI knee without and with IV contrast	Usually not appropriate	Strong	0 0 mSv	0 0 mSv [ped]	1	1	16	0	0	0	1	0	0	0	0	0	0	0
			References		Study Quality													
			26 (16861540)		3													
			21 (23677359)		2													
			62 (21446625)		3													
			32 (28432395)		4													
			29 (27775896)		3													
			31 (28293698)		4													
			27 (27097792)		4													
			28 (27177438)		3													
US knee	Usually not appropriate	Expert Consensus	0 0 mSv	0 0 mSv [ped]	1	1	14	1	0	0	0	2	0	0	0			
Bone scan with SPECT or SPECT/CT knee	Usually not appropriate	Expert Consensus	☼☼☼ 1-10 mSv	☼☼☼☼ 3-10 mSv [ped]	1	1	16	1	0	0	0	0	0	0	0			

Appendix Key

A more complete discussion of the items presented below can be found by accessing the supporting documents at the designated hyperlinks.

Appropriateness Category: The panel's recommendation for a procedure based on the assessment of the risks and benefits of performing the procedure for the specified clinical scenario.

SOE: Strength of Evidence. The assessment of the amount and quality of evidence found in the peer reviewed medical literature for an appropriateness recommendation.

- **References:** The citation number and PMID for the reference(s) associated with the recommendation.
- **Study Quality:** The assessment of the quality of an individual reference based on the number of study quality elements described in the reference.

RRL: Relative Radiation Level. A population based assessment of the amount of radiation a typical patient may be exposed to during the specified procedure.

Rating: The final rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

Median: The median rating (1-9 scale) for the procedure as determined by the panel during rating rounds.

Final tabulations: A histogram showing the number of panel members who rated the procedure as noted in the column heading (ie, 1, 2, 3, etc.).

Additional supporting documents about the AC methodology and processes can be found at www.acr.org/ac.